

On q -asymptotics for linear q -difference-differential equations with Fuchsian and irregular singularities

Alberto Lastra, Stéphane Malek, Javier Sanz

November, 14 2011

Abstract

We consider a Cauchy problem for some family of linear q -difference-differential equations with Fuchsian and irregular singularities, that admit a unique formal power series solution in two variables $\hat{X}(t, z)$ for given formal power series initial conditions. Under suitable conditions and by the application of certain q -Borel and Laplace transforms (introduced by J.-P. Ramis and C. Zhang), we are able to deal with the small divisors phenomenon caused by the Fuchsian singularity, and to construct actual holomorphic solutions of the Cauchy problem whose q -asymptotic expansion in t , uniformly for z in the compact sets of \mathbb{C} , is $\hat{X}(t, z)$. The small divisors's effect is an increase in the order of q -exponential growth and the appearance of a power of the factorial in the corresponding q -Gevrey bounds in the asymptotics.

Key words: q -difference-differential equations, q -Laplace transform, formal power series solutions, q -Gevrey asymptotic expansions, small divisors, Fuchsian and irregular singularities.

2010 MSC: 34K25, 34M25, 34M30, 33E30.

1 Introduction

The second author has studied partial differential equations of the form

$$(1) \quad t^{2r_2} \partial_t^{r_2} (z \partial_z)^{r_1} \partial_z^S u(t, z) = F(t, z, \partial_t, \partial_z) u(t, z)$$

where S, r_1, r_2 are nonnegative integers and F is some differential operator with polynomial coefficients. These equations belong to a class of partial differential equations with both irregular singularity at $t = 0$ (in the sense of T. Mandai [21]) and Fuchsian singularity at $z = 0$. These kind of problems have been extensively studied in the literature, see for example [1, 4, 9, 11, 20, 30] for Fuchsian partial differential equations, and [6, 21, 22] for irregular singularities.

It is possible to construct formal power series solutions for the equation (1) of the form $\hat{u}(t, z) = \sum_{m \geq 0} \hat{u}_m(t) z^m / m!$, with coefficients in $\mathbb{C}[[t]]$, for given initial data

$$(2) \quad (\partial_z^j \hat{u})(t, 0) = \hat{u}_j(t) \in \mathbb{C}[[t]],$$

$0 \leq j \leq S - 1$, which are assumed to be 1-Borel summable with respect to t in some direction $d \in \mathbb{R}$.

In the case $r_1 = 0$, it was shown in [18] that the formal series $\hat{u}(t, z)$ is 1-Borel summable with respect to t in the direction d if d is well chosen, as series with coefficients in the Banach space of holomorphic functions near the origin with the supremum norm.

In the paper [15] the case $r_1 \neq 0$ was treated, noticing that the formal series \hat{u} is in general no longer 1–Borel summable, but one can show the existence of actual holomorphic solutions $u(t, z)$ which are Gevrey asymptotic of order larger than 1 to $\hat{u}(t, z)$ with respect to t in sectors centered at 0 with finite radius in well chosen directions $d \in \mathbb{R}$. The reason for this different behaviour is the presence of small divisors introduced by the Fuchsian operator $(z\partial_z)^{r_1}$. More precisely, the singularities of the 1–Borel transforms $\mathbb{B}\hat{u}_m(\tau)$ accumulate to the origin in \mathbb{C} as m tends to infinity, so that the 1–Borel transform $(\mathbb{B}\hat{u})(\tau, z)$ with respect to t is only holomorphic on a sector with infinite radius centered at 0 with respect to τ . This induces a large multiplier effect on the l –th derivatives with respect to t of the actual solution of (1) constructed with the classical Borel-Laplace procedure $u(t, z) = (\mathbb{L}\mathbb{B}\hat{u})(t, z)$ which grows like $CK^l l! \Gamma(1 + \gamma l)$ for some $C, K > 0$ and some $\gamma > 2$ that can be expressed in terms of r_1, r_2 , for all $l \geq 0$.

In this paper, we study a q –analog of the problem (1), (2), discretized with respect to the variable t , where ∂_t is replaced by the operator $(f(qt) - f(t))/(qt - t)$ for $q \in \mathbb{C}$ (which formally tends to ∂_t as q tends to 1). Namely, we will consider the following linear q –difference-differential equation

$$(3) \quad ((z\partial_z + 1)^{r_1} (t\sigma_q)^{r_2} + 1) \partial_z^S \hat{X}(t, z) = \sum_{k=0}^{S-1} b_k(z) (t\sigma_q)^{m_{0,k}} (\partial_z^k \hat{X})(t, zq^{-m_{1,k}})$$

with given initial conditions

$$(4) \quad (\partial_z^j \hat{X})(t, 0) = \hat{X}_j(t) \in \mathbb{C}[[t]], \quad 0 \leq j \leq S - 1,$$

where $S, m_{0,k}, m_{1,k}$ are nonnegative integers, for $0 \leq k \leq S - 1$ and where $q \in \mathbb{C}$ such that $|q| > 1$, σ_q is the dilation operator defined by $(\sigma_q \hat{X})(t, z) = \hat{X}(qt, z)$, and $b_k(z)$ are polynomials in z . As in previous works [16], [17], the map $(t, z) \mapsto (q^{m_{0,k}} t, zq^{-m_{1,k}})$ is assumed to be a volume shrinking map, meaning that the modulus of the Jacobian determinant $|q|^{m_{0,k} - m_{1,k}}$ is less than 1. We will always assume that $r_2 \geq 1$, while $r_1 \geq 0$.

Advanced/delayed partial differential equations have also been widely studied, see for example [12, 13, 14, 23, 29, 32], and some authors have considered the use of special functions transforms for the study of the asymptotic properties of the solutions of q –difference-differential equations [10, 24]. Our present work is a contribution to this area.

It is not difficult to show (see Lemma 5) that this Cauchy problem has a unique formal power series solution of the form

$$\hat{X}(t, z) = \sum_{h \geq 0} \hat{X}_h(t) \frac{z^h}{h!},$$

where $\hat{X}_h(t) = \sum_{m \geq 0} f_{m,h} t^m \in \mathbb{C}[[t]]$, $h \geq 0$. Our purpose is to construct actual holomorphic solutions of this problem that are asymptotically represented by $\hat{X}(t, z)$ in a precise sense.

The key idea in our approach is the study of a related Cauchy problem,

$$(5) \quad ((z\partial_z + 1)^{r_1} \tau^{r_2} + 1) \partial_z^S \hat{W}(\tau, z) = \sum_{k=0}^{S-1} b_k(z) \tau^{m_{0,k}} (\partial_z^k \hat{W})(\tau, zq^{-m_{1,k}})$$

with initial conditions

$$(6) \quad (\partial_z^j \hat{W})(\tau, 0) = \hat{W}_j(\tau) \in \mathbb{C}[[\tau]], \quad 0 \leq j \leq S - 1,$$

which, by the application of a q –Laplace transform in the variable τ , provides information on our initial problem (see Lemma 6). The q –Laplace transform we consider was introduced by

J.-P. Ramis and C. Zhang in [28], and in recent years it has been used with great success in the study of the asymptotic properties of solutions of q -difference equations, see [8], in much the same way as the classical Laplace-Borel transform has been applied to the asymptotic study of formal solutions to differential equations and singular perturbation problems in the complex domain (see the works of W. Balser [2, 3], B. Malgrange [19], J.-P. Ramis [26] or O. Costin [7]). This new Cauchy problem (5), (6) is studied in two respects.

Firstly, assuming the initial conditions W_j are holomorphic and have q -exponential growth (of order 2) in a set $Vq^{\mathbb{Z}} = \{vq^h : v \in V, h \in \mathbb{Z}\}$, V being a well chosen bounded open set in $\mathbb{C} \setminus \{0\}$, and with some restriction on the argument of q , we prove in Theorem 2 that there exists a unique solution of (5), (6), of the form

$$(7) \quad W(\tau, z) = \sum_{h \geq 0} W_h(\tau) \frac{z^h}{h!},$$

holomorphic on $Vq^{\mathbb{Z}} \times \mathbb{C}$ and of q -exponential growth (of order 1) in τ , in the terminology of [28], uniformly for z in any compact set of \mathbb{C} . The increase in the order may be seen as an effect of the small divisors appearing in the problem.

Secondly, assuming the initial conditions W_j , $0 \leq j \leq S-1$, are holomorphic near the origin, we prove in Theorem 4 that the solution in (7) has coefficients W_h holomorphic in discs D_h whose radii tend to 0 as h tends to infinity, in such a way that there exist constants $C_1, T_1, X_1 > 0$ such that

$$\sup_{\tau \in \overline{D}_j} |\partial^n W_j(\tau)| \leq C_1 \left(\frac{1}{T_1}\right)^n \left(\frac{1}{X_1}\right)^j n! j! (j+1)^{\frac{r_1 n}{r_2}} |q|^{-j^2/2},$$

for every $n, j \geq 0$. The important fact here is the q -exponential decrease of these bounds with respect to j , what will turn out to be crucial in the following. These two results allow us to analyze the q -asymptotic expansion of the q -Laplace transforms of the W_h (Proposition 3), which is shown to hold in a common domain $\mathcal{T}_{\lambda, q, \delta, r_0}$ (see (9) for its definition) for all $h \geq 0$.

We are prepared to turn now to our main objective. Departing from formal initial conditions \hat{X}_j , $0 \leq j \leq S-1$, whose q -Borel transforms $W_j(\tau)$ (in the terminology of [28]) satisfy all the conditions in the previous two results, we are finally able to find a solution of our problem (Theorem 5) in the form

$$X(t, z) = \sum_{h \geq 0} \mathcal{L}_q^\lambda(W_h)(t) \frac{z^h}{h!},$$

which is holomorphic in $\mathcal{T}_{\lambda, q, \delta, r_0} \times \mathbb{C}$, and such that given $R > 0$, there exist constants $\tilde{C} > 0$, $\tilde{D} > 0$ such that for every $n \in \mathbb{N}$, $n \geq 1$, one has

$$\left| X(t, z) - \sum_{h \geq 0} \sum_{m=0}^{n-1} f_{m,h} t^m \frac{z^h}{h!} \right| \leq \tilde{C} \tilde{D}^n \Gamma\left(\frac{r_1}{r_2}(n+1)\right) |q|^{n(n-1)/2} |t|^n$$

for every $t \in \mathcal{T}_{\lambda, q, \delta, r_0}$, $z \in D(0, R)$. Again one may note that the small divisors phenomenon has caused the appearance of the term $\Gamma\left(\frac{r_1}{r_2}(n+1)\right)$.

The paper is organized as follows. Section 2 provides the facts concerning the q -Laplace transform. Section 3 is devoted to the study of a first auxiliary Cauchy problem in suitable weighted Banach spaces of formal Laurent series. This is needed in the following section, devoted to the proof of Theorem 2. A second Cauchy problem in weighted Banach spaces of formal Taylor series (Section 5) is applied in the next Section, which contains Theorem 4. Finally, Section 7

consists of the construction of the solution, and it also contains some remarks on the nature of the solution in the special case that $r_1 = 0$, in which no small divisors appear.

We fix some conventions. \mathbb{C}^* stands for $\mathbb{C} \setminus \{0\}$, and \mathbb{N} for the set $\{0, 1, 2, \dots\}$. $D(0, r)$ denotes the open disc with center 0 and radius $r > 0$. Given a set $V \subset \mathbb{C}$ and $q \in \mathbb{C}$, we define

$$Vq^{\mathbb{Z}} = \{vq^h : v \in V, h \in \mathbb{Z}\}, \quad Vq^{\mathbb{N}} = \{vq^h : v \in V, h \in \mathbb{N}\}.$$

2 A q -analogue of the Laplace transform and q -asymptotic expansion

In this section, we recall the definition of a q -analogue of the Laplace transform introduced in the papers [28, 31] and some of its properties that will be useful in the sequel. For the sake of clarity, we include the proof of these results (mainly available in [31]), since they contain important estimates that will be used in the proof of our main result (Theorem 5).

Proposition 1 *Let $q \in \mathbb{C}$ such that $|q| > 1$. Let V be an open and bounded set in \mathbb{C}^* and $D(0, \rho_0)$ a disc such that $V \cap D(0, \rho_0) \neq \emptyset$. Let $(\mathbb{F}, \|\cdot\|_{\mathbb{F}})$ be a complex Banach space. Let $\phi : Vq^{\mathbb{N}} \cup D(0, \rho_0) \rightarrow \mathbb{F}$ be a holomorphic function which satisfies the following estimates : there exist $C, M > 0$ such that*

$$(8) \quad \|\phi(xq^m)\|_{\mathbb{F}} \leq M|q|^{m^2/2}C^m$$

for all $m \geq 0$, all $x \in V$. Let Θ be the Jacobi Theta function defined in \mathbb{C}^* by

$$\Theta(x) = \sum_{n \in \mathbb{Z}} q^{-n(n-1)/2} x^n.$$

Let $\delta > 0$ and $\lambda \in V \cap D(0, \rho_0)$. We denote by

$$(9) \quad \mathcal{R}_{\lambda, q, \delta} = \{t \in \mathbb{C}^* : |1 + \frac{\lambda}{tq^k}| > \delta, \forall k \in \mathbb{Z}\}, \quad \mathcal{T}_{\lambda, q, \delta, r_1} = \mathcal{R}_{\lambda, q, \delta} \cap D(0, r_1).$$

The q -Laplace transform of ϕ in the direction $\lambda q^{\mathbb{Z}}$ is defined by

$$\mathcal{L}_q^\lambda(\phi)(t) := \sum_{m \in \mathbb{Z}} \phi(q^m \lambda) / \Theta\left(\frac{q^m \lambda}{t}\right)$$

for all $t \in \mathcal{T}_{\lambda, q, \delta, r_1}$, if $r_1 < |\lambda q^{1/2}|/C$. Moreover, $\mathcal{L}_q^\lambda(\phi)(t)$ defines a bounded holomorphic function on $\mathcal{T}_{\lambda, q, \delta, r_1}$ with values in \mathbb{F} when $r_1 < |\lambda q^{1/2}|/C$. Assume that the function ϕ has the following Taylor expansion

$$(10) \quad \phi(\tau) = \sum_{n \geq 0} \frac{f_n}{q^{n(n-1)/2}} \tau^n$$

on $D(0, \rho_0)$, where $f_n \in \mathbb{F}$, $n \geq 0$. Then, there exist two constants $D, B > 0$ such that

$$(11) \quad \|\mathcal{L}_q^\lambda(\phi)(t) - \sum_{m=0}^{n-1} f_m t^m\|_{\mathbb{F}} \leq DB^n |q|^{n(n-1)/2} |t|^n$$

for all $n \geq 1$, for all $t \in \mathcal{T}_{\lambda, q, \delta, r_1}$.

Remark: In the situation described by (11) it is said that $\mathcal{L}_q^\lambda(\phi)$ admits the series $\sum_{m=0}^{\infty} f_m t^m$ as q -Gevrey asymptotic expansion of order 1 (whenever the exponent of $|q|$ in the bounds is $n(n-1)/(2r)$ the order is said to be r). Analogously, a function that satisfies estimates such as (8) is said to have q -exponential growth of order 1 in $Vq^{\mathbb{N}}$.
If $\phi(z) = \sum_{n \geq 0} a_n z^n$ is an entire function such that there exists $C > 0$ such that

$$|a_n| \leq C \exp(-(n - \alpha)^2/2)$$

for all $n \geq 0$ and some $\alpha \geq 0$, then ϕ satisfies the estimates (8). For a reference, see [25].

Proof Since the Theta function $\Theta(x)$ satisfies the q -difference equation $\Theta(qx) = qx\Theta(x)$ for all $x \in \mathbb{C}^*$, we get that

$$(12) \quad \Theta\left(\frac{q^m \lambda}{t}\right) = q^{m(m+1)/2} \left(\frac{\lambda}{t}\right)^m \Theta\left(\frac{\lambda}{t}\right)$$

for all $t \in \mathbb{C}^*$. Moreover, from Lemma 4.6 of [27], there exists $K_1 > 0$ such that

$$(13) \quad |\Theta(q^m \lambda/t)| \geq K_1 \delta \sum_{n \in \mathbb{Z}} |q|^{-n(n-1)/2} \left|\frac{q^m \lambda}{t}\right|^n$$

for all $t \in \mathcal{R}_{\lambda, q, \delta}$, all $m \in \mathbb{Z}$.

In the proof, we will show the estimates (11). From them one may easily deduce that the series defining $\mathcal{L}_q^\lambda(\phi)(t)$ converges and defines a bounded holomorphic function on $\mathcal{T}_{\lambda, q, \delta, r_1}$. We would like to point out that many of the series following are initially formal, but we will finally prove their convergence.

Let $K \geq 0$ be an integer. First of all, we give estimates for the sum $\sum_{m > 0} \phi(q^m \lambda)/\Theta(q^m \lambda/t)$. From the estimates (13), we have that

$$(14) \quad \left|\Theta\left(\frac{\lambda}{t}\right)\right| \geq K_1 \delta |q|^{-K(K-1)/2} \left|\frac{\lambda}{t}\right|^K$$

for all $t \in \mathcal{R}_{\lambda, q, \delta}$. Using (8), (12) and (14), we get the estimates

$$\left\| \frac{\phi(q^m \lambda)}{\Theta(q^m \lambda/t)} \right\|_{\mathbb{F}} \leq \frac{M}{K_1 \delta} \left(\frac{1}{|\lambda|}\right)^K |q|^{K(K-1)/2} |t|^K \left(\frac{C|t|}{|\lambda||q|^{1/2}}\right)^m$$

for all $m > 0$, all $t \in \mathcal{R}_{\lambda, q, \delta}$. So that if we choose a positive real number $r_1 < |\lambda||q|^{1/2}/C$, then there exist $D_1, B_1 > 0$ (independent of K) such that

$$(15) \quad \sum_{m > 0} \left\| \frac{\phi(q^m \lambda)}{\Theta(q^m \lambda/t)} \right\|_{\mathbb{F}} \leq D_1 (B_1)^K |q|^{K(K-1)/2} |t|^K$$

for all $t \in \mathcal{T}_{\lambda, q, \delta, r_1}$.

In a second step, we give estimates for the sum $\sum_{m \leq 0} \phi(q^m \lambda)/\Theta(q^m \lambda/t) - \sum_{n=0}^K f_n t^n$, where the f_n are defined in the Taylor expansion (10). From the formula

$$q^{n(n-1)/2} t^n = \sum_{m \in \mathbb{Z}} \frac{(q^m \lambda)^n}{\Theta(q^m \lambda/t)}$$

for all $n \geq 0$, given in [28], we can write (at least formally)

$$(16) \quad \sum_{m \leq 0} \phi(q^m \lambda) / \Theta(q^m \lambda / t) - \sum_{n=0}^K f_n t^n = \sum_{m \leq 0} \frac{1}{\Theta(q^m \lambda / t)} \left(\sum_{n \geq K+1} \frac{f_n}{q^{n(n-1)/2}} (q^m \lambda)^n \right) \\ - \sum_{n=0}^K \frac{f_n}{q^{n(n-1)/2}} \left(\sum_{m > 0} \frac{(q^m \lambda)^n}{\Theta(q^m \lambda / t)} \right)$$

for all $t \in \mathbb{C}^*$. From the fact that ϕ has convergent expansion (10) on $D(0, \rho_0)$, and since $|\lambda| < \rho_0$, there exist $C, A > 0$, with $A < 1/|\lambda|$, such that

$$(17) \quad \left\| \frac{f_n}{q^{n(n-1)/2}} \right\|_{\mathbb{F}} \leq CA^n$$

for all $n \geq 0$. From (16) and (17), we deduce that

$$(18) \quad \left\| \sum_{m \leq 0} \phi(q^m \lambda) / \Theta(q^m \lambda / t) - \sum_{n=0}^K f_n t^n \right\|_{\mathbb{F}} \leq \mathcal{A}(t) + \mathcal{B}(t)$$

where

$$\mathcal{A}(t) = \sum_{m \leq 0} \frac{1}{|\Theta(q^m \lambda / t)|} \left(\sum_{n \geq K+1} CA^n (|q|^m |\lambda|)^n \right)$$

and

$$\mathcal{B}(t) = \sum_{n=0}^K CA^n \left(\sum_{m > 0} \frac{|(q^m \lambda)^n|}{|\Theta(q^m \lambda / t)|} \right),$$

for all $t \in \mathbb{C}^*$.

We give estimates for $\mathcal{A}(t)$. By summing up the geometric series (convergent because $A|q|^m |\lambda| \leq A|\lambda| < 1$ for all $m \leq 0$) and changing m into $-m$, we first have that there exists $D > 0$ such that

$$(19) \quad \mathcal{A}(t) \leq DA^{K+1} \sum_{m \geq 0} \frac{(|q|^{-m} |\lambda|)^{K+1}}{|\Theta(q^{-m} \lambda / t)|}$$

for all $t \in \mathbb{C}^*$. From (13), we have that

$$|\Theta(q^{-m} \lambda / t)| \geq K_1 \delta |q|^{-K(K-1)/2} \left| \frac{q^{-m} \lambda}{t} \right|^K$$

for all $m \geq 0$, all $t \in \mathcal{R}_{\lambda, q, \delta}$. We then have that

$$(20) \quad \frac{(|q|^{-m} |\lambda|)^{K+1}}{|\Theta(q^{-m} \lambda / t)|} \leq \frac{|\lambda|}{K_1 \delta} |q|^{K(K-1)/2} |t|^K \left(\frac{1}{|q|} \right)^m$$

for all $m \geq 0$, all $t \in \mathcal{R}_{\lambda, q, \delta}$. From (19) and (20), we deduce that there exist $D_2, B_2 > 0$ (independent of K) such that

$$(21) \quad \mathcal{A}(t) \leq D_2 (B_2)^K |q|^{K(K-1)/2} |t|^K$$

for all $t \in \mathcal{R}_{\lambda, q, \delta}$.

In the next step, we get estimates for $\mathcal{B}(t)$. From (13), we have that

$$|\Theta(q^m \lambda/t)| \geq K_1 \delta |q|^{-(K+1)K/2} \left| \frac{q^m \lambda}{t} \right|^{K+1}$$

for all $m > 0$, all $t \in \mathcal{R}_{\lambda, q, \delta}$. We deduce that

$$(22) \quad \frac{|(q^m \lambda)^n|}{|\Theta(q^m \lambda/t)|} \leq \frac{|\lambda|^n}{K_1 \delta} \left(\frac{1}{|\lambda|} \right)^{K+1} |q|^{(K+1)K/2} |t|^{K+1} \left(\frac{1}{|q|} \right)^m$$

for all $m > 0$, all $0 \leq n \leq K$. From (22), the equality $(K+1)K/2 = K + K(K-1)/2$ and the fact that $|t| < r_1$ whenever $t \in \mathcal{T}_{\lambda, q, \delta, r_1}$, we obtain that there exist $D_3, B_3 > 0$ (independent of K) such that

$$(23) \quad \mathcal{B}(t) \leq D_3 (B_3)^K |q|^{K(K-1)/2} |t|^K$$

for all $t \in \mathcal{T}_{\lambda, q, \delta, r_1}$.

Finally, using the estimates

$$\begin{aligned} \left\| \sum_{m \in \mathbb{Z}} \phi(q^m \lambda) / \Theta(q^m \lambda/t) - \sum_{n=0}^K f_n t^n \right\|_{\mathbb{F}} &\leq \left\| \sum_{m \leq 0} \phi(q^m \lambda) / \Theta(q^m \lambda/t) - \sum_{n=0}^K f_n t^n \right\|_{\mathbb{F}} \\ &\quad + \left\| \sum_{m > 0} \phi(q^m \lambda) / \Theta(q^m \lambda/t) \right\|_{\mathbb{F}} \end{aligned}$$

we deduce from (15), (18), (21), (23) that

$$(24) \quad \left\| \sum_{m \in \mathbb{Z}} \phi(q^m \lambda) / \Theta(q^m \lambda/t) - \sum_{n=0}^K f_n t^n \right\|_{\mathbb{F}} \leq D_4 (B_4)^K |q|^{K(K-1)/2} |t|^K$$

for some $D_4, B_4 > 0$ (independent of K). Now, for $K \in \mathbb{N}$, $K \geq 1$ one may write

$$\left\| \sum_{m \in \mathbb{Z}} \phi(q^m \lambda) / \Theta(q^m \lambda/t) - \sum_{n=0}^{K-1} f_n t^n \right\|_{\mathbb{F}} \leq \left\| \sum_{m \in \mathbb{Z}} \phi(q^m \lambda) / \Theta(q^m \lambda/t) - \sum_{n=0}^K f_n t^n \right\|_{\mathbb{F}} + \|f_K t^K\|_{\mathbb{F}}$$

and take into account (24) and (17) in order to obtain (11), as desired. \square

Proposition 2 *Let V be an open and bounded set in \mathbb{C}^* and $D(0, \rho_0)$ be a disc such that $V \cap D(0, \rho_0) \neq \emptyset$. Let ϕ be a holomorphic function on $Vq^{\mathbb{N}} \cup D(0, \rho_0)$ with values in $(\mathbb{F}, \|\cdot\|_{\mathbb{F}})$ which satisfies the estimates : There exist $C, K > 0$, such that*

$$(25) \quad \|\phi(xq^m)\|_{\mathbb{F}} \leq K |q|^{m^2/2} C^m$$

for all $m \geq 0$, all $x \in V$. Then, the function $M\phi(\tau) := \tau\phi(\tau)$ is holomorphic on $Vq^{\mathbb{N}} \cup D(0, \rho_0)$ and satisfies estimates of the form (8). Let $\lambda \in V \cap D(0, \rho_0)$. We have the following equality

$$\mathcal{L}_q^\lambda(M\phi)(t) = t\mathcal{L}_q^\lambda(\phi)(qt)$$

for all $t \in \mathcal{T}_{\lambda, q, \delta, r_1}$, if $r_1 < |\lambda q^{1/2}|/(C|q|)$.

Proof From the estimates (25), we get a constant $r > 0$ such that

$$\|(M\phi)(xq^m)\|_{\mathbb{F}} \leq rK|q|^{m^2/2}(|q|C)^m$$

for all $m \geq 0$, all $x \in V$. From Proposition 1, we deduce that $\mathcal{L}_q^\lambda(M\phi)(t)$ defines a holomorphic function on $\mathcal{T}_{\lambda,q,\delta,r_1}$, if $r_1 < |\lambda q^{1/2}|/(C|q|)$. On the other hand,

$$(26) \quad t\mathcal{L}_q^\lambda(\phi)(qt) = \sum_{m \in \mathbb{Z}} \frac{t\phi(q^m\lambda)}{\Theta(q^m\lambda/(qt))}.$$

But we have that

$$\frac{t}{\Theta(q^m\lambda/(qt))} = \frac{q^m\lambda}{\Theta(q^m\lambda/t)}$$

for all $m \in \mathbb{Z}$. Indeed, put $y = q^m\lambda/(qt)$ in the identity $\Theta(qy) = qy\Theta(y)$. From (26), we get that

$$t\mathcal{L}_q^\lambda(\phi)(qt) = \sum_{m \in \mathbb{Z}} \frac{q^m\lambda\phi(q^m\lambda)}{\Theta(q^m\lambda/t)} = \mathcal{L}_q^\lambda(M\phi)(t)$$

for all $t \in \mathcal{T}_{\lambda,q,\delta,r_1}$. □

For convenience, we recall the following concepts.

Definition 1 A series $\hat{f}(t) = \sum_{n \geq 0} f_n t^n \in \mathbb{C}[[t]]$ is said to be q -Gevrey of order 1 if its so-called formal q -Borel transform of order 1,

$$\hat{\mathcal{B}}_q \hat{f}(\tau) = \sum_{n \geq 0} \frac{f_n}{q^{n(n-1)/2}} \tau^n,$$

converges (i.e. it has positive radius of convergence).

The formal q -Laplace transform of order 1 of a series $\hat{g}(\tau) = \sum_{n \geq 0} g_n \tau^n \in \mathbb{C}[[\tau]]$ is defined as

$$\hat{\mathcal{L}}_q \hat{g}(t) = \sum_{n \geq 0} q^{n(n-1)/2} g_n t^n,$$

so that these formal transforms are inverse of each other.

It is immediate to check that, in agreement with Proposition 2, we have that for every $\hat{g} \in \mathbb{C}[[\tau]]$,

$$(27) \quad \hat{\mathcal{L}}_q(\tau \hat{g})(t) = t \hat{\mathcal{L}}_q \hat{g}(qt).$$

3 A Cauchy problem in a weighted Banach space of formal Laurent series

With the help of the q -Laplace transform we will change our initial problem (3), (4) into an equivalent one (5), (6), whose study will require the consideration of two auxiliary Cauchy problems. The first of them, which we are going to present in this Section, will be crucial in the study of the q -exponential growth of the coefficients of a solution of (5), (6). Although our equation involves a complex number q with $|q| > 1$, in this Section and in Section 5 we will be only concerned with the value $|q|$, so we directly work with a real value $q > 1$.

Definition 2 We consider the vector space $\mathbb{E}_{q,(T,X)}$ of formal Laurent power series

$$(28) \quad V(\xi, x) = \sum_{l \in \mathbb{Z}, h \geq 0} v_{l,h} \xi^l \frac{x^h}{h!} \in \mathbb{C}[[\xi, \xi^{-1}, x]]$$

such that

$$\|V(\xi, x)\|_{(T,X)} := \sum_{l \in \mathbb{Z}, h \geq 0} \frac{|v_{l,h}|}{q^{P(l,h)}} T^l \frac{X^h}{h!} < \infty,$$

where $T, X > 0$, $q > 1$ are positive real numbers and where

$$P(l, h) = \begin{cases} \frac{1}{4}l^2 + \frac{1}{2}lh - \frac{1}{2}h^2 & \text{if } l \geq 0, h \geq 0, \\ -(1/2)h^2 & \text{if } l \leq 0, h \geq 0. \end{cases}$$

The space $(\mathbb{E}_{q,(T,X)}, \|\cdot\|_{(T,X)})$ is a Banach space.

Remark: Notice that we have a continuous inclusion $(\mathbb{E}_{q,(T,X')}, \|\cdot\|_{(T,X')}) \hookrightarrow (\mathbb{E}_{q,(T,X)}, \|\cdot\|_{(T,X)})$ when $0 < X \leq X'$.

We consider the integration operator ∂_x^{-1} defined on $\mathbb{C}[[\xi, \xi^{-1}, x]]$ by

$$\partial_x^{-1}(V(\xi, x)) := \sum_{l \in \mathbb{Z}, h \geq 1} v_{l,h-1} \xi^l \frac{x^h}{h!} \in \mathbb{C}[[\xi, \xi^{-1}, x]]$$

Lemma 1 Let $m_1, s, h_1, h_2 \geq 0$ be nonnegative integers. Let $T, X > 0$. Assume that the inequalities hold

$$(29) \quad s + h_2 \geq 2h_1 \quad , \quad m_1 \geq s + h_2.$$

Then, there exist $C > 0$ (depending q, s, h_1, h_2, m_1) such that

$$(30) \quad \|x^s (\partial_x^{-h_2} V)(q^{h_1} \xi, \frac{x}{q^{m_1}})\|_{(T,X)} \leq CX^{(s+h_2)} \|V(\xi, x)\|_{(T,X)}$$

for all $V(\xi, x) \in \mathbb{E}_{q,(T,X)}$.

Proof Let $V(\xi, x) \in \mathbb{C}[[\xi, \xi^{-1}, x]]$ as in (28). We have that

$$x^s (\partial_x^{-h_2} V)(q^{h_1} \xi, \frac{x}{q^{m_1}}) = \sum_{l \in \mathbb{Z}, h \geq h_2+s} v_{l, h-(s+h_2)} \frac{q^{h_1 l} h!}{q^{m_1(h-s)} (h-s)!} \xi^l \frac{x^h}{h!}$$

From the definition of the norm $\|\cdot\|_{(T,X)}$, we get that

$$(31) \quad \|x^s (\partial_x^{-h_2} V)(q^{h_1} \xi, \frac{x}{q^{m_1}})\|_{(T,X)} = \sum_{l \in \mathbb{Z}, h \geq h_2+s} \frac{|v_{l, h-(s+h_2)}|}{q^{P(l, h-(s+h_2))}} T^l \frac{X^{h-(s+h_2)}}{(h-(s+h_2))!} \times \left\{ \frac{1}{q^{P(l, h)-P(l, h-(s+h_2))-h_1 l+m_1(h-s)}} \frac{(h-(s+h_2))!}{(h-s)!} X^{s+h_2} \right\}.$$

In the rest of the proof, we will show that there exists a constant $C > 0$ (depending on q, s, h_1, h_2, m_1) such that

$$(32) \quad \frac{1}{q^{P(l, h)-P(l, h-(s+h_2))-h_1 l+m_1(h-s)}} \leq C$$

for all $l \in \mathbb{Z}$, $h \geq 0$. Indeed, if $l \geq 0$, then

$$P(l, h) - P(l, h - (s + h_2)) - h_1 l + m_1(h - s) = l\left(\frac{s + h_2}{2} - h_1\right) + h(m_1 - (s + h_2)) - m_1 s + \frac{(s + h_2)^2}{2}$$

for all $h \geq 0$. From the assumption (29), we deduce that the inequalities (32) hold for $l \geq 0$, $h \geq 0$.

If $l \leq 0$, then

$$P(l, h) - P(l, h - (s + h_2)) - h_1 l + m_1(h - s) = l(-h_1) + h(m_1 - (s + h_2)) - m_1 s + \frac{(s + h_2)^2}{2}$$

for all $h \geq 0$. From the assumption (29), we deduce that the inequalities (32) hold for $l \leq 0$, $h \geq 0$.

Finally, the inequality (30) follows from the expression (31) and the estimates (32). \square

Lemma 2 *Let $s, h_1 \geq 0$ and $T_0, X_0 > 0$. Then, there exists a constant $C_1 > 0$ (depending on q, s, h_1, T_0, X_0) such that for all $0 < X_1 \leq X_0 q^{-s}$ and for all $T_1 > 0$ satisfying*

$$(33) \quad q^{-h_1} T_0 \leq T_1 \leq T_0 q^{\frac{s}{2} - h_1}$$

one has

$$(34) \quad \|x^s V(q^{h_1} \xi, x)\|_{(T_1, X_1)} \leq C_1 \|V(\xi, x)\|_{(T_0, X_0)}$$

for all $V(\xi, x) \in \mathbb{E}_{q, (T_0, X_0)}$.

Proof Let $V(\xi, x) \in \mathbb{C}[[\xi, \xi^{-1}, x]]$. From the definition of the norm $\|\cdot\|_{(T, X)}$, one can write

$$(35) \quad \|x^s V(q^{h_1} \xi, x)\|_{(T_1, X_1)} = \sum_{l \in \mathbb{Z}, h \geq s} \frac{|v_{l, h-s}|}{q^{P(l, h-s)}} T_0^l \frac{X_0^{(h-s)}}{(h-s)!} \times \left\{ \frac{1}{q^{P(l, h) - P(l, h-s) - h_1 l}} \left(\frac{T_1}{T_0}\right)^l \left(\frac{X_1}{X_0}\right)^h X_0^s \right\}$$

In the rest of the proof, we will show that there exists a constant $C_1 > 0$ (depending on q, s, h_1) such that for all $0 < X_1 \leq X_0 q^{-s}$ and all T_1 satisfying (33), one has

$$(36) \quad \frac{1}{q^{P(l, h) - P(l, h-s) - h_1 l}} \left(\frac{T_1}{T_0}\right)^l \left(\frac{X_1}{X_0}\right)^h \leq C_1$$

for all $l \in \mathbb{Z}$, all $h \geq 0$.

If $l \geq 0$, then

$$P(l, h) - P(l, h - s) - h_1 l = l\left(\frac{s}{2} - h_1\right) + h(-s) + \frac{s^2}{2}$$

for all $h \geq 0$. So, we get that (36) holds for all $l \geq 0$, all $h \geq 0$.

If $l \leq 0$, then

$$P(l, h) - P(l, h - s) - h_1 l = l(-h_1) + h(-s) + \frac{s^2}{2}$$

for all $h \geq 0$. Hence, we get that (36) holds for all $l \leq 0$, all $h \geq 0$.

Finally, the inequality (34) follows from the expression (35) and the estimates (36). \square

Lemma 3 Let $h_2 \geq 0$ and $T_0, X_0 > 0$. Then, there exists a constant $C_2 > 0$ (depending on q, h_2, T_0, X_0) such that for all $0 < X_1 \leq X_0 q^{-h_2}$ and for all $T_1 > 0$ satisfying

$$(37) \quad T_0 \leq T_1 \leq T_0 q^{h_2/2},$$

one has

$$(38) \quad \|\partial_x^{-h_2} V(\xi, x)\|_{(T_1, X_1)} \leq C_2 \|V(\xi, x)\|_{(T_0, X_0)}$$

for all $V(\xi, x) \in \mathbb{E}_{q, (T_0, X_0)}$.

Proof Let $V(\xi, x) \in \mathbb{C}[[\xi, \xi^{-1}, x]]$. From the definition of the norm $\|\cdot\|_{(T, X)}$, one can write

$$(39) \quad \|\partial_x^{-h_2} V(\xi, x)\|_{(T_1, X_1)} = \sum_{l \in \mathbb{Z}, h \geq h_2} \frac{|v_{l, h-h_2}|}{q^{P(l, h-h_2)}} T_0^l \frac{X_0^{(h-h_2)}}{(h-h_2)!} \times \left\{ \frac{1}{q^{P(l, h)-P(l, h-h_2)}} \left(\frac{T_1}{T_0}\right)^l \left(\frac{X_1}{X_0}\right)^h \frac{(h-h_2)!}{h!} X_0^{h_2} \right\}$$

In the rest of the proof, we will show that there exists a constant $C_2 > 0$ (depending on q, h_2) such that for all $0 < X_1 \leq X_0 q^{-h_2}$ and T_1 satisfying (37) one has

$$(40) \quad \frac{1}{q^{P(l, h)-P(l, h-h_2)}} \left(\frac{T_1}{T_0}\right)^l \left(\frac{X_1}{X_0}\right)^h \leq C_2$$

for all $l \in \mathbb{Z}$, all $h \geq 0$.

If $l \geq 0$, then

$$P(l, h) - P(l, h-h_2) = l \left(\frac{h_2}{2}\right) + h(-h_2) + \frac{h_2^2}{2},$$

for all $h \geq 0$. Hence, we get that (40) holds for all $l \geq 0$, all $h \geq 0$.

If $l \leq 0$, then

$$P(l, h) - P(l, h-h_2) = h(-h_2) + \frac{1}{2} h_2^2,$$

for all $h \geq 0$, and we get that (40) holds for all $l \leq 0$, all $h \geq 0$.

Finally, the inequality (38) follows from the expression (39) and the estimates (40). \square

Let $S, m_{0,k}, m_{1,k}$, $0 \leq k \leq S-1$ be positive integers. Let \mathcal{D} the linear operator from $\mathbb{C}[[\xi, \xi^{-1}, x]]$ into $\mathbb{C}[[\xi, \xi^{-1}, x]]$ defined by

$$\mathcal{D}(V(\xi, x)) := \partial_x^S V(\xi, x) - \sum_{k=0}^{S-1} a_k(x) (\partial_x^k V)(q^{m_{0,k}} \xi, x/q^{m_{1,k}}),$$

for all $V \in \mathbb{C}[[\xi, \xi^{-1}, x]]$, where $a_k(x) = \sum_{s \in I_k} a_{ks} x^s \in \mathbb{C}[x]$, with I_k be a finite subset of \mathbb{N} , for $0 \leq k \leq S-1$.

We make the following hypothesis.

Assumption (A) For all $0 \leq k \leq S-1$, for all $s \in I_k$, we have

$$s + S - k \geq 2m_{0,k} \quad , \quad m_{1,k} \geq s + S - k.$$

We consider the operator \mathcal{A} from $\mathbb{C}[[\xi, \xi^{-1}, x]]$ into $\mathbb{C}[[\xi, \xi^{-1}, x]]$ defined by

$$\mathcal{A}(V(\xi, x)) = V(\xi, x) - \mathcal{D}(\partial_x^{-S} V(\xi, x)) = \sum_{k=0}^{S-1} a_k(x) (\partial_x^{k-S} V)(q^{m_{0,k}} \xi, x/q^{m_{1,k}})$$

for all $V \in \mathbb{C}[[\xi, \xi^{-1}, x]]$.

From Lemma 1, we deduce the following

Lemma 4 *Let $T > 0$. Then, there exists $X > 0$ such that \mathcal{A} is a linear bounded operator from $(\mathbb{E}_{q,(T,X)}, \|\cdot\|_{(T,X)})$ into itself. Moreover, we have that*

$$\|\mathcal{A}(V(\xi, x))\|_{(T,X)} \leq \frac{1}{2} \|V(\xi, x)\|_{(T,X)},$$

for all $V \in \mathbb{E}_{q,(T,X)}$.

From Lemma 4, we deduce the next

Corollary 1 *Let $T > 0$. Then, there exists $X > 0$ such that $\mathcal{D} \circ \partial_x^{-S}$ is an invertible linear operator from $(\mathbb{E}_{q,(T,X)}, \|\cdot\|_{(T,X)})$ into itself. In particular, there exists $C > 0$ such that*

$$\|\mathcal{D}(\partial_x^{-S} b(\xi, x))\|_{(T,X)} \leq C \|b(\xi, x)\|_{(T,X)}$$

for all $b(\xi, x) \in \mathbb{E}_{q,(T,X)}$.

Definition 3 *Let $T_{0,j} > 0$, $0 \leq j \leq S-1$ be real numbers. We say that $(T_{0,j})_{0 \leq j \leq S-1}$ satisfies the assumption **(B)** if*

1) *The set*

$$\mathbb{T} := \bigcap_{0 \leq k \leq S-1} \bigcap_{k \leq j \leq S-1, s \in I_k} [q^{-m_{0,k}} T_{0,j}, T_{0,j} q^{\frac{s+j-k-2m_{0,k}}{2}}]$$

is not empty,

2) *There exists $T_1 \in \mathbb{T}$ such that*

$$\mathbb{T}_1 := [T_1, T_1 q^{S/2}] \bigcap \bigcap_{j=0}^{S-1} [T_{0,j}, T_{0,j} q^{j/2}]$$

is not empty.

Example: Let $S \geq 1$. For all $0 \leq k \leq S-1$, let $I_k \subset \mathbb{N}$ such that $s \in I_k$ implies $s \geq 2m_{0,k} + k$. For all $0 \leq j \leq S-1$, we put $T_{0,j} = T_0 > 0$. Then, we have that $T_0 \in \mathbb{T}$ and that $\mathbb{T}_1 = \{T_0\}$. So that $(T_{0,j})_{0 \leq j \leq S-1}$ satisfies the assumption **(B)**.

Theorem 1 *Let $S \geq 1$ be an integer. For all $0 \leq k \leq S-1$, let $m_{0,k}, m_{1,k}$ be positive integers and $a_k(x) = \sum_{s \in I_k} a_{ks} x^s \in \mathbb{C}[x]$. We make the hypothesis that the assumption **(A)** holds.*

We consider the following functional equation

$$(41) \quad \partial_x^S V(\xi, x) = \sum_{k=0}^{S-1} a_k(x) (\partial_x^k V)(q^{m_{0,k}} \xi, x/q^{m_{1,k}})$$

with initial conditions

$$(42) \quad (\partial_x^j V)(\xi, 0) = \phi_j(\xi) \quad , \quad 0 \leq j \leq S-1.$$

We assume that $\phi_j(\xi) \in \mathbb{E}_{q,(T_{0,j},X_0)}$, for $0 \leq j \leq S-1$, where $X_0 > 0$ and $(T_{0,j})_{0 \leq j \leq S-1}$ satisfy the assumption **(B)**. Then, there exists $X > 0$ and for $T \in \mathbb{T}_1$, the problem (41), (42) has a unique solution $V(\xi, x) \in \mathbb{E}_{q,(T,X)}$. Moreover, there exists $C > 0$ (depending on $S, q, a_k(x), m_{0,k}, m_{1,k}$, for $0 \leq k \leq S-1$ and $X_0, T_{0,j}$, for $0 \leq j \leq S-1$) such that

$$\|V(\xi, x)\|_{(T,X)} \leq C \sum_{j=0}^{S-1} \|\phi_j(\xi)\|_{(T_{0,j},X_0)}.$$

Proof A formal series $V(\xi, x) \in \mathbb{C}[[\xi, \xi^{-1}, x]]$ which satisfies (42) can be written in the form $V(\xi, x) = \partial_x^{-S} U(\xi, x) + I(\xi, x)$ where

$$I(\xi, x) = \sum_{j=0}^{S-1} \phi_j(\xi) \frac{x^j}{j!}$$

and $U(\xi, x) \in \mathbb{C}[[\xi, \xi^{-1}, x]]$. A formal series $V(\xi, x) \in \mathbb{C}[[\xi, \xi^{-1}, x]]$ is a solution of the problem (41), (42) if and only if $U(\xi, x)$ satisfies the equation

$$(43) \quad \mathcal{D}(\partial_x^{-S} U(\xi, x)) = -\mathcal{D}(I(\xi, x)).$$

By construction, we have that

$$-\mathcal{D}(I(\xi, x)) = \sum_{k=0}^{S-1} \sum_{j=k}^{S-1} \sum_{s \in I_k} \frac{a_{ks}}{q^{m_{1,k}(j-k)}(j-k)!} x^{s+j-k} \phi_j(q^{m_{0,k}} \xi)$$

From Lemma 2 and the assumption **(B)**, there exists $X_1 > 0$ such that for all $T_1 \in \mathbb{T}$ we have that

$$x^{s+j-k} \phi_j(q^{m_{0,k}} \xi) \in \mathbb{E}_{q,(T_1, X_1)}$$

for all $0 \leq k \leq S-1$, all $k \leq j \leq S-1$, all $s \in I_k$. Moreover, there exists $C_1 > 0$ (depending on $I_k, j, m_{0,k}, X_0, T_{0,j}$) such that

$$(44) \quad \|x^{s+j-k} \phi_j(q^{m_{0,k}} \xi)\|_{(T_1, X_1)} \leq C_1 \|\phi_j(\xi)\|_{(T_{0,j}, X_0)}.$$

We deduce that $\mathcal{D}(I(\xi, x)) \in \mathbb{E}_{q,(T_1, X_1)}$ and from (44) there exists a constant $C'_1 > 0$ (depending on $q, a_k(x), m_{0,k}, m_{1,k}$, for $0 \leq k \leq S-1$ and $X_0, T_{0,j}$, for $0 \leq j \leq S-1$) such that

$$(45) \quad \|\mathcal{D}(I(\xi, x))\|_{(T_1, X_1)} \leq C'_1 \sum_{j=0}^{S-1} \|\phi_j(\xi)\|_{(T_{0,j}, X_0)}.$$

From Corollary 1, we deduce that the equation (43) has a unique solution $U(\xi, x) \in \mathbb{E}_{q,(T_1, X_1)}$. Moreover, there exists a constant $C_2 > 0$ (depending on $q, a_k(x), m_{0,k}, m_{1,k}$, for $0 \leq k \leq S-1$) such that

$$(46) \quad \|U(\xi, x)\|_{(T_1, X_1)} \leq C_2 \|\mathcal{D}(I(\xi, x))\|_{(T_1, X_1)}.$$

Now, from the assumption **(B)**, we choose $T_1 \in \mathbb{T}$ in such a way that \mathbb{T}_1 is not empty. Let $T_2 \in \mathbb{T}_1$. From Lemma 3, there exists $X_2 < X_1$ such that $\partial_x^{-S} U(\xi, x) \in \mathbb{E}_{q,(T_2, X_2)}$. Moreover, there exists a constant $C_3 > 0$ (depending on q, S, T_1, X_1) such that

$$(47) \quad \|\partial_x^{-S} U(\xi, x)\|_{(T_2, X_2)} \leq C_3 \|U(\xi, x)\|_{(T_1, X_1)}$$

From Lemma 2, there exists $X_3 < X_2$ such that $I(\xi, x) \in \mathbb{E}_{q, (T_2, X_3)}$. Moreover, there exists a constant $C_4 > 0$ (depending on $S, q, T_{0,j}, X_0$, for $0 \leq j \leq S-1$) such that

$$(48) \quad \|I(\xi, x)\|_{(T_2, X_3)} \leq C_4 \sum_{j=0}^{S-1} \|\phi_j(\xi)\|_{(T_{0,j}, X_0)}.$$

Finally, the formal series $V(\xi, x) = \partial_x^{-S} U(\xi, x) + I(\xi, x)$, solution of the problem (41), (42), belongs to $\mathbb{E}_{q, (T_2, X_3)}$. Moreover, from the inequalities (45), (46), (47) and (48), we get a constant C_5 (depending on $S, q, a_k(x), m_{0,k}, m_{1,k}$, for $0 \leq k \leq S-1$ and $X_0, T_{0,j}$, for $0 \leq j \leq S-1$) such that

$$\|V(\xi, x)\|_{(T_2, X_3)} \leq C_5 \sum_{j=0}^{S-1} \|\phi_j(\xi)\|_{(T_{0,j}, X_0)}.$$

□

4 A Cauchy problem in analytic spaces of q -exponential growth

Let $S \geq 1$, $r_1, r_2 \geq 0$ be integers. For all $0 \leq k \leq S-1$, let $m_{0,k}, m_{1,k}$ be positive integers and $b_k(z) = \sum_{s \in I_k} b_{ks} z^s$ be a polynomial in z , where I_k is a subset of \mathbb{N} .

Lemma 5 *For every choice of formal series $\hat{X}_j \in \mathbb{C}[[t]]$, $0 \leq j \leq S-1$, the Cauchy problem (3), (4) has a unique solution in the form of a formal power series $\hat{X}(t, z) = \sum_{h \geq 0} \hat{X}_h(t) \frac{z^h}{h!}$, where $\hat{X}_h \in \mathbb{C}[[t]]$ for every $h \geq 0$.*

Proof Let us put $\hat{X}_h(t) = \sum_{m \geq 0} f_{m,h} t^m$, $h \geq 0$, $f_{m,h} \in \mathbb{C}$. By substituting \hat{X} in (3), one can check that the left hand side turns out to be

$$\sum_{h \geq 0} \left(\sum_{m=0}^{r_2-1} f_{m,h+S} t^m + \sum_{m=r_2}^{\infty} (f_{m,h+S} + (h+1)^{r_1} f_{m-r_2, h+S} q^{r_2(r_2-1)/2} q^{r_2(m-r_2)}) t^m \right) \frac{z^h}{h!},$$

while the right hand side is

$$\sum_{h \geq 0} \left(\sum_{k=0}^{S-1} \sum_{h_1+h_2=h, h_1 \in I_k} \frac{b_{kh_1} t^{m_{0,k}} \hat{X}_{h_2+k}(q^{m_{0,k}} t)}{h_2! q^{m_{1,k} h_2}} \right) z^h.$$

The values $f_{m,h}$, $m \geq 0$, $0 \leq h \leq S-1$, are given by the initial conditions. We begin obtaining $f_{m,S}$, $m \geq 0$. We look at the coefficients of z^0 at both sides and impose them to be equal:

$$\sum_{m=0}^{r_2-1} f_{m,S} t^m + \sum_{m=r_2}^{\infty} (f_{m,S} + f_{m-r_2,S} q^{r_2(r_2-1)/2} q^{r_2(m-r_2)}) t^m = \sum_{k=0}^{S-1} \sum_{0 \in I_k} b_{k0} t^{m_{0,k}} \hat{X}_k(q^{m_{0,k}} t).$$

Since the right hand side is determined by the \hat{X}_j , $0 \leq j \leq S-1$, we can recursively obtain the $f_{m,S}$, $m \geq 0$, by imposing the equality of the coefficients of t^m in each side for every $m = 0, 1, 2, \dots$

We can repeat this argument in a similar way as the second index in $f_{m,h}$ increases. □

With the help of the q -Laplace transform, we reformulate our problem. Consider the Cauchy problem

$$(49) \quad ((z\partial_z + 1)^{r_1}\tau^{r_2} + 1)\partial_z^S \hat{W}(\tau, z) = \sum_{k=0}^{S-1} b_k(z)\tau^{m_{0,k}}(\partial_z^k \hat{W})(\tau, zq^{-m_{1,k}})$$

with initial conditions

$$(50) \quad (\partial_z^j \hat{W})(\tau, 0) = \hat{W}_j(\tau) \in \mathbb{C}[[\tau]], \quad 0 \leq j \leq S-1.$$

Lemma 6 *The formal series $\hat{X}(t, z) = \sum_{h \geq 0} \hat{X}_h(t) \frac{z^h}{h!}$, where $\hat{X}_h \in \mathbb{C}[[t]]$ for every $h \geq 0$, satisfies*

the Cauchy problem (3), (4) if, and only if, the formal series $\hat{W}(\tau, z) = \sum_{h \geq 0} \hat{\mathcal{B}}_q \hat{X}_h(\tau) \frac{z^h}{h!}$ satisfies

the Cauchy problem (49), (50) with $W_j(\tau) = \hat{\mathcal{B}}_q \hat{X}_j$, $0 \leq j \leq S-1$.

Conversely, $\hat{W}(\tau, z) = \sum_{h \geq 0} \hat{W}_h(\tau) \frac{z^h}{h!}$, with $\hat{W}_h \in \mathbb{C}[[\tau]]$ for every $h \geq 0$, satisfies the Cauchy

problem (49), (50) if, and only if, the formal series $\hat{X}(t, z) = \sum_{h \geq 0} \hat{\mathcal{L}}_q \hat{W}_h(t) \frac{z^h}{h!}$ satisfies the Cauchy

problem (3), (4) with $\hat{X}_j(t) = \hat{\mathcal{L}}_q \hat{W}_j(t)$ for $0 \leq j \leq S-1$.

Proof It suffices to insert each series in the corresponding Cauchy problem and apply (27). \square

Let V be an open and bounded set in \mathbb{C}^* , and $q \in \mathbb{C}$ with $|q| > 1$. In the following result we study the q -exponential growth of the coefficients of a solution to the Cauchy problem (49), (50). We will depart from initial conditions W_j , $0 \leq j \leq S-1$, holomorphic in $Vq^{\mathbb{Z}}$. We make the assumptions **(A)** and **(B)** in the previous Section, so that we may apply Theorem 1, and we also suitably choose q and V in order to deal with a small divisors problem.

Theorem 2 *Let the assumption **(A)** (of Section 3) be fulfilled by the sets I_k and the integers $m_{0,k}, m_{1,k}$, for $0 \leq k \leq S-1$.*

1) We make the following assumptions on q and on the open set V : q is of the form $q = |q|e^{i\theta}$, with $\theta = \frac{2\pi}{br_2}$ for some $b \in \mathbb{N}$, $b \geq 1$. If we denote

$$V^{r_2} = \{x^{r_2} : x \in V\},$$

we assume that there exists $\varepsilon \in (0, \min\{\pi/b, \pi/2\})$ such that

$$V^{r_2} \cap \left(\bigcup_{l=0}^{b-1} S(-\pi + \frac{2\pi l}{b}, 2\varepsilon) \right) = \emptyset,$$

where $S(d, \varphi)$ stands for the unbounded sector in \mathbb{C} with vertex at 0, bisected by direction d and with opening φ .

*2) The following assumptions on the initial conditions hold: Let $(T_{0,j})_{0 \leq j \leq S-1}$ be a sequence satisfying the assumption **(B)**, there exists a constant $K_0 > 0$ such that*

$$(51) \quad \sup_{x \in V} |W_j(xq^l)| \leq K_0 |q|^{\frac{1}{4}l^2} \left(\frac{1}{T_{0,j}} \right)^l \frac{1}{1+l^2}, \quad \sup_{x \in V} |W_j(xq^{-l})| \leq K_0 (T_{0,j})^l \frac{1}{1+l^2}$$

for all $0 \leq j \leq S-1$, all $l \geq 0$.

Then, there exists a unique solution of (49), (50)

$$(\tau, z) \mapsto W(\tau, z) = \sum_{h \geq 0} W_h(\tau) \frac{z^h}{h!}$$

which is holomorphic on $Vq^{\mathbb{Z}} \times \mathbb{C}$. Moreover, for all $\rho > 0$, there exist two constants $C, T > 0$ (depending on $\rho, S, |q|, b_k(z), m_{0,k}, m_{1,k}$, for $0 \leq k \leq S-1$ and $T_{0,j}$, for $0 \leq j \leq S-1$) such that

$$(52) \quad \sup_{x \in V, z \in D(0, \rho)} |W(xq^l, z)| \leq CK_0 |q|^{\frac{1}{2}l^2} \left(\frac{1}{T}\right)^l, \quad \sup_{x \in V, z \in D(0, \rho)} |W(xq^{-l}, z)| \leq CK_0 T^l$$

for all $l \geq 0$ (where $K_0 > 0$ is defined in (51)).

Proof From the hypothesis 1) in the statement, there exists $\delta > 0$ such that

$$(53) \quad |(h+1)^{r_1} x^{r_2} q^{r_2 l} + 1| > \delta$$

for all $l \in \mathbb{Z}$, all $h \geq 0$, all $x \in V$. We consider the sequence of functions $W_h(\tau)$, $h \geq S$, defined as follows

$$(54) \quad \frac{W_{h+S}(xq^l)}{h!} = \sum_{k=0}^{S-1} \sum_{h_1+h_2=h, h_1 \in I_k} \frac{b_{kh_1} x^{m_{0,k}} q^{m_{0,k}l}}{((h+1)^{r_1} x^{r_2} q^{r_2 l} + 1)} \frac{W_{h_2+k}(xq^l)}{h_2! |q|^{m_{1,k}h_2}}$$

for all $h \geq 0$, all $l \in \mathbb{Z}$, all $x \in V$. One checks that the sequence $W_h(\tau)$, $h \geq 0$, of holomorphic functions on $Vq^{\mathbb{Z}}$, satisfies the recursion (54) if and only if the formal series

$$W(\tau, z) = \sum_{h \geq 0} W_h(\tau) \frac{z^h}{h!}$$

in the z variable, satisfies the problem (49), (50). From this we deduce that the solution W , if it exists, is unique.

According to (51) and (54), we can recursively prove that the sequence $(w_{l,h})_{l \in \mathbb{Z}, h \geq 0}$ defined by

$$(55) \quad w_{l,h} = \sup_{x \in V} |W_h(xq^l)|,$$

for all $l \in \mathbb{Z}$, all $h \geq 0$, consists of positive real numbers. Due to (53), the sequence $(w_{l,h})_{l \in \mathbb{Z}, h \geq 0}$ satisfies the following inequalities: There exists $r > 0$ (depending on $m_{0,k}$, V) such that

$$\frac{w_{l,h+S}}{h!} \leq \sum_{k=0}^{S-1} \sum_{h_1+h_2=h, h_1 \in I_k} \frac{|b_{kh_1}| r |q|^{m_{0,k}l}}{\delta} \frac{w_{l,h_2+k}}{h_2! |q|^{m_{1,k}h_2}}$$

for all $l \in \mathbb{Z}$, all $h \geq 0$.

We consider the sequence of real numbers $(v_{l,h})_{l \in \mathbb{Z}, h \geq 0}$ defined by the following recursion

$$(56) \quad \frac{v_{l,h+S}}{h!} = \sum_{k=0}^{S-1} \sum_{h_1+h_2=h, h_1 \in I_k} \frac{|b_{kh_1}| r |q|^{m_{0,k}l}}{\delta} \frac{v_{l,h_2+k}}{h_2! |q|^{m_{1,k}h_2}}$$

with initial conditions $v_{l,j} = w_{l,j}$, for $0 \leq j \leq S-1$, all $l \in \mathbb{Z}$. By construction, we have that

$$(57) \quad w_{l,h} \leq v_{l,h}$$

for all $l \in \mathbb{Z}$, all $h \geq 0$.

In the following, we put $a_k(x) = \sum_{s \in I_k} (|b_{ks}|r/\delta)x^s$ for $0 \leq k \leq S-1$ and we consider the formal Laurent series

$$V(\xi, x) = \sum_{l \in \mathbb{Z}, h \geq 0} v_{l,h} \xi^l \frac{x^h}{h!}.$$

From the recursion (56), we get that $V(\xi, x)$ satisfies the following Cauchy problem

$$(58) \quad \partial_x^S V(\xi, x) = \sum_{k=0}^{S-1} a_k(x) (\partial_x^k V)(\xi |q|^{m_{0,k}}, x/|q|^{m_{1,k}})$$

with initial conditions

$$(59) \quad (\partial_x^j V)(\xi, 0) = \phi_j(\xi) := \sum_{l \in \mathbb{Z}} w_{l,j} \xi^l$$

From the hypothesis (51), we get that $\phi_j(\xi)$ belongs to $\mathbb{E}_{|q|, (T_{0,j}, X_0)}$, for all $X_0 > 0$. By hypothesis, the assumption **(A)** holds for the sets I_k and the numbers $m_{0,k}, m_{1,k}$ and the assumption **(B)** is fulfilled for the sequence $T_{0,j}$, $0 \leq j \leq S-1$. From Theorem 1, we deduce that the unique solution $V(\xi, x)$ of the problem (58), (59) satisfies $V(\xi, x) \in \mathbb{E}_{|q|, (T, X)}$ for a real number $X > 0$ and $T \in \mathbb{T}_1$. Moreover, there exists a constant $C > 0$ (depending on $S, |q|, a_k(x), m_{0,k}, m_{1,k}$, for $0 \leq k \leq S-1$ and $X_0, T_{0,j}$, for $0 \leq j \leq S-1$) such that

$$(60) \quad \|V(\xi, x)\|_{(T, X)} \leq C \sum_{j=0}^{S-1} \|\phi_j(\xi)\|_{(T_{0,j}, X_0)}.$$

From the inequality $P(l, h) \leq \frac{l^2}{2} - \frac{h^2}{4}$, for all $l \in \mathbb{Z}$, $h \geq 0$, and (60) we get that there exists a constant $C' > 0$ (depending on $S, |q|, a_k(x), m_{0,k}, m_{1,k}$, for $0 \leq k \leq S-1$ and $X_0, T_{0,j}$, for $0 \leq j \leq S-1$) such that

$$(61) \quad |v_{l,h}| \leq K_0 C' |q|^{\frac{l^2}{2}} |q|^{\frac{-h^2}{4}} h! \left(\frac{1}{T}\right)^l \left(\frac{1}{X}\right)^h, \quad |v_{-l,h}| \leq K_0 C' |q|^{\frac{-h^2}{2}} T^l h! \left(\frac{1}{X}\right)^h$$

for all $l \geq 0$, all $h \geq 0$, where K_0 is the constant introduced in (51). From the inequalities (57) and (61), we get that

$$\begin{aligned} \sup_{x \in V, z \in D(0, \rho)} |W(xq^l, z)| &\leq K_0 C' |q|^{\frac{l^2}{2}} \left(\frac{1}{T}\right)^l \left(\sum_{h \geq 0} |q|^{-h^2/4} \left(\frac{\rho}{X}\right)^h\right), \\ \sup_{x \in V, z \in D(0, \rho)} |W(xq^{-l}, z)| &\leq K_0 C' T^l \left(\sum_{h \geq 0} |q|^{-h^2/2} \left(\frac{\rho}{X}\right)^h\right) \end{aligned}$$

for all $l \geq 0$, all $\rho > 0$. So that the estimates (52) hold. \square

Remark: Condition 1) in the previous statement could be replaced by a more general condition, namely: Let q and V be such that (53) is verified for some $\delta > 0$ and for all $l \in \mathbb{Z}$, all $h \geq 0$, all $x \in V$. However, we preferred to use 1) because of its easy geometrical interpretation.

5 Second auxiliary Cauchy problem

We now suppose that the initial conditions W_h , $0 \leq h \leq S - 1$, of (49), (50) are holomorphic in suitably small neighbourhoods of 0. Our next aim is to obtain information on the rate of decreasing of the derivatives of the functions W_h , $h \geq 0$, coefficients of the solution constructed in Theorem 2, near the origin. This will be done in the next Section, where we will need the second auxiliary Cauchy problem we deal with in this Section.

Definition 4 *Let $q > 1$ be given. Let us consider the space $\mathbb{H}_{(T,X)}$ of formal power series*

$$V(\xi, x) = \sum_{l \geq 0, h \geq 0} v_{l,h} \xi^l \frac{x^h}{h!} \in \mathbb{C}[[\xi, x]]$$

such that

$$|V(\xi, x)|'_{(T,X)} := \sum_{l \geq 0, h \geq 0} |v_{l,h}| T^l q^{h^2/2} \frac{X^h}{h!} < \infty,$$

where T, X are positive real numbers.

The space $(\mathbb{H}_{(T,X)}, |\cdot|'_{(T,X)})$ is a Banach algebra.

Remark: We have a continuous inclusion $(\mathbb{H}_{(T,X')}, |\cdot|'_{(T,X')}) \hookrightarrow (\mathbb{H}_{(T,X)}, |\cdot|'_{(T,X)})$ whenever $0 < X \leq X'$.

We can easily prove the following result, along the same lines as Lemma 1 in Section 3.

Lemma 7 *Let $m, s, h \geq 0$ be nonnegative integers such that $m \geq s + h$, and let $T, X > 0$. Then, there exists $C > 0$ such that*

$$|x^s (\partial_x^{-h} V)(\xi, \frac{x}{q^m})|'_{(T,X)} \leq C X^{s+h} |V(\xi, x)|'_{(T,X)}$$

for all $V(\xi, x) \in \mathbb{H}_{(T,X)}$.

The following is immediate from the definition of $\mathbb{H}_{(T,X)}$.

Lemma 8 *Let $T, X > 0$. The series $R(\xi) = \sum_{l=0}^{\infty} 2^{l+1} \xi^l$, to be considered next, belongs to $\mathbb{H}_{(T,X)}$ if, and only if, $T < 1/2$.*

Let $S, m_{1,k}$, $0 \leq k \leq S - 1$, be positive integers. Let \mathcal{F} be the linear operator from $\mathbb{C}[[\xi, x]]$ into $\mathbb{C}[[\xi, x]]$ defined by

$$\mathcal{F}(V(\xi, x)) := \partial_x^S V(\xi, x) - \sum_{k=0}^{S-1} c_k(x) R(\xi) (\partial_x^k V)(\xi, x/q^{m_{1,k}}),$$

for all $V \in \mathbb{C}[[\xi, x]]$, where

$$c_k(x) = \sum_{s \in I_k} |b_{ks}| x^s, \quad 0 \leq k \leq S - 1,$$

and $R(\xi)$ is the one in Lemma 8.

We consider the operator \mathcal{B} from $\mathbb{C}[[\xi, x]]$ into $\mathbb{C}[[\xi, x]]$ defined by

$$\mathcal{B}(V(\xi, x)) = V(\xi, x) - \mathcal{F}(\partial_x^{-S}V(\xi, x)) = \sum_{k=0}^{S-1} c_k(x)R(\xi)(\partial_x^{k-S}V)(\xi, x/q^{m_{1,k}})$$

for all $V \in \mathbb{C}[[\xi, x]]$.

From now on in this Section, we will make the following

Assumption (A2) For all $0 \leq k \leq S-1$, for all $s \in I_k$, we have

$$m_{1,k} \geq s + S - k.$$

From Lemmas 7 and 8 we deduce the following result.

Lemma 9 *Let $T \in (0, 1/2)$. Then, there exists $X > 0$ such that \mathcal{B} is a linear bounded operator from $(\mathbb{H}_{(T,X)}, |\cdot|'_{(T,X)})$ into itself, and*

$$|\mathcal{B}(V(\xi, x))|'_{(T,X)} \leq \frac{1}{2}|V(\xi, x)|'_{(T,X)},$$

for all $V \in \mathbb{H}_{(T,X)}$.

From Lemma 9, we deduce the next

Corollary 2 *Let $T \in (0, 1/2)$. Then, there exists $X > 0$ such that $\mathcal{F} \circ \partial_x^{-S}$ is an invertible linear operator from $(\mathbb{H}_{(T,X)}, |\cdot|'_{(T,X)})$ into itself. In particular, there exists $C > 0$ such that*

$$|\mathcal{F}(\partial_x^{-S}b(\xi, x))|'_{(T,X)} \leq C|b(\xi, x)|'_{(T,X)}$$

for all $b(\xi, x) \in \mathbb{H}_{(T,X)}$.

Theorem 3 *Let us consider the Cauchy problem*

$$(62) \quad \partial_x^S V(\xi, x) = \sum_{k=0}^{S-1} c_k(x)R(\xi)(\partial_x^k V)(\xi, x/|q|^{m_{1,k}})$$

with initial conditions

$$(63) \quad (\partial_x^j V)(\xi, 0) = \phi_j(\xi), \quad 0 \leq j \leq S-1,$$

and assume that $\phi_j(\xi) \in \mathbb{H}_{(T_{0,j}, X_0)}$, $0 \leq j \leq S-1$, where $X_0 > 0$ and $T_{0,j} > 0$ for $j = 0, 1, \dots, S-1$. Then, for every positive number $T_1 < \min\{T_{0,1}, \dots, T_{0,S-1}, 1/2\}$, there exists $X_1 > 0$ such that the problem (62), (63) has a unique solution $V(\xi, x) \in \mathbb{H}_{(T_1, X_1)}$. Moreover, there exists $C > 0$ (depending on S, q, X_0 , and $c_k(x), m_{1,k}, T_{0,k}$ for $0 \leq k \leq S-1$) such that

$$|V(\xi, x)|'_{(T_1, X_1)} \leq C \sum_{j=0}^{S-1} |\phi_j(\xi)|'_{(T_{0,j}, X_0)}.$$

Proof It heavily resembles that of Theorem 1, so we omit it. □

6 Estimates for the derivatives of W_j near the origin

In the Cauchy problem (49), (50) we consider initial conditions W_j which are holomorphic functions respectively defined in open sets containing the closed disc

$$\overline{D}_j = \{\tau : |\tau| \leq 1/(2(j+1)^{r_1/r_2})\}$$

for $0 \leq j \leq S-1$ (for the sake of brevity, we say that W_j is holomorphic in \overline{D}_j). Then, Cauchy's integral formula for the derivatives allows us to obtain constants $A_j > 0$ such that for every natural number $n \geq 0$ we have

$$\max_{\tau \in \overline{D}_j} |\partial^n W_j(\tau)| \leq A_j^n n!.$$

So, the assumptions in the following result are not restrictive.

Theorem 4 *Consider the Cauchy problem (49), (50). Suppose $W_j(\tau)$, $0 \leq j \leq S-1$, are holomorphic functions in \overline{D}_j such that there exist constants $T_{0,j} > 0$ and a constant $K > 0$ such that*

$$\max_{\tau \in \overline{D}_j} |\partial^n W_j(\tau)| \leq K \left(\frac{1}{T_{0,j}} \right)^n \frac{n!}{1+n^2},$$

for $n \geq 0$, $j = 0, 1, \dots, S-1$. Then there exists a formal solution of (49), (50),

$$W(\tau, z) = \sum_{h \geq 0} W_h(\tau) \frac{z^h}{h!},$$

where W_h is a holomorphic function in $\overline{D}_h = \{\tau : |\tau| \leq 1/(2(h+1)^{r_1/r_2})\}$, $h \geq S$. Moreover, there exist constants $T_1, X_1 > 0$ such that

$$(64) \quad \sup_{\tau \in \overline{D}_j} |\partial^n W_j(\tau)| \leq C_1 \left(\frac{1}{T_1} \right)^n \left(\frac{1}{X_1} \right)^j n! j! (j+1)^{r_1 n / r_2} |q|^{-j^2/2},$$

for every $n, j \geq 0$, where C_1 is a positive constant (depending on $S, q, b_k(z), m_{1,k}$, for $0 \leq k \leq S-1$ and $T_{0,j}$, for $0 \leq j \leq S-1$).

Proof We look for formal series solutions of (49), (50) of the form

$$W(\tau, z) = \sum_{h \geq 0} W_h(\tau) \frac{z^h}{h!},$$

which leads to the equalities

$$(65) \quad \frac{W_{h+S}(\tau)}{h!} = \sum_{k=0}^{S-1} \sum_{h_1+h_2=h, h_1 \in I_k} \frac{b_{kh_1} \tau^{m_{0,k}}}{((h+1)^{r_1} \tau^{r_2} + 1)} \frac{W_{h_2+k}(\tau)}{h_2! q^{m_{1,k} h_2}},$$

for all $h \geq 0$. These equations recursively define in a unique way the sequence $\{W_h\}_{h \geq 0}$, and we easily see that W_h is holomorphic in $\overline{D}_h = \{\tau : |\tau| \leq 1/(2(h+1)^{r_1/r_2})\}$, $h \geq 0$. We aim at estimating the rate of growth of the derivatives of W_h in \overline{D}_h .

Let n_0 be a natural number. Differentiating n_0 times in (65) we get

$$(66) \quad \frac{\partial^{n_0} W_{h+S}(\tau)}{h!} = \sum_{k=0}^{S-1} \sum_{h_1+h_2=h, h_1 \in I_k} b_{kh_1} \sum_{l_1+l_2=n_0} \frac{n_0!}{l_1! l_2!} \partial^{l_1} \left(\frac{\tau^{m_{0,k}}}{((h+1)^{r_1} \tau^{r_2} + 1)} \right) \frac{\partial^{l_2} W_{h_2+k}(\tau)}{h_2! q^{m_{1,k} h_2}}.$$

It is clear that

$$(67) \quad \partial^{l_1} \left(\frac{\tau^{m_{0,k}}}{((h+1)^{r_1} \tau^{r_2} + 1)} \right) = \sum_{\lambda_1 + \lambda_2 = l_1, \lambda_1 \leq m_{0,k}} \frac{l_1!}{\lambda_1! \lambda_2!} \frac{m_{0,k}!}{(m_{0,k} - \lambda_1)!} \tau^{m_{0,k} - \lambda_1} \partial^{\lambda_2} \left(\frac{1}{((h+1)^{r_1} \tau^{r_2} + 1)} \right).$$

Following the proof of Lemma 7 in [15], we get that for all $\lambda_2 \geq 0$, all $\tau \in \bar{D}_h$,

$$(68) \quad \left| \partial^{\lambda_2} \left(\frac{1}{((h+1)^{r_1} \tau^{r_2} + 1)} \right) \right| \leq \lambda_2! 2^{\lambda_2+1} (h+1)^{\frac{r_1}{r_2} \lambda_2}.$$

We take (68) into (67) to obtain that

$$\begin{aligned} \left| \partial^{l_1} \left(\frac{\tau^{m_{0,k}}}{((h+1)^{r_1} \tau^{r_2} + 1)} \right) \right| &\leq \sum_{\lambda_1 + \lambda_2 = l_1, \lambda_1 \leq m_{0,k}} \frac{l_1!}{\lambda_1! \lambda_2!} \frac{m_{0,k}!}{(m_{0,k} - \lambda_1)!} |\tau|^{m_{0,k} - \lambda_1} \lambda_2! 2^{\lambda_2+1} (h+1)^{\frac{r_1}{r_2} \lambda_2} \\ &= 2l_1! \sum_{\lambda_1=0}^{m_{0,k}} \frac{m_{0,k}!}{\lambda_1! (m_{0,k} - \lambda_1)!} |\tau|^{m_{0,k} - \lambda_1} 2^{l_1 - \lambda_1} (h+1)^{\frac{r_1}{r_2} (l_1 - \lambda_1)} \\ &= l_1! 2^{l_1+1} (h+1)^{\frac{r_1}{r_2} l_1} \left(|\tau| + \frac{1}{2(h+1)^{r_1/r_2}} \right)^{m_{0,k}} \leq l_1! 2^{l_1+1} (h+1)^{\frac{r_1}{r_2} l_1}. \end{aligned}$$

The previous estimates may be applied in (66) and they let us write

$$\frac{|\partial^{n_0} W_{h+S}(\tau)|}{n_0! h!} \leq \sum_{k=0}^{S-1} \sum_{h_1+h_2=h, h_1 \in I_k} |b_{kh_1}| \sum_{l_1+l_2=n_0} 2^{l_1+1} (h+1)^{\frac{r_1}{r_2} l_1} \frac{|\partial^{l_2} W_{h_2+k}(\tau)|}{l_2! h_2! |q|^{m_{1,k} h_2}},$$

which may be rewritten as

$$(69) \quad \frac{|\partial^{n_0} W_{h+S}(\tau)|}{n_0! h! (h+S+1)^{\frac{r_1}{r_2} n_0}} \leq \sum_{k=0}^{S-1} \sum_{h_1+h_2=h, h_1 \in I_k} |b_{kh_1}| \sum_{l_1+l_2=n_0} 2^{l_1+1} \frac{(h+1)^{\frac{r_1}{r_2} n_0}}{(h+S+1)^{\frac{r_1}{r_2} n_0}} \\ \times \frac{(h_2+k+1)^{\frac{r_1}{r_2} l_2}}{(h+1)^{\frac{r_1}{r_2} l_2}} \frac{|\partial^{l_2} W_{h_2+k}(\tau)|}{l_2! h_2! (h_2+k+1)^{\frac{r_1}{r_2} l_2} |q|^{m_{1,k} h_2}}.$$

Let us put

$$w_{n_0, h} := \sup_{\tau \in \bar{D}_h} \frac{|\partial^{n_0} W_h(\tau)|}{(h+1)^{r_1 n_0 / r_2}}, \quad n_0 \geq 0, h \geq 0.$$

From (69) we deduce that

$$\frac{w_{n_0, h+S}}{n_0! h!} \leq \sum_{k=0}^{S-1} \sum_{h_1+h_2=h, h_1 \in I_k} |b_{kh_1}| \sum_{l_1+l_2=n_0} 2^{l_1+1} \frac{w_{l_2, h_2+k}}{l_2! h_2! |q|^{m_{1,k} h_2}}.$$

Now we define a multi-sequence $\{v_{l,h}\}_{l,h}$ by

$$v_{l,h} = w_{l,h}, \quad l \geq 0, 0 \leq h \leq S-1,$$

and by the following recurrence relations for $n_0 \geq 0, h \geq 0$:

$$(70) \quad \frac{v_{n_0, h+S}}{n_0! h!} = \sum_{k=0}^{S-1} \sum_{h_1+h_2=h, h_1 \in I_k} |b_{kh_1}| \sum_{l_1+l_2=n_0} 2^{l_1+1} \frac{v_{l_2, h_2+k}}{l_2! h_2! |q|^{m_{1,k} h_2}}.$$

It is clear that $w_{l,h} \leq v_{l,h}$ for all $l \geq 0$, all $h \geq 0$. Let us consider the functions

$$c_k(x) = \sum_{s \in I_k} |b_{ks}| x^s, \quad 0 \leq k \leq S-1,$$

and

$$R(\xi) = \sum_{l=0}^{\infty} 2^{l+1} \xi^l = \frac{2}{1-2\xi}.$$

Due to the recursions (70), one can check that the formal power series $\hat{V}(\xi, x) = \sum_{l,h \geq 0} \frac{v_{l,h}}{l!} \xi^l \frac{x^h}{h!}$ is a formal solution of the Cauchy problem (62), (63) with initial conditions

$$(\partial_x^j V)(\xi, 0) = \phi_j(\xi) := \sum_{l \geq 0} \frac{w_{l,j}}{l!} \xi^l, \quad 0 \leq j \leq S-1.$$

It is immediate to check that, for any $X_0 > 0$, we have $\phi_j(\xi) \in \mathbb{H}_{(T_{0,j}, X_0)}$ for $0 \leq j \leq S-1$. From Theorem 3 and the fact that $\hat{V}(\xi, x) = \sum_{n,j \geq 0} \frac{v_{n,j}}{n!} \xi^n \frac{x^j}{j!}$ is the unique formal solution of (62), (63), we can find $X_1 > 0$ and $T_1 > 0$ such that

$$|\hat{V}(\xi, x)|'_{(T_1, X_1)} \leq C \sum_{j=0}^{S-1} |\phi_j(\xi)|'_{(T_{0,j}, X_0)}$$

for a certain $C > 0$ (depending on $S, q, c_k(x), m_{1,k}$, for $0 \leq k \leq S-1$ and $X_0, T_{0,j}$, for $0 \leq j \leq S-1$). From the last expression, one can obtain that

$$w_{n,j} \leq v_{n,j} \leq C \left(\frac{1}{T_1} \right)^n \left(\frac{1}{X_1} \right)^j n! j! q^{-j^2/2} \left[\sum_{l=0}^{S-1} |\phi_l(\xi)|'_{(T_{0,l}, X_0)} \right] \leq C_1 \left(\frac{1}{T_1} \right)^n \left(\frac{1}{X_1} \right)^j n! j! |q|^{-j^2/2}.$$

We conclude by the very definition of the multi-sequence $\{w_{n,j}\}_{n,j \geq 0}$, since

$$\max_{\tau \in \bar{D}_j} |\partial^n W_j(\tau)| = w_{n,j} (j+1)^{r_1 n / r_2} \leq C_1 \left(\frac{1}{T_1} \right)^n \left(\frac{1}{X_1} \right)^j n! j! (j+1)^{r_1 n / r_2} |q|^{-j^2/2},$$

as desired. \square

7 Analytic solutions of the Cauchy problem with Fuchsian and irregular singularities

Let W_h be the initial data in the Cauchy problem (49), (50), and suppose they are subject to the hypotheses of Theorem 2 and to the hypotheses in Theorem 4. Those results give us a sequence of functions $\{W_h\}_{h \geq 0}$, holomorphic in $Vq^{\mathbb{Z}} \cup D_h$ for each $h \geq 0$, and such that the series

$$W(\tau, z) = \sum_{h \geq 0} W_h(\tau) \frac{z^h}{h!}$$

defines a holomorphic function on $Vq^{\mathbb{Z}} \times \mathbb{C}$ which solves the Cauchy problem.

Moreover, from (55), (57) and (61) in the proof of Theorem 2 we know that

$$(71) \quad \sup_{x \in V} |W_h(xq^l)| \leq K_0 C' |q|^{\frac{l^2}{2}} |q|^{\frac{-h^2}{4}} h! \left(\frac{1}{T} \right)^l \left(\frac{1}{X} \right)^h$$

for all $l, h \geq 0$.

Let us choose $\lambda \in V$ and $\delta > 0$. By (71) we see that every W_h verifies estimates as those in (8). If we choose an integer $n(h)$ in such a way that $\lambda q^{n(h)} \in D_h$, then, according to Proposition 1, the q -Laplace transform of W_h in the direction $\lambda q^{n(h)} q^{\mathbb{Z}}$, which clearly equals $\lambda q^{\mathbb{Z}}$, is given by

$$\mathcal{L}_q^{\lambda q^{n(h)}}(W_h)(t) = \sum_{m \in \mathbb{Z}} \frac{W_h(q^m \lambda q^{n(h)})}{\Theta\left(\frac{q^m \lambda q^{n(h)}}{t}\right)} = \sum_{m \in \mathbb{Z}} \frac{W_h(q^m \lambda)}{\Theta\left(\frac{q^m \lambda}{t}\right)},$$

so that it deserves to be denoted by $\mathcal{L}_q^\lambda(W_h)(t)$. This function is well defined and holomorphic in the set $\mathcal{T}_{\lambda q^{n(h)}, q, \delta, r(h)}$, which is equal to $\mathcal{T}_{\lambda, q, \delta, r(h)}$, whenever $r(h) < |\lambda q^{n(h)} q^{1/2}| T$. We will show that these radii $r(h)$ can be taken independent of h , equal to $r_0 = |\lambda q^{1/2}| T / |q| = |\lambda q^{-1/2}| T$ for every $h \geq 0$, and we will obtain precise estimates for the corresponding q -asymptotic expansions.

Let us assume that the function W_h has the following Taylor expansion at 0,

$$(72) \quad W_h(\tau) = \sum_{n \geq 0} \frac{f_{n,h}}{q^{n(n-1)/2}} \tau^n,$$

where $f_{n,h} \in \mathbb{C}$, $n, h \geq 0$, and $\tau \in \overline{D}_h$.

Proposition 3 *In the situation assumed in this Section, there exist constants $B(h), D(h) > 0$ (to be specified) such that*

$$(73) \quad \left| \mathcal{L}_q^\lambda(W_h)(t) - \sum_{m=0}^{n-1} f_{m,h} t^m \right| \leq D(h) B(h)^n |q|^{n(n-1)/2} |t|^n$$

for all $n \geq 1$, for all $t \in \mathcal{T}_{\lambda, q, \delta, r_0}$.

Proof

According to the estimates (64) in Theorem 4, we can write

$$(74) \quad \left| \frac{f_{n,h}}{q^{n(n-1)/2}} \right| = \left| \frac{1}{n!} \partial^n W_h(0) \right| \leq C_1 \left(\frac{1}{T_1} \right)^n \left(\frac{1}{X_1} \right)^h h! (h+1)^{r_1 n / r_2} |q|^{-h^2/2} = C(h) A(h)^n$$

for every $n, h \geq 0$, where we have put, for short,

$$(75) \quad C(h) = C_1 \left(\frac{1}{X_1} \right)^h h! |q|^{-h^2/2}, \quad A(h) = \frac{1}{T_1} (h+1)^{r_1 / r_2}, \quad h \geq 0.$$

For each $h \geq 0$ we define

$$m_h := \max \left\{ m \in \mathbb{Z} : |q^m \lambda| < \frac{1}{2A(h)} \right\},$$

so that

$$(76) \quad |q^m \lambda| A(h) < \frac{1}{2}, \quad m \leq m_h.$$

Also, we recall from Theorem 3 that $T_1 < 1/2$, so

$$\frac{1}{A(h)} < \frac{1}{2(h+1)^{\frac{r_1}{r_2}}},$$

and we deduce that

$$(77) \quad q^m \lambda \in D_h, \quad m \leq m_h.$$

Moreover, by the very definition of m_h , we have that

$$(78) \quad m_h + 1 \geq \frac{-\log(2|\lambda|A(h))}{\log(|q|)}, \quad |q|^{m_h+1} \geq \frac{1}{2|\lambda|A(h)}.$$

Let $K \geq 0$ be a fixed integer. Firstly, we give estimates for $\sum_{m>m_h} W_h(q^m \lambda)/\Theta(q^m \lambda/t)$. Using (71), (12) and (14), we get

$$\left| \frac{W_h(q^m \lambda)}{\Theta(q^m \lambda/t)} \right| \leq \frac{C' K_0 h!}{K_1 \delta} \left(\frac{1}{|\lambda|} \right)^K |q|^{K(K-1)/2} |t|^K \left(\frac{|t|}{T|\lambda||q|^{1/2}} \right)^m \left(\frac{1}{X} \right)^h |q|^{-h^2/4}$$

for all $m > m_h$, all $t \in \mathcal{R}_{\lambda, q, \delta}$. For every $t \in \mathcal{T}_{\lambda, q, \delta, r_0}$ we have $|t| < r_0 < T|\lambda||q|^{-1/2}$. Using (78), we obtain that

$$\sum_{m>m_h} \left(\frac{|t|}{T|\lambda||q|^{1/2}} \right)^m \leq \sum_{m>m_h} \left(\frac{1}{|q|} \right)^m = \frac{1}{1-|q|^{-1}} \frac{1}{|q|^{m_h+1}} \leq \frac{2|\lambda|A(h)}{1-|q|^{-1}},$$

hence

$$(79) \quad \sum_{m>m_h} \left| \frac{W_h(q^m \lambda)}{\Theta(q^m \lambda/t)} \right| \leq \frac{2|\lambda|C'K_0}{K_1\delta(1-|q|^{-1})} A(h)h! \left(\frac{1}{X} \right)^h |q|^{-h^2/4} \left(\frac{1}{|\lambda|} \right)^K |q|^{K(K-1)/2} |t|^K$$

for all $t \in \mathcal{T}_{\lambda, q, \delta, r_0}$.

In a second step, we give estimates for the sum $\sum_{m \leq m_h} W_h(q^m \lambda)/\Theta(q^m \lambda/t) - \sum_{n=0}^K f_{n,h} t^n$, where the $f_{n,h}$ are defined in the Taylor expansion (72). Taking into account (72) and (77), we can formally write, as we did in the proof of Proposition 1,

$$(80) \quad \sum_{m \leq m_h} W_h(q^m \lambda)/\Theta(q^m \lambda/t) - \sum_{n=0}^K f_{n,h} t^n = \sum_{m \leq m_h} \frac{1}{\Theta(q^m \lambda/t)} \left(\sum_{n \geq K+1} \frac{f_{n,h}}{q^{n(n-1)/2}} (q^m \lambda)^n \right) - \sum_{n=0}^K \frac{f_{n,h}}{q^{n(n-1)/2}} \left(\sum_{m > m_h} \frac{(q^m \lambda)^n}{\Theta(q^m \lambda/t)} \right)$$

for all $t \in \mathbb{C}^*$.

From (80) and (74), we deduce that

$$(81) \quad \left| \sum_{m \leq m_h} W_h(q^m \lambda)/\Theta(q^m \lambda/t) - \sum_{n=0}^K f_{n,h} t^n \right| \leq \mathcal{A}(t) + \mathcal{B}(t)$$

where

$$\mathcal{A}(t) = \sum_{m \leq m_h} \frac{1}{|\Theta(q^m \lambda/t)|} \left(\sum_{n \geq K+1} C(h) A(h)^n (|q|^m |\lambda|)^n \right)$$

and

$$\mathcal{B}(t) = \sum_{n=0}^K C(h) A(h)^n \left(\sum_{m > m_h} \frac{|(q^m \lambda)^n|}{|\Theta(q^m \lambda/t)|} \right),$$

for all $t \in \mathbb{C}^*$.

We give estimates for $\mathcal{A}(t)$. Taking into account (76), we have that

$$(82) \quad \mathcal{A}(t) \leq C(h) \sum_{m \leq m_h} \frac{1}{|\Theta(q^m \lambda/t)|} \frac{(A(h)|q|^m |\lambda|)^{K+1}}{1 - A(h)|q|^m |\lambda|} \leq \frac{C(h)A(h)^{K+1}}{1 - A(h)|q|^{m_h} |\lambda|} \sum_{m \geq -m_h} \frac{(|q|^{-m} |\lambda|)^{K+1}}{|\Theta(q^{-m} \lambda/t)|}$$

for all $t \in \mathbb{C}^*$. From (13), we have that

$$|\Theta(q^{-m} \lambda/t)| \geq K_1 \delta |q|^{-K(K-1)/2} \left| \frac{q^{-m} \lambda}{t} \right|^K$$

for all $m \geq -m_h$, all $t \in \mathcal{R}_{\lambda, q, \delta}$. We deduce that

$$(83) \quad \frac{(|q|^{-m} |\lambda|)^{K+1}}{|\Theta(q^{-m} \lambda/t)|} \leq \frac{|\lambda|}{K_1 \delta} |q|^{K(K-1)/2} |t|^K \left(\frac{1}{|q|} \right)^m$$

for all $m \geq -m_h$, all $t \in \mathcal{R}_{\lambda, q, \delta}$. From (82), (83) and (76), we conclude that

$$(84) \quad \begin{aligned} \mathcal{A}(t) &\leq \frac{C(h)A(h)}{1 - A(h)|q|^{m_h} |\lambda|} \frac{|\lambda|}{K_1 \delta} \frac{|q|^{m_h}}{1 - |q|^{-1}} A(h)^K |q|^{K(K-1)/2} |t|^K \\ &\leq \frac{C(h)}{K_1 \delta (1 - |q|^{-1})} A(h)^K |q|^{K(K-1)/2} |t|^K \end{aligned}$$

for all $t \in \mathcal{R}_{\lambda, q, \delta}$.

In the next step, we get estimates for $\mathcal{B}(t)$. From (13), we have that

$$|\Theta(q^m \lambda/t)| \geq K_1 \delta |q|^{-(K+1)K/2} \left| \frac{q^m \lambda}{t} \right|^{K+1}$$

for all $m > m_h$, all $t \in \mathcal{R}_{\lambda, q, \delta}$. We deduce that

$$\frac{|(q^m \lambda)^n|}{|\Theta(q^m \lambda/t)|} \leq \frac{|\lambda|^n}{K_1 \delta} \left(\frac{1}{|\lambda|} \right)^{K+1} |q|^{(K+1)K/2} |t|^{K+1} \left(\frac{1}{|q|^{K+1-n}} \right)^m$$

for all $m > m_h$, all $0 \leq n \leq K$. Then,

$$(85) \quad \sum_{m > m_h} \frac{|(q^m \lambda)^n|}{|\Theta(q^m \lambda/t)|} \leq \frac{|\lambda|^n}{K_1 \delta} \left(\frac{1}{|\lambda|} \right)^{K+1} |q|^{(K+1)K/2} |t|^{K+1} \frac{\left(\frac{1}{|q|^{K+1-n}} \right)^{m_h+1}}{1 - \frac{1}{|q|^{K+1-n}}}.$$

It is clear that $|q|^{K+1-n} \geq |q|$ for $0 \leq n \leq K$, hence

$$1 - \frac{1}{|q|^{K+1-n}} \geq 1 - |q|^{-1}, \quad 0 \leq n \leq K.$$

If we write

$$\left(\frac{1}{|q|^{K+1-n}} \right)^{m_h+1} = \left(\frac{1}{|q|^{m_h+1}} \right)^{K+1} (|q|^{m_h+1})^n,$$

from (85) we deduce that

$$\begin{aligned} \mathcal{B}(t) &\leq \frac{C(h)}{K_1 \delta (1 - |q|^{-1})} \left(\frac{1}{|\lambda| |q|^{m_h+1}} \right)^{K+1} |q|^{(K+1)K/2} |t|^{K+1} \sum_{n=0}^K (A(h)|\lambda| |q|^{m_h+1})^n \\ &\leq \frac{C(h)}{K_1 \delta (1 - |q|^{-1})} \left(\frac{1}{|\lambda| |q|^{m_h+1}} \right)^{K+1} |q|^{(K+1)K/2} |t|^{K+1} \sum_{n=0}^K (2A(h)|\lambda| |q|^{m_h+1})^n. \end{aligned}$$

By (78) we know that $2A(h)|\lambda||q|^{m_h+1} \geq 1$. For every real number $x \geq 1$ we have

$$\sum_{n=0}^K x^n \leq \sum_{n=0}^K \binom{K}{n} x^n x^{K-n} = (2x)^K,$$

and we deduce that $\sum_{n=0}^K (2A(h)|\lambda||q|^{m_h+1})^n \leq (4A(h)|\lambda||q|^{m_h+1})^K$. Also, we have $|t| < |\lambda q^{-1/2}|T$ whenever $t \in \mathcal{T}_{\lambda,q,\delta,r_0}$, and $(K+1)K/2 = K + K(K-1)/2$. Gathering all these facts and using (78), we deduce that

$$(86) \quad \begin{aligned} \mathcal{B}(t) &\leq \frac{|t|C(h)}{K_1\delta(1-|q|^{-1})|\lambda||q|^{m_h+1}} (4A(h)|q|)^K |q|^{K(K-1)/2} |t|^K \\ &\leq \frac{2|\lambda|TA(h)C(h)}{K_1\delta|q|^{1/2}(1-|q|^{-1})} (4A(h)|q|)^K |q|^{K(K-1)/2} |t|^K \end{aligned}$$

for all $t \in \mathcal{T}_{\lambda,q,\delta,r_0}$.

Finally, using the estimates

$$\begin{aligned} \left| \sum_{m \in \mathbb{Z}} W_h(q^m \lambda) / \Theta(q^m \lambda / t) - \sum_{n=0}^K f_{n,h} t^n \right| &\leq \left| \sum_{m \leq m_h} W_h(q^m \lambda) / \Theta(q^m \lambda / t) - \sum_{n=0}^K f_{n,h} t^n \right| \\ &\quad + \left| \sum_{m > m_h} W_h(q^m \lambda) / \Theta(q^m \lambda / t) \right| \end{aligned}$$

we deduce from (79), (81), (84), (86) that

$$(87) \quad \left| \mathcal{L}_q^\lambda(W_h)(t) - \sum_{n=0}^K f_{n,h} t^n \right| \leq D_1(h) B_1(h)^K |q|^{K(K-1)/2} |t|^K$$

for all $K \geq 0$, for all $t \in \mathcal{T}_{\lambda,q,\delta,r_0}$, with

$$(88) \quad B_1(h) = B_1(h+1)^{r_1/r_2}, \quad D_1(h) = B_2(h+1)^{r_1/r_2} h! B_3^h |q|^{-h^2/4},$$

where B_1, B_2 and B_3 are positive constants that do not depend on h . In order to conclude, it suffices to write, for $K \geq 1$,

$$\left| \mathcal{L}_q^\lambda(W_h)(t) - \sum_{n=0}^{K-1} f_{n,h} t^n \right| \leq \left| \mathcal{L}_q^\lambda(W_h)(t) - \sum_{n=0}^K f_{n,h} t^n \right| + |f_{K,h} t^K|,$$

and take into account (87) and (74). According to the expressions (75) and (88), one obtains the estimates (73) with

$$(89) \quad B(h) = A_1(h+1)^{r_1/r_2}, \quad D(h) = A_2(h+1)^{r_1/r_2} h! A_3^h |q|^{-h^2/4},$$

where A_1, A_2 and A_3 are again positive constants that do not depend on h . \square

We are ready to obtain our main result.

Theorem 5 *Suppose $\hat{X}_j(t) = \sum_{m \geq 0} f_{m,j} t^m \in \mathbb{C}[[t]]$, $0 \leq j \leq S-1$, are given initial conditions for the Cauchy problem (3), (4), and let*

$$\hat{X}(t, z) = \sum_{h \geq 0} \hat{X}_h(t) \frac{z^h}{h!} = \sum_{h \geq 0} \sum_{m \geq 0} f_{m,h} t^m \frac{z^h}{h!}$$

be the only formal series solution of the problem (see Lemma 5). We suppose that the series $\hat{X}_j(t)$, $0 \leq j \leq S-1$, are q -Gevrey of order 1, and that their formal q -Borel transforms of order 1, $W_j(\tau) = \hat{\mathcal{B}}_q \hat{X}_j(\tau)$, which are holomorphic functions around 0, indeed satisfy the assumptions of Theorems 2 and 4. We also assume that the rest of hypotheses of Theorem 2 are satisfied. Let

$$W(\tau, z) = \sum_{h \geq 0} W_h(\tau) \frac{z^h}{h!}$$

be the solution of the Cauchy problem (49), (50), corresponding to the initial conditions W_j , $0 \leq j \leq S-1$. Then, we have that:

1) The function $X(t, z) = \sum_{h \geq 0} \mathcal{L}_q^\lambda(W_h)(t) \frac{z^h}{h!}$ is holomorphic in $\mathcal{T}_{\lambda, q, \delta, r_0} \times \mathbb{C}$.

2) The function $X(t, z)$ solves the Cauchy problem (3), (4).

3) If $r_1 \geq 1$, given $R > 0$ there exist constants $\tilde{C} > 0$, $\tilde{D} > 0$ such that for every $n \in \mathbb{N}$, $n \geq 1$, one has

$$(90) \quad \left| X(t, z) - \sum_{h \geq 0} \sum_{m=0}^{n-1} f_{m,h} t^m \frac{z^h}{h!} \right| \leq \tilde{C} \tilde{D}^n \Gamma\left(\frac{r_1}{r_2}(n+1)\right) |q|^{n(n-1)/2} |t|^n$$

for every $t \in \mathcal{T}_{\lambda, q, \delta, r_0}$, $z \in D(0, R)$.

If $r_1 = 0$, given $R > 0$ there exist constants $\tilde{C} > 0$, $\tilde{D} > 0$ such that for every $n \in \mathbb{N}$, $n \geq 1$, one has

$$(91) \quad \left| X(t, z) - \sum_{h \geq 0} \sum_{m=0}^{n-1} f_{m,h} t^m \frac{z^h}{h!} \right| \leq \tilde{C} \tilde{D}^n |q|^{n(n-1)/2} |t|^n$$

for every $t \in \mathcal{T}_{\lambda, q, \delta, r_0}$, $z \in D(0, R)$.

Remark: Due to the estimates (90) and (91), we may say that the function $X(t, z)$ admits the series $\sum_{h \geq 0} \sum_{m \geq 0} f_{m,h} t^m \frac{z^h}{h!}$ as q -asymptotic expansion of order 1 in t , uniformly for z in the compact subsets of \mathbb{C} . It may be noted that, because of the small divisors problem we have dealt with, a new factor appears in the estimates, in terms of the Eulerian Gamma function. The value r_1/r_2 may be thought of as a sub-order, or a second-level order, in the asymptotic expansion.

Proof 1) In view of (73), for $n = 1$, and (89) we have that

$$|\mathcal{L}_q^\lambda(W_h)(t) - f_{0,h}| \leq D(h)B(h)|t| \leq A_1(h+1)^{2r_1/r_2} A_2 h! A_3^h |q|^{-h^2/4} r_0$$

for every $h \geq 0$, every $t \in \mathcal{T}_{\lambda, q, \delta, r_0}$. On the other hand, by (74) we have that

$$|f_{0,h}| \leq C_1 \left(\frac{1}{X_1}\right)^h h! |q|^{-h^2/2}$$

for every $h \geq 0$. So, we conclude that there exist $A_4, A_5 > 0$ such that

$$|\mathcal{L}_q^\lambda(W_h)(t)| \leq A_4 A_5^h h! |q|^{-h^2/4}$$

for every $h \geq 0$, every $t \in \mathcal{T}_{\lambda, q, \delta, r_0}$. Then, for $z \in D(0, R)$ we have

$$\left| \sum_{h \geq 0} \mathcal{L}_q^\lambda(W_h)(t) \frac{z^h}{h!} \right| \leq \sum_{h \geq 0} A_4 (A_5 R)^h |q|^{-h^2/4} < \infty,$$

so that the series converges and the function it defines is holomorphic in $\mathcal{T}_{\lambda, q, \delta, r_0} \times \mathbb{C}$.

2) Since the series $\sum_{h \geq 0} W_h(\tau) \frac{z^h}{h!}$ is a solution of (49), (50), one can guarantee that $X(t, z)$ is a solution of the Cauchy problem (3), (4) by Proposition 2.

3) For every $n \geq 1$, every $(t, z) \in \mathcal{T}_{\lambda, q, \delta, r_0} \times D(0, R)$, the sum

$$\sum_{h \geq 0} \sum_{m=0}^{n-1} f_{m,h} t^m \frac{z^h}{h!}$$

is convergent, as we see from (74). One may take into account (73) and (89) and write

$$\begin{aligned} \left| X(t, z) - \sum_{h \geq 0} \sum_{m=0}^{n-1} f_{m,h} t^m \frac{z^h}{h!} \right| &\leq \sum_{h \geq 0} \left| \mathcal{L}_q^\lambda(W_h)(t) - \sum_{m=0}^{n-1} f_{m,h} t^m \right| \frac{R^h}{h!} \\ &\leq A_2 A_1^n |q|^{n(n-1)/2} |t|^n \sum_{h \geq 0} (h+1)^{r_1(n+1)/r_2} (A_3 R)^h |q|^{-h^2/4} \\ (92) \qquad \qquad \qquad &= \frac{A_2}{A_3 R} A_1^n |q|^{n(n-1)/2} |t|^n \sum_{h \geq 1} h^{r_1(n+1)/r_2} (A_3 R)^h |q|^{-(h-1)^2/4}. \end{aligned}$$

In case $r_1 = 0$, the conclusion easily follows, since the last sum is convergent and independent of n . In case $r_1 \geq 1$, we follow an idea of B. Braaksma and L. Stolovitch [5]. Let $\varepsilon > 0$, and let γ be a contour that goes from $\infty e^{-i\pi}$ to $-\varepsilon$ along the negative real axis, then it turns once around 0 in the positive sense, and it goes from $-\varepsilon$ to $\infty e^{i\pi}$ again along the negative real axis. For

$$(93) \qquad \qquad \qquad \mu = \frac{r_1(n+1)}{r_2} > 0,$$

Hankel's formula allows us to write

$$\frac{h^\mu}{\Gamma(\mu+1)} = \frac{1}{2\pi i} \int_\gamma e^{hs} s^{-\mu-1} ds,$$

so that the sum in (92) may be written as

$$\begin{aligned} (94) \quad \frac{\Gamma(\mu+1)}{2\pi i} \sum_{h \geq 1} (A_3 R)^h |q|^{-(h-1)^2/4} \int_\gamma e^{hs} s^{-\mu-1} ds \\ = \frac{\Gamma(\mu+1)}{2\pi i} \sum_{h \geq 1} \int_\gamma s^{-\mu-1} |q|^{-(h-1)^2/4} (A_3 R e^s)^h ds. \end{aligned}$$

We consider now the entire function

$$F(z) = \sum_{h \geq 1} |q|^{-(h-1)^2/4} z^h, \quad z \in \mathbb{C}.$$

The series converges uniformly in every closed disc. Observe that as s runs over γ , its real part remains bounded above, and the same is valid for the modulus of A_3Re^s . So, we may write

$$F(A_3Re^s) = \sum_{h \geq 1} |q|^{-(h-1)^2/4} (A_3Re^s)^h$$

uniformly in γ , and the dominated convergence theorem ensures that

$$(95) \quad \sum_{h \geq 1} \int_{\gamma} s^{-\mu-1} |q|^{-(h-1)^2/4} (A_3Re^s)^h ds = \int_{\gamma} s^{-\mu-1} \sum_{h \geq 1} |q|^{-(h-1)^2/4} (A_3Re^s)^h ds \\ = \int_{\gamma} s^{-\mu-1} F(A_3Re^s) ds.$$

Moreover, $F(A_3Re^s)$ remains bounded as s runs over γ , say by $M > 0$, and it is easy to obtain, estimating on each of the three parts of γ , that

$$(96) \quad \left| \int_{\gamma} s^{-\mu-1} F(A_3Re^s) ds \right| \leq 2 \frac{M}{\mu \varepsilon^{\mu}} + \frac{2\pi M}{\varepsilon^{\mu}} \leq \frac{\tilde{M}^{\mu}}{\mu \varepsilon^{\mu}},$$

where $\tilde{M} > 0$ is some suitable constant independent of h . Gathering (92), (94), (95) and (96), we see that

$$\left| X(t, z) - \sum_{h \geq 0} \sum_{m=0}^{n-1} f_{m,h} t^m \frac{z^h}{h!} \right| \leq \frac{A_2}{A_3 R} A_1^n |q|^{n(n-1)/2} |t|^n \frac{\Gamma(\mu+1)}{2\pi i} \frac{\tilde{M}^{\mu}}{\mu \varepsilon^{\mu}}.$$

It suffices to recall that $\Gamma(\mu+1) = \mu\Gamma(\mu)$ and the definition of μ , (93), in order to conclude. \square

Remark: All the results in this work are valid for any $r_1 \geq 0$, but the case $r_1 = 0$, as it may be seen in the last Theorem, deserves some attention, since the Fuchsian singularity at $z = 0$ does not appear any more. The most important consequence of this fact is the disappearance of the small divisors phenomenon we had in general.

Moreover, the condition 1) in Theorem 2, concerning the argument of q and the set V , can be relaxed. Indeed, the estimates (53) hold if one assumes that there exists $\delta > 0$ such that

$$(97) \quad \text{dist}(V^{r_2} q^{r_2 \mathbb{Z}}, \{-1\}) > \delta,$$

where dist is the Euclidean distance between two sets in \mathbb{C} . For example, suppose V is such that there exist R_1, R_2 with

$$0 < R_1 \leq |x^{r_2}| \leq R_2$$

for all $x \in V$, and suppose that $R_2 < |q|R_1$ and there exists $j \in \mathbb{Z}$ such that

$$|q|^{r_2 j} R_2 < 1 < |q|^{r_2(j+1)} R_1.$$

Then, one can easily check that the condition (97) holds.

In Theorem 4 all the functions W_h are holomorphic in a common disc, say \bar{D} , and there exist constants $T_1, X_1 > 0$ such that

$$\sup_{\tau \in \bar{D}} |\partial^n W_j(\tau)| \leq C_1 \left(\frac{1}{T_1} \right)^n \left(\frac{1}{X_1} \right)^j n! j! |q|^{-j^2/2}$$

for every $n, j \geq 0$. The proof of Proposition 3 admits some simplification, and one obtains that

$$|\mathcal{L}_q^\lambda(W_h)(t) - \sum_{m=0}^{n-1} f_{m,h} t^m| \leq A_2 h! A_3^h |q|^{-h^2/4} A_1^n |q|^{n(n-1)/2} |t|^n,$$

for every $h \geq 0, n \geq 1$. Finally, no sub-order appears in the q -asymptotic expansion of the solution $X(t, z)$.

References

- [1] S. Alinhac, *Problèmes de Cauchy pour des opérateurs singuliers*, (French) Bull. Soc. Math. France 102 (1974), 289–315.
- [2] W. Balsler, From divergent power series to analytic functions. Theory and application of multisummable power series. Lecture Notes in Math. 1582. Springer-Verlag, Berlin, 1994.
- [3] W. Balsler, Formal power series and linear systems of meromorphic ordinary differential equations. Universitext. Springer-Verlag, New York, 2000.
- [4] A. Bove, J. Lewis, C. Parenti, Propagation of singularities for Fuchsian operators, Lecture Notes in Math. 984. Springer-Verlag, Berlin, 1983.
- [5] B. Braaksma, L. Stolovitch, *Small divisors and large multipliers*, Ann. Inst. Fourier (Grenoble) 57 (2007), no. 2, 603–628.
- [6] H. Chen, Z. Luo, H. Tahara, *Formal solutions of nonlinear first order totally characteristic type PDE with irregular singularity*, Ann. Inst. Fourier (Grenoble) 51 (2001), no. 6, 1599–1620.
- [7] O. Costin, Asymptotics and Borel summability. Chapman&Hall, CRC Monographs and Surveys in Pure and Applied Mathematics, 141. CRC Press, Boca Raton, FL, 2009.
- [8] L. Di Vizio, J.-P. Ramis, J. Sauloy, C. Zhang, *Équations aux q -différences*, Gaz. Math. no. 96 (2003), 20–49.
- [9] R. Gérard, H. Tahara, Singular nonlinear partial differential equations, Aspects of Mathematics. Friedr. Vieweg and Sohn, Braunschweig, 1996.
- [10] C.-L. Ho, *On the use of Mellin transform to a class of q -difference-differential equations*, Phys. Lett. A 268 (2000), no. 4-6, 217–223.
- [11] K. Igari, *Removable singularities of holomorphic solutions of linear partial differential equations*, J. Math. Soc. Japan 56 (2004), no. 1, 87–113.
- [12] T. Kato, *Asymptotic behavior of solutions of the functional differential equation $y'(x) = ay(\lambda x) + by(x)$* , Delay and functional differential equations and their applications (Proc. Conf., Park City, Utah, 1972), pp. 197–217. Academic Press, New York, 1972.
- [13] M. Kawagishi, T. Yamanaka, *On the Cauchy problem for non linear PDEs in the Gevrey class with shrinkings*, J. Math. Soc. Japan 54 (2002), no. 3, 649–677.
- [14] M. Kawagishi, T. Yamanaka, *The heat equation and the shrinking*, Electron. J. Differential Equations 2003, no. 97, 1–14.
- [15] S. Malek, *On Gevrey functions solutions of partial differential equations with Fuchsian and irregular singularities*, J. Dyn. Control Syst. 15 (2009), no. 2, 277–305.
- [16] S. Malek, *On complex singularity analysis for linear q -difference-differential equations*. J. Dyn. Control Syst. 15 (2009), no. 1, 83–98.
- [17] S. Malek, *On functional linear partial differential equations in Gevrey spaces of holomorphic functions*. Ann. Fac. Sci. Toulouse Math. (6) 16 (2007), no. 2, 285–302.

- [18] S. Malek, *On singularly perturbed partial integro-differential equations with irregular singularity*, J. Dyn. Control Syst. 13 (2007), no. 3, 419–449.
- [19] B. Malgrange, *Sommation des séries divergentes*, Expo. Math. 13 (1995), no. 2-3, 163–222.
- [20] T. Mandai, *The method of Frobenius to Fuchsian partial differential equations*, J. Math. Soc. Japan 52 (2000), no. 3, 645–672.
- [21] T. Mandai, *Existence and nonexistence of null-solutions for some non-Fuchsian partial differential operators with T -dependent coefficients*, Nagoya Math. J. 122 (1991), 115–137.
- [22] S. Ouchi, *Multisummability of formal solutions of some linear partial differential equations*, J. Differential Equations 185 (2002), no. 2, 513–549.
- [23] D. W. Pravica, M. J. Spurr, *Analytic continuation into the future*, Dynamical systems and differential equations (Wilmington, NC, 2002). Discrete Contin. Dyn. Syst. 2003, suppl., 709–716.
- [24] D. W. Pravica, M. J. Spurr, *Unique summing of formal power series solutions to advanced and delayed differential equations*, Discrete Contin. Dyn. Syst. 2005, suppl., 730–737.
- [25] J.-P. Ramis, *About the growth of entire functions solutions of linear algebraic q -difference equations*, Ann. Fac. Sci. Toulouse Math. (6) 1 (1992), no. 1, 53–94.
- [26] J.-P. Ramis, *Déviissage Gevrey*, Astérisque, 59-60 (1978), 173–204.
- [27] J.-P. Ramis, J. Sauloy, C. Zhang, *Local analytic classification of q -difference equations*, preprint ArXiv, 2009.
- [28] J.-P. Ramis, C. Zhang, *Développement asymptotique q -Gevrey et fonction thêta de Jacobi*. C. R. Math. Acad. Sci. Paris 335 (2002), no. 11, 899–902.
- [29] T. Yamanaka, M. Kawagishi, *A Cauchy-Kovalevskaja type theory in the Gevrey class for PDEs with shrinkings*, Nonlinear Anal. 64 (2006), 1860–1884.
- [30] H. Yamane, *Global Fuchsian Cauchy problem*, J. Math. Sci. Univ. Tokyo 7 (2000), no. 1, 147–162.
- [31] C. Zhang, *Une sommation discrète pour des équations aux q -différences linéaires et à coefficients analytiques: théorie générale et exemples*, (French) [Discrete summation for linear q -difference equations with analytic coefficients: general theory and examples] Differential equations and the Stokes phenomenon, 309–329, World Sci. Publ., River Edge, NJ, 2002.
- [32] O. Zubelevich, *An elliptic equation with perturbed p -Laplace operator*, J. Math. Anal. Appl. 328 (2007), no. 2, 1188–1195.