

On complex singularity analysis for some linear partial differential equations in \mathbb{C}^3

A. Lastra*, S. Malek†, C. Stenger,

University of Alcalá, Departamento de Matemáticas,
Ap. de Correos 20, E-28871 Alcalá de Henares (Madrid), Spain,

University of Lille 1, Laboratoire Paul Painlevé,
59655 Villeneuve d'Ascq cedex, France,

University of La Rochelle, Avenue Michel Crépeau,
17042 La Rochelle cedex France

alberto.lastra@uah.es

Stephane.Malek@math.univ-lille1.fr

catherine.stenger@univ-lr.fr

June, 3 2013

Abstract

We investigate the existence of local holomorphic solutions Y of linear partial differential equations in three complex variables whose coefficients are holomorphic on some polydisc in \mathbb{C}^2 outside some singular set Θ . The coefficients are written as linear combinations of powers of a solution X of some first order nonlinear partial differential equation following an idea we have initiated in a previous work [20]. The solutions Y are shown to develop singularities along Θ with estimates of exponential type depending on the growth's rate of X near the singular set. We construct these solutions with the help of series of functions with infinitely many variables which involve derivatives of all orders of X in one variable. Convergence and bounds estimates of these series are studied using a majorant series method which leads to an auxiliary functional equation that contains differential operators in infinitely many variables. Using a fixed point argument, we show that these functional equations actually have solutions in some Banach spaces of formal power series.

Key words: singular linear partial differential equations, linear partial differential equations with infinitely many variables, formal series with infinitely many variables, singularity analysis. 2000 MSC: 35C10, 35C20.

*The author is partially supported by the project MTM2012-31439 of Ministerio de Ciencia e Innovacion, Spain

†The author is partially supported by the french ANR-10-JCJC 0105 project and the PHC Polonium 2013 project No. 28217SG.

1 Introduction

In this paper, we study a family of linear partial differential equations of the form

$$(1) \quad \partial_w^S Y(t, z, w) = \sum_{k \in \mathcal{S}} (a_{1,k}(t, z, w) \partial_t \partial_w^k Y(t, z, w) \\ + a_{2,k}(t, z, w) \partial_z \partial_w^k Y(t, z, w) + a_{3,k}(t, z, w) \partial_w^k Y(t, z, w))$$

for given initial data $\partial_w^j Y(t, z, 0) = \varphi_j(t, z)$, $0 \leq j \leq S - 1$, where \mathcal{S} is a subset of \mathbb{N}^2 and S is an integer which satisfy the constraints (118). The coefficients $a_{m,k}(t, z, w)$ are holomorphic functions on some domain $(D(0, r)^2 \setminus \Theta) \times D(0, \bar{w})$ where Θ is some singular set of $D(0, r)^2$ (where $D(0, \delta)$ denotes the disc centered at 0 in \mathbb{C} with radius $\delta > 0$) and the initial data $\varphi_j(t, z)$ are assumed to be holomorphic functions on the polydisc $D(0, r)^2$.

In order to avoid cumbersome statements and tedious computations, the authors have chosen to restrict their study to equations (1) that involve at most first order derivatives with respect to t and z but the method proposed in this work can also be extended to higher order derivatives too.

In this work, we plan to construct holomorphic solutions of the problem (1) on $(D(0, r)^2 \setminus \Theta) \times D(0, \bar{w})$ and we will give precise growth's rate for these solutions near the singular set Θ of the coefficients $a_{m,k}(t, z, w)$ (Theorem 1).

There exists a huge literature on the study of complex singularities and analytic continuation of solutions to linear partial differential equations starting from the fundamental contributions of J. Leray in [15]. Many important results are known for singular initial data and concern equations with bounded holomorphic coefficients. In that context, the singularities of the solution are generally contained in characteristic hypersurfaces issued from the singular locus of the initial conditions. For meromorphic initial data, we may refer to [3], [8], [24], [25] and for more general ramified multivalued initial data, we may cite [9], [10], [26], [31]. In our framework, the initial data are assumed to be non singular and the coefficients of the equation now carry the singularities. To the best knowledge of the authors, few results have been worked out in that case. For instance, the research of so-called *fuchsian singularities* in the context of partial differential equations is widely developed, we provide [1], [2], [7], [22] as examples of references in this direction. It turns out that the situation we consider is actually close to a singular perturbation problem since the nature of the equation changes nearby the singular locus of it's coefficients.

This work is a continuation of our previous study [20]. In the paper [20], the authors focused on linear partial differential equations in \mathbb{C}^2 . They have constructed local holomorphic solutions with a careful study of their asymptotic behaviour near the singular locus of the initial data. These initial data were chosen to be polynomial in t, z and a function $u(t)$ satisfying some nonlinear differential equation of first order on some punctured disc $D(t_0, r) \setminus \{t_0\} \subset \mathbb{C}$ and owning an isolated singularity at t_0 which is either a pole or an algebraic branch point according to a result of P. Painlevé. Inspired by the classical *tanh method* introduced in [21], they have considered formal series solutions of the form

$$(2) \quad u(t, z) = \sum_{l \geq 0} u_l(t, z) (u(t))^l$$

where u_l are holomorphic functions on $D(t_0, r) \times D$ where $D \subset \mathbb{C}$ is a small disc centered at 0. They have given suitable conditions for these series to be well defined and holomorphic for t in a sector S with vertex t_0 and moreover as t tends to t_0 the solutions $u(t, z)$ are shown to

carry at most exponential bounds estimates of the form $C \exp(M|t - t_0|^{-\mu})$ for some constants $C, M, \mu > 0$.

In this work, the coefficients $a_{m,k}(t, z, w)$ are constructed as polynomials in some function $X(t, z)$ with holomorphic coefficients in (t, z, w) , where $X(t, z)$ is now assumed to solve some nonlinear partial differential equation of first order and is asked to be holomorphic on a domain $D(0, r)^2 \setminus \Theta$ and to be singular along the set Θ . The class of functions in which one can choose the coefficients $a_{m,k}(t, z, w)$ is quite large since it contains meromorphic and multivalued holomorphic functions in (t, z) (see the example of Section 2.1).

In our setting, one cannot achieve the goal only dealing with formal expansions involving the function $X(t, z)$ like (2) since the derivatives of $X(t, z)$ with respect to t or z cannot be expressed only in term of $X(t, z)$. In order to get suitable recursion formulas, it turns out that we need to deal with series expansions that take into account all the derivatives of $X(t, z)$ with respect to z . For this reason, the construction of the solutions will follow the one introduced in a recent work of H. Tahara and will involve Banach spaces of holomorphic functions with infinitely many variables.

In the paper [27], H. Tahara introduced a new equivalence problem connecting two given nonlinear partial differential equations of first order in the complex domain. He showed that the equivalence maps have to satisfy so called coupling equations which are nonlinear partial differential equations of first order but with infinitely many variables. It is worthwhile saying that within the framework of mathematical physics, spaces of functions of infinitely many variables play a fundamental role in the study of nonlinear integrable partial differential equations known as solitons equations as described in the theory of M. Sato. See [23] for an introduction.

The layout of the paper is as follows. In a first step described in Section 2.2, we construct formal series of the form

$$(3) \quad U(t, z, w) = \sum_{\alpha \geq 0} \phi_{\alpha}(t, z, (\frac{\partial_z^h X(t, z)}{h! \nu^h})_{0 \leq h \leq \alpha}) \frac{w^{\alpha}}{\alpha!},$$

solutions of some auxiliary non-homogeneous integro-differential equation (13) with polynomial coefficients in $X(t, z)$. The coefficients ϕ_{α} , $\alpha \geq 0$, are holomorphic functions on some polydisc in $\mathbb{C}^{\alpha+3}$ that satisfy some differential recursion (Proposition 1).

In Section 2.3, we establish a sequence of inequalities for the modulus of the differentials of arbitrary order of the functions ϕ_{α} denoted $\varphi_{\alpha, n_0, n_1, (l_h)_{0 \leq h \leq \alpha}}$ for all non-negative integers α, n_0, n_1, l_h with $0 \leq h \leq \alpha$ (Proposition 2). In the next section, we construct a sequence of coefficients $\psi_{\alpha, n_0, n_1, (l_h)_{0 \leq h \leq \alpha}}$ which is larger than the latter sequence

$$\varphi_{\alpha, n_0, n_1, (l_h)_{0 \leq h \leq \alpha}} \leq \psi_{\alpha, n_0, n_1, (l_h)_{0 \leq h \leq \alpha}}$$

for any non-negative integers α, n_0, n_1, l_h with $0 \leq h \leq \alpha$ and whose generating formal series satisfies some integro-differential functional equation (37) that involves differential operators with infinitely many variables (Propositions 3 and 4). The idea of considering recursions over the complete family of derivatives and the use of majorant series which lead to auxiliary Cauchy problems were already applied in former papers by the authors of this work, see [14], [17], [18], [19], [20].

In Section 3, we solve the functional equation (37) by applying a fixed point argument in some Banach space of formal series with infinitely many variables (Proposition 10). The definition of these Banach spaces (Definition 2) is inspired from formal series spaces introduced in our previous work [20]. The core of the proof is based on continuity properties of linear integro-differential operators in infinitely many variables explained in Section 3.1 and constitutes the most technical part of the paper.

Finally, in Section 4, we prove the main result of our work. Namely, we construct analytic functions $Y(t, z, w)$, solutions of (1) for the prescribed initial data, defined on sets $K \times D(0, \bar{w})$ for any compact set $K \subset D(0, r)^2 \setminus \Theta$ with precise bounds of exponential type in term of the maximum value of $|X(t, z)|$ over K (Theorem 1). The proof puts together all the constructions performed in the previous sections. More precisely, for some specific choice of the non-homogeneous term in the equation (13), a formal solution (3) of (13) gives rise to a formal solution $Y(t, z, w)$ of (1) with the given initial data that can be written as the sum of the integral $\partial_w^{-S} U(t, z, w)$ and a polynomial in w having the initial data φ_j as coefficients. Owing to the fact that the generating series of the sequence $\psi_{\alpha, n_0, n_1, (l_h)_{0 \leq h \leq \alpha}}$, solution of (37), belongs to the Banach spaces mentioned above, we get estimates for the holomorphic functions ϕ_α with precise bounds of exponential type in term of the radii of the polydiscs where they are defined, see (136). As a result, the formal solution $U(t, z, w)$ is actually convergent for w near the origin and for (t, z) belonging to any compact set of $D(0, r) \setminus \Theta$. Moreover, exponential bounds are achieved, see (137). The same properties then hold for $Y(t, z, w)$.

2 Formal series solutions of linear integro-differential equations

2.1 Some nonlinear partial differential equation

We consider the following nonlinear partial differential equation

$$(4) \quad \partial_t X(t, z) = a(t, z) \partial_z X(t, z) + \sum_{p=0}^d a_p(t, z) X^p(t, z)$$

where $d \geq 2$ is some integer, the coefficients $a(t, z)$, $a_p(t, z)$ are holomorphic functions on some polydisc $D(0, R')^2 \subset \mathbb{C}^2$ such that $a_d(t, z)$ is not identically equal to zero on $D(0, R')^2$.

Notice that the equation (4) can be solved by using the classical method of characteristics which is described in some classical textbooks like [4], p. 118 or [6], p. 100. However, the solutions of (4) cannot in general be expressed in closed form. Nevertheless, we can mention some general results concerning qualitative properties of holomorphic solutions to equation (4) and even to more general first order partial differential equations of the form

$$(5) \quad \partial_t u(t, x) = F(t, x, u(t, x), \partial_x u(t, x))$$

for $(t, x) \in \mathbb{C} \times \mathbb{C}^n$ where F is some holomorphic function and $n \geq 1$ an integer. For the construction of holomorphic functions to (5) with singularities located on some specific hypersurfaces (like $\{t = 0\}$), see [28] and [29]. For the existence of local multivalued holomorphic solutions ramified around some singular sets, we may refer to [11] and [13]. Concerning the study of the analytic continuation of singular solutions bounded on some hypersurface, we cite [30] and with prescribed upper estimates, we quote [12] and [16].

In this work, we make the assumption that (4) has a holomorphic solution $X(t, z)$ on $D(0, R')^2 \setminus \Theta$ where Θ is some set of $D(0, R')^2$ (Θ will be called a singular set in the sequel).

In the next example, we show that a large class of functions can be obtained as solutions of equations of the form (4).

Example : Let $n \geq 1$ be an integer and let $g : D(0, R')^2 \rightarrow \mathbb{C}$ be a holomorphic function which is not identically equal to zero. We consider

$$a_{n+1}(t, z) = \frac{1}{n} (\partial_z g(t, z) - \partial_t g(t, z))$$

which defines a holomorphic function on $D(0, R')^2$. Then, the function $X(t, z) = 1/(g(t, z))^{1/n}$ is a holomorphic solution of the equation

$$\partial_t X(t, z) = \partial_z X(t, z) + a_{n+1}(t, z)X^{n+1}(t, z)$$

on $D(0, R')^2 \setminus \Theta$ where Θ is the singular set defined by $\Theta = \{(t, z) \in D(0, R')^2 / g(t, z) \in L_\theta\}$ and L_θ is some halfline $\mathbb{R}_+ e^{i\theta}$ with $\theta \in \mathbb{R}$ depending on the choice of the determination of the logarithm.

2.2 Composition series

Let X be as in the previous subsection. In the following, we choose a compact subset K_0 with non-empty interior of $D(0, R')^2 \setminus \Theta$ for some $R < R'$ and we consider a real number $\rho > 1$ such that

$$\sup_{(t,z) \in K_0} |X(t, z)| \leq \rho/2.$$

Let $K \subsetneq K_0$ be a compact set with non-empty interior $\text{Int}(K)$. From the Cauchy formula, there exists a real number $\nu > 0$ such that

$$(6) \quad \sup_{(t,z) \in \text{Int}(K)} \frac{|\partial_z^h X(t, z)|}{h! \nu^h} \leq \rho/2$$

for all integers $h \geq 0$. For all integers $\alpha \geq 0$, we denote $I(\alpha) = \{0, \dots, \alpha\}$. We consider a sequence of functions $\phi_\alpha(v_0, v_1, (u_h)_{h \in I(\alpha)})$ which are holomorphic and bounded on the polydisc $D(0, R)^2 \prod_{h \in I(\alpha)} D(0, \rho)$, for all $\alpha \geq 0$.

We define the formal series in the w variable,

$$(7) \quad U(t, z, w) = \sum_{\alpha \geq 0} \phi_\alpha(t, z, (\frac{\partial_z^h X(t, z)}{h! \nu^h})_{h \in I(\alpha)}) \frac{w^\alpha}{\alpha!}.$$

For all $\alpha \geq 0$, we consider a holomorphic and bounded function $\tilde{\omega}_\alpha(v_0, v_1, (u_h)_{h \in I(\alpha)})$ on the product $D(0, R')^2 \prod_{h \in I(\alpha)} D(0, \rho)$. We define the formal series

$$(8) \quad \tilde{\omega}(t, z, w) = \sum_{\alpha \geq 0} \tilde{\omega}_\alpha(t, z, (\frac{\partial_z^h X(t, z)}{h! \nu^h})_{h \in I(\alpha)}) \frac{w^\alpha}{\alpha!}.$$

Let \mathcal{S} be a finite subset of \mathbb{N} and let $S \geq 1$ be an integer which satisfies the property that

$$(9) \quad S > k$$

for all $k \in \mathcal{S}$. For all $k \in \mathcal{S}$, $m = 1, 2, 3$ and all integers $\alpha \geq 0$, we define a function $b_{m,k,\alpha}(t, z, u_0)$ which is holomorphic on $D(0, R')^2 \times \mathbb{C}$ and satisfies estimates of the form: there exist two constants $D_{m,k} > 0$, $\hat{D}_{m,k} > 0$ and an integer $d_{m,k} \geq 0$ such that

$$(10) \quad \sup_{|t| < R', |z| < R', |u_0| \leq \rho} |b_{m,k,\alpha}(t, z, u_0)| \leq D_{m,k} \rho^{d_{m,k}} \hat{D}_{m,k}^\alpha \alpha!$$

for all $\alpha \geq 0$, all $\rho \geq 1$. In particular, each function $u_0 \mapsto b_{m,k,\alpha}(t, z, u_0)$ is a polynomial of degree at most $d_{m,k}$ for all $(t, z) \in D(0, R')^2$. Finally, for all $k \in \mathcal{S}$, $m = 1, 2, 3$, we consider the series

$$(11) \quad b_{m,k}(t, z, u_0, w) = \sum_{\alpha \geq 0} b_{m,k,\alpha}(t, z, u_0) \frac{w^\alpha}{\alpha!}$$

which define holomorphic functions on $D(0, R')^2 \times \mathbb{C} \times D(0, \bar{w})$, for any $0 < \bar{w} \leq 1/\hat{D}_{m,k}$.

with

$$\begin{aligned} \partial_z(\phi_{\alpha_2+k-S}(t, z, (\frac{\partial^h X(t, z)}{h! \nu^h})_{h \in I(\alpha_2+k-S)})) &= (\partial_{v_1} \phi_{\alpha_2+k-S})(t, z, (\frac{\partial^h X(t, z)}{h! \nu^h})_{h \in I(\alpha_2+k-S)}) \\ &+ \sum_{j \in I(\alpha_2+k-S)} (j+1) \nu \frac{\partial_z^{j+1} X(t, z)}{(j+1)! \nu^{j+1}} (\partial_{u_j} \phi_{\alpha_2+k-S})(t, z, (\frac{\partial^h X(t, z)}{h! \nu^h})_{h \in I(\alpha_2+k-S)}), \end{aligned}$$

for all $(t, z) \in \text{Int}(K)$. We also get that

$$\begin{aligned} &b_{1,k}(t, z, X(t, z), w) \partial_t \partial_w^{-S+k} U(t, z, w) \\ &= \sum_{\alpha \geq 0} \left(\sum_{\alpha_1 + \alpha_2 = \alpha, \alpha_2 \geq S-k} \alpha! \frac{b_{1,k,\alpha_1}(t, z, X(t, z))}{\alpha_1!} \frac{\partial_t(\phi_{\alpha_2+k-S}(t, z, (\frac{\partial^h X(t, z)}{h! \nu^h})_{h \in I(\alpha_2+k-S)}))}{\alpha_2!} \right) \frac{w^\alpha}{\alpha!} \end{aligned}$$

with

$$\begin{aligned} \partial_t(\phi_{\alpha_2+k-S}(t, z, (\frac{\partial^h X(t, z)}{h! \nu^h})_{h \in I(\alpha_2+k-S)})) &= (\partial_{v_0} \phi_{\alpha_2+k-S})(t, z, (\frac{\partial^h X(t, z)}{h! \nu^h})_{h \in I(\alpha_2+k-S)}) \\ &+ \sum_{j \in I(\alpha_2+k-S)} \frac{\partial_t \partial_z^j X(t, z)}{j! \nu^j} (\partial_{u_j} \phi_{\alpha_2+k-S})(t, z, (\frac{\partial^h X(t, z)}{h! \nu^h})_{h \in I(\alpha_2+k-S)}), \end{aligned}$$

for all $(t, z) \in \text{Int}(K)$. Now, from (4) and the classical Schwarz's result on equality of mixed partial derivatives, we get that

$$\frac{\partial_t \partial_z^j X(t, z)}{j! \nu^j} = \frac{\partial_z^j \partial_t X(t, z)}{j! \nu^j} = \frac{1}{j! \nu^j} \partial_z^j (a(t, z) \partial_z X(t, z)) + \sum_{p=0}^d a_p(t, z) X^p(t, z)$$

and from the Leibniz formula, we can write

$$\frac{1}{j! \nu^j} \partial_z^j (a(t, z) \partial_z X(t, z)) = \sum_{l_1+l_2=j} \frac{\partial_z^{l_1} a(t, z)}{l_1! \nu^{l_1}} (l_2+1) \nu \frac{\partial_z^{l_2+1} X(t, z)}{(l_2+1)! \nu^{l_2+1}}$$

and

$$\frac{1}{j! \nu^j} \partial_z^j (a_p(t, z) X^p(t, z)) = \sum_{j_0+\dots+j_p=j} \frac{\partial_z^{j_0} a_p(t, z)}{j_0! \nu^{j_0}} \prod_{l=1}^p \frac{\partial_z^{j_l} X(t, z)}{j_l! \nu^{j_l}},$$

for all $(t, z) \in \text{Int}(K)$. Finally, gathering all the equalities above and using the recursion (12), one gets the integro-differential equation (13). \square

2.3 Recursion for the derivatives of the functions ϕ_α , $\alpha \geq 0$

We consider a sequence of functions $\phi_\alpha(v_0, v_1, (u_h)_{h \in I(\alpha)})$, $\alpha \geq 0$, which are holomorphic and bounded on some polydisc $D(0, R)^2 \prod_{h \in I(\alpha)} D(0, \rho)$ for some real numbers $R > 0$ and $\rho > 1$ and which satisfy the equalities (12). We introduce the sequences

$$(14) \quad \varphi_{\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}} = \sup_{|v_0| < R, |v_1| < R, |u_h| < \rho, h \in I(\alpha)} |\partial_{v_0}^{n_0} \partial_{v_1}^{n_1} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_h} \phi_\alpha(v_0, v_1, (u_h)_{h \in I(\alpha)})|$$

for all $n_0, n_1 \geq 0$, all $l_h \geq 0$, $h \in I(\alpha)$, for all $\alpha \geq 0$. We define also the following sequences

$$(15) \quad b_{m,k,\alpha,n_0,n_1,l_0} = \sup_{|v_0| < R, |v_1| < R, |u_0| < \rho} |\partial_{v_0}^{n_0} \partial_{v_1}^{n_1} \partial_{u_0}^{l_0} b_{m,k,\alpha}(v_0, v_1, u_0)|,$$

$$\tilde{\omega}_{\alpha,n_0,n_1,(l_h)_{h \in I(\alpha)}} = \sup_{|v_0| < R, |v_1| < R, |u_h| < \rho, h \in I(\alpha)} |\partial_{v_0}^{n_0} \partial_{v_1}^{n_1} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_h} \tilde{\omega}_{\alpha}(v_0, v_1, (u_h)_{h \in I(\alpha)})|$$

for $m = 1, 2, 3$ and $k \in \mathcal{S}$. We put

$$(16) \quad A_j(v_0, v_1, (u_h)_{h \in I(\alpha+1)}) = \sum_{l_1+l_2=j} \frac{\partial_{v_1}^{l_1} a(v_0, v_1)}{l_1! \nu^{l_1}} (l_2 + 1) \nu u_{l_2+1}$$

$$+ \sum_{p=0}^d \sum_{j_0+\dots+j_p=j} \frac{\partial_{v_1}^{j_0} a_p(v_0, v_1)}{j_0! \nu^{j_0}} \prod_{l=1}^p u_{j_l}$$

and

$$(17) \quad B_j(v_0, v_1, (u_h)_{h \in I(\alpha+1)}) = (j + 1) \nu u_{j+1}$$

for all $j \in I(\alpha)$, $v_0, v_1 \in D(0, R')$ and $u_h \in \mathbb{C}$, $h \in I(\alpha)$. We define the sequences

$$A_{j,\alpha,n_0,n_1,(l_h)_{h \in I(\alpha+1)}} = \sup_{|v_0| < R, |v_1| < R, |u_h| < \rho, h \in I(\alpha)} |\partial_{v_0}^{n_0} \partial_{v_1}^{n_1} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_h} A_j(v_0, v_1, (u_h)_{h \in I(\alpha+1)})|$$

and

$$B_{j,\alpha,n_0,n_1,(l_h)_{h \in I(\alpha+1)}} = \sup_{|v_0| < R, |v_1| < R, |u_h| < \rho, h \in I(\alpha)} |\partial_{v_0}^{n_0} \partial_{v_1}^{n_1} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_h} B_j(v_0, v_1, (u_h)_{h \in I(\alpha+1)})|$$

for all $j \in I(\alpha)$, all $n_0, n_1 \geq 0$, all $l_h \geq 0$, $h \in I(\alpha+1)$, for all $\alpha \geq 0$. We also recall the definition of the Kronecker symbol $\delta_{0,l}$ which is equal to 0 if $l \neq 0$ and equal to 1 if $l = 0$.

Proposition 2 *The sequence $\varphi_{\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}}$ satisfies the following inequality:*

$$\begin{aligned}
(18) \quad & \frac{\varphi_{\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}}}{\alpha!} \leq \sum_{k \in \mathcal{S}} \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ \alpha_2 \geq \mathcal{S} - k}} \sum_{\substack{n_{0,1} + n_{0,2} = n_0, n_{1,1} + n_{1,2} = n_1 \\ l_{h,1} + l_{h,2} = l_h, h \in I(\alpha)}} \frac{n_0! n_1! \prod_{h \in I(\alpha)} l_h!}{n_{0,1}! n_{0,2}! n_{1,1}! n_{1,2}! \prod_{h \in I(\alpha)} l_{h,1}! l_{h,2}!} \\
& \frac{b_{1,k, \alpha_1, n_{0,1}, n_{1,1}, l_{0,1}}}{\alpha_1!} \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_{h,1}} \times \frac{\varphi_{\alpha_2 + k - S, n_{0,2} + 1, n_{1,2}, (l_{h,2})_{h \in I(\alpha_2 + k - S)}}}{\alpha_2!} \prod_{h \in I(\alpha) \setminus I(\alpha_2 + k - S)} \delta_{0, l_{h,2}} \\
& + \sum_{j \in I(\alpha_2 + k - S)} \sum_{\substack{n_{0,1} + n_{0,2} + n_{0,3} = n_0, n_{1,1} + n_{1,2} + n_{1,3} = n_1 \\ l_{h,1} + l_{h,2} + l_{h,3} = l_h, h \in I(\alpha)}} \frac{n_0! n_1! \prod_{h \in I(\alpha)} l_h!}{n_{0,1}! n_{0,2}! n_{0,3}! n_{1,1}! n_{1,2}! n_{1,3}! \prod_{h \in I(\alpha)} l_{h,1}! l_{h,2}! l_{h,3}!} \\
& \frac{b_{1,k, \alpha_1, n_{0,1}, n_{1,1}, l_{0,1}}}{\alpha_1!} \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_{h,1}} \times A_{j, \alpha_2 + k - S + 1, n_{0,2}, n_{1,2}, (l_{h,2})_{h \in I(\alpha_2 + k - S + 1)}} \\
& \times \prod_{h \in I(\alpha) \setminus I(\alpha_2 + k - S + 1)} \delta_{0, l_{h,2}} \times \frac{\varphi_{\alpha_2 + k - S, n_{0,3}, n_{1,3}, (l_{h,3})_{h \in I(\alpha_2 + k - S), h \neq j}, l_{j,3} + 1}}{\alpha_2!} \times \prod_{h \in I(\alpha) \setminus I(\alpha_2 + k - S)} \delta_{0, l_{h,3}} \\
& + \sum_{k \in \mathcal{S}} \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ \alpha_2 \geq \mathcal{S} - k}} \sum_{\substack{n_{0,1} + n_{0,2} = n_0, n_{1,1} + n_{1,2} = n_1 \\ l_{h,1} + l_{h,2} = l_h, h \in I(\alpha)}} \frac{n_0! n_1! \prod_{h \in I(\alpha)} l_h!}{n_{0,1}! n_{0,2}! n_{1,1}! n_{1,2}! \prod_{h \in I(\alpha)} l_{h,1}! l_{h,2}!} \\
& \frac{b_{2,k, \alpha_1, n_{0,1}, n_{1,1}, l_{0,1}}}{\alpha_1!} \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_{h,1}} \times \frac{\varphi_{\alpha_2 + k - S, n_{0,2}, n_{1,2} + 1, (l_{h,2})_{h \in I(\alpha_2 + k - S)}}}{\alpha_2!} \prod_{h \in I(\alpha) \setminus I(\alpha_2 + k - S)} \delta_{0, l_{h,2}} \\
& + \sum_{j \in I(\alpha_2 + k - S)} \sum_{\substack{n_{0,1} + n_{0,2} + n_{0,3} = n_0, n_{1,1} + n_{1,2} + n_{1,3} = n_1 \\ l_{h,1} + l_{h,2} + l_{h,3} = l_h, h \in I(\alpha)}} \frac{n_0! n_1! \prod_{h \in I(\alpha)} l_h!}{n_{0,1}! n_{0,2}! n_{0,3}! n_{1,1}! n_{1,2}! n_{1,3}! \prod_{h \in I(\alpha)} l_{h,1}! l_{h,2}! l_{h,3}!} \\
& \frac{b_{2,k, \alpha_1, n_{0,1}, n_{1,1}, l_{0,1}}}{\alpha_1!} \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_{h,1}} \times B_{j, \alpha_2 + k - S + 1, n_{0,2}, n_{1,2}, (l_{h,2})_{h \in I(\alpha_2 + k - S + 1)}} \\
& \times \prod_{h \in I(\alpha) \setminus I(\alpha_2 + k - S + 1)} \delta_{0, l_{h,2}} \times \frac{\varphi_{\alpha_2 + k - S, n_{0,3}, n_{1,3}, (l_{h,3})_{h \in I(\alpha_2 + k - S), h \neq j}, l_{j,3} + 1}}{\alpha_2!} \times \prod_{h \in I(\alpha) \setminus I(\alpha_2 + k - S)} \delta_{0, l_{h,3}} \\
& + \sum_{k \in \mathcal{S}} \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ \alpha_2 \geq \mathcal{S} - k}} \sum_{\substack{n_{0,1} + n_{0,2} = n_0, n_{1,1} + n_{1,2} = n_1 \\ l_{h,1} + l_{h,2} = l_h, h \in I(\alpha)}} \frac{n_0! n_1! \prod_{h \in I(\alpha)} l_h!}{n_{0,1}! n_{0,2}! n_{1,1}! n_{1,2}! \prod_{h \in I(\alpha)} l_{h,1}! l_{h,2}!} \\
& \frac{b_{3,k, \alpha_1, n_{0,1}, n_{1,1}, l_{0,1}}}{\alpha_1!} \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_{h,1}} \times \frac{\varphi_{\alpha_2 + k - S, n_{0,2}, n_{1,2}, (l_{h,2})_{h \in I(\alpha_2 + k - S)}}}{\alpha_2!} \prod_{h \in I(\alpha) \setminus I(\alpha_2 + k - S)} \delta_{0, l_{h,2}} \\
& + \frac{\tilde{\omega}_{\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}}}{\alpha!}
\end{aligned}$$

for all $\alpha \geq 0$, all $n_0, n_1, l_h \geq 0$ for $h \in I(\alpha)$.

Proof In order to get the inequality (18), we apply the differential operator $\partial_{v_0}^{n_0} \partial_{v_1}^{n_1} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_h}$ on the left and right handside of the recursion (12) and we use the expansions that are computed below.

From the Leibniz formula, we deduce that

$$\begin{aligned}
(19) \quad & \partial_{v_0}^{n_0} \partial_{v_1}^{n_1} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_h} (b_{3,k, \alpha_1}(v_0, v_1, u_0) \phi_{\alpha_2 + k - S}(v_0, v_1, (u_h)_{h \in I(\alpha_2 + k - S)})) = \\
& \sum_{\substack{n_{0,1} + n_{0,2} = n_0, n_{1,1} + n_{1,2} = n_1 \\ l_{h,1} + l_{h,2} = l_h, h \in I(\alpha)}} \frac{n_0! n_1! \prod_{h \in I(\alpha)} l_h!}{n_{0,1}! n_{0,2}! n_{1,1}! n_{1,2}! \prod_{h \in I(\alpha)} l_{h,1}! l_{h,2}!} \partial_{v_0}^{n_{0,1}} \partial_{v_1}^{n_{1,1}} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_{h,1}} (b_{3,k, \alpha_1}(v_0, v_1, u_0)) \\
& \times \partial_{v_0}^{n_{0,2}} \partial_{v_1}^{n_{1,2}} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_{h,2}} (\phi_{\alpha_2 + k - S}(v_0, v_1, (u_h)_{h \in I(\alpha_2 + k - S)}))
\end{aligned}$$

and

$$(20) \quad \partial_{v_0}^{n_0} \partial_{v_1}^{n_1} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_h} (b_{1,k,\alpha_1}(v_0, v_1, u_0) \partial_{v_0} \phi_{\alpha_2+k-S}(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S)})) =$$

$$\sum_{\substack{n_{0,1}+n_{0,2}=n_0, n_{1,1}+n_{1,2}=n_1 \\ l_{h,1}+l_{h,2}=l_h, h \in I(\alpha)}} \frac{n_0! n_1! \prod_{h \in I(\alpha)} l_h!}{n_{0,1}! n_{0,2}! n_{1,1}! n_{1,2}! \prod_{h \in I(\alpha)} l_{h,1}! l_{h,2}!} \partial_{v_0}^{n_{0,1}} \partial_{v_1}^{n_{1,1}} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_{h,1}} (b_{1,k,\alpha_1}(v_0, v_1, u_0))$$

$$\times \partial_{v_0}^{n_{0,2}+1} \partial_{v_1}^{n_{1,2}} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_{h,2}} (\phi_{\alpha_2+k-S}(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S)})).$$

Moreover, we can write

$$(21) \quad \partial_{v_0}^{n_0,1} \partial_{v_1}^{n_{1,1}} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_{h,1}} (b_{3,k,\alpha_1}(v_0, v_1, u_0))$$

$$= \partial_{v_0}^{n_0,1} \partial_{v_1}^{n_{1,1}} \partial_{u_0}^{l_{0,1}} b_{3,k,\alpha_1}(v_0, v_1, u_0) \times \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0,l_{h,1}}$$

with

$$(22) \quad \partial_{v_0}^{n_0,2} \partial_{v_1}^{n_{1,2}} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_{h,2}} (\phi_{\alpha_2+k-S}(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S)}))$$

$$= \partial_{v_0}^{n_0,2} \partial_{v_1}^{n_{1,2}} \prod_{h \in I(\alpha_2+k-S)} \partial_{u_h}^{l_{h,2}} (\phi_{\alpha_2+k-S}(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S)}))$$

$$\times \prod_{h \in I(\alpha) \setminus I(\alpha_2+k-S)} \delta_{0,l_{h,2}}$$

and

$$(23) \quad \partial_{v_0}^{n_0,1} \partial_{v_1}^{n_{1,1}} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_{h,1}} (b_{1,k,\alpha_1}(v_0, v_1, u_0))$$

$$= \partial_{v_0}^{n_0,1} \partial_{v_1}^{n_{1,1}} \partial_{u_0}^{l_{0,1}} b_{1,k,\alpha_1}(v_0, v_1, u_0) \times \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0,l_{h,1}}$$

with

$$(24) \quad \partial_{v_0}^{n_0,2+1} \partial_{v_1}^{n_{1,2}} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_{h,2}} (\phi_{\alpha_2+k-S}(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S)}))$$

$$= \partial_{v_0}^{n_0,2+1} \partial_{v_1}^{n_{1,2}} \prod_{h \in I(\alpha_2+k-S)} \partial_{u_h}^{l_{h,2}} (\phi_{\alpha_2+k-S}(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S)}))$$

$$\times \prod_{h \in I(\alpha) \setminus I(\alpha_2+k-S)} \delta_{0,l_{h,2}}.$$

By construction, we have

$$(25) \quad A_j(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S+1)}) = \sum_{l_1+l_2=j} \frac{\partial_{v_1}^{l_1} a(v_0, v_1)}{l_1! \nu^{l_1}} (l_2+1) \nu u_{l_2+1}$$

$$+ \sum_{p=0}^d \sum_{j_0+\dots+j_p=j} \frac{\partial_{v_1}^{j_0} a_p(v_0, v_1)}{j_0! \nu^{j_0}} \prod_{l=1}^p u_{j_l}$$

for all $j \in I(\alpha_2+k-S)$. Again, by the Leibniz formula, we get that

$$(26) \quad \partial_{v_0}^{n_0} \partial_{v_1}^{n_1} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_h} (b_{1,k,\alpha_1}(v_0, v_1, u_0) A_j(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S+1)}))$$

$$\times \partial_{u_j} \phi_{\alpha_2+k-S}(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S)})$$

$$= \sum_{\substack{n_{0,1}+n_{0,2}+n_{0,3}=n_0, n_{1,1}+n_{1,2}+n_{1,3}=n_1 \\ l_{h,1}+l_{h,2}+l_{h,3}=l_h, h \in I(\alpha)}} \frac{n_0! n_1! \prod_{h \in I(\alpha)} l_h!}{n_{0,1}! n_{0,2}! n_{0,3}! n_{1,1}! n_{1,2}! n_{1,3}! \prod_{h \in I(\alpha)} l_{h,1}! l_{h,2}! l_{h,3}!}$$

$$\partial_{v_0}^{n_{0,1}} \partial_{v_1}^{n_{1,1}} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_{h,1}} (b_{1,k,\alpha_1}(v_0, v_1, u_0))$$

$$\times \partial_{v_0}^{n_{0,2}} \partial_{v_1}^{n_{1,2}} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_{h,2}} (A_j(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S+1)}))$$

$$\times \partial_{v_0}^{n_{0,3}} \partial_{v_1}^{n_{1,3}} (\prod_{h \in I(\alpha), h \neq j} \partial_{u_h}^{l_{h,3}}) \partial_{u_j}^{l_{j,3}+1} \phi_{\alpha_2+k-S}(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S)}).$$

Inside the formula (26), we can rewrite the relations (23) and

$$(27) \quad \begin{aligned} & \partial_{v_0}^{n_0,2} \partial_{v_1}^{n_1,2} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_{h,2}} A_j(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S+1)}) \\ &= \partial_{v_0}^{n_0,2} \partial_{v_1}^{n_1,2} \prod_{h \in I(\alpha_2+k-S+1)} \partial_{u_h}^{l_{h,2}} A_j(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S+1)}) \times \prod_{h \in I(\alpha) \setminus I(\alpha_2+k-S+1)} \delta_{0,l_{h,2}} \end{aligned}$$

with

$$(28) \quad \begin{aligned} & \partial_{v_0}^{n_0,3} \partial_{v_1}^{n_1,3} (\prod_{h \in I(\alpha), h \neq j} \partial_{u_h}^{l_{h,3}}) \partial_{u_j}^{l_{j,3}+1} \phi_{\alpha_2+k-S}(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S)}) \\ &= \partial_{v_0}^{n_0,3} \partial_{v_1}^{n_1,3} (\prod_{h \in I(\alpha_2+k-S), h \neq j} \partial_{u_h}^{l_{h,3}}) \partial_{u_j}^{l_{j,3}+1} \phi_{\alpha_2+k-S}(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S)}) \\ & \quad \times \prod_{h \in I(\alpha) \setminus I(\alpha_2+k-S)} \delta_{0,l_{h,3}}. \end{aligned}$$

In the same way, one gets the next equalities

$$(29) \quad \begin{aligned} & \partial_{v_0}^{n_0} \partial_{v_1}^{n_1} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_h} (b_{2,k,\alpha_1}(v_0, v_1, u_0) \partial_{v_1} \phi_{\alpha_2+k-S}(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S)})) = \\ & \sum_{\substack{n_{0,1}+n_{0,2}=n_0, n_{1,1}+n_{1,2}=n_1 \\ l_{h,1}+l_{h,2}=l_h, h \in I(\alpha)}} \frac{n_0!n_1! \prod_{h \in I(\alpha)} l_h!}{n_{0,1}!n_{0,2}!n_{1,1}!n_{1,2}! \prod_{h \in I(\alpha)} l_{h,1}!l_{h,2}!} \partial_{v_0}^{n_{0,1}} \partial_{v_1}^{n_{1,1}} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_{h,1}} (b_{2,k,\alpha_1}(v_0, v_1, u_0)) \\ & \quad \times \partial_{v_0}^{n_{0,2}} \partial_{v_1}^{n_{1,2}+1} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_{h,2}} (\phi_{\alpha_2+k-S}(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S)})) \end{aligned}$$

with the factorizations

$$(30) \quad \begin{aligned} & \partial_{v_0}^{n_0,1} \partial_{v_1}^{n_1,1} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_{h,1}} (b_{2,k,\alpha_1}(v_0, v_1, u_0)) \\ &= \partial_{v_0}^{n_0,1} \partial_{v_1}^{n_1,1} \partial_{u_0}^{l_{0,1}} b_{2,k,\alpha_1}(v_0, v_1, u_0) \times \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0,l_{h,1}} \end{aligned}$$

and

$$(31) \quad \begin{aligned} & \partial_{v_0}^{n_0,2} \partial_{v_1}^{n_1,2+1} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_{h,2}} (\phi_{\alpha_2+k-S}(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S)})) \\ &= \partial_{v_0}^{n_0,2} \partial_{v_1}^{n_1,2+1} \prod_{h \in I(\alpha_2+k-S)} \partial_{u_h}^{l_{h,2}} (\phi_{\alpha_2+k-S}(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S)})) \\ & \quad \times \prod_{h \in I(\alpha) \setminus I(\alpha_2+k-S)} \delta_{0,l_{h,2}}. \end{aligned}$$

We recall that

$$(32) \quad B_j(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S+1)}) = (j+1)\nu u_{j+1}$$

for all $j \in I(\alpha_2+k-S)$ and we deduce that

$$(33) \quad \begin{aligned} & \partial_{v_0}^{n_0} \partial_{v_1}^{n_1} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_h} (b_{2,k,\alpha_1}(v_0, v_1, u_0) B_j(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S+1)})) \\ & \quad \times \partial_{u_j} \phi_{\alpha_2+k-S}(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S)}) \\ &= \sum_{\substack{n_{0,1}+n_{0,2}+n_{0,3}=n_0, n_{1,1}+n_{1,2}+n_{1,3}=n_1 \\ l_{h,1}+l_{h,2}+l_{h,3}=l_h, h \in I(\alpha)}} \frac{n_0!n_1! \prod_{h \in I(\alpha)} l_h!}{n_{0,1}!n_{0,2}!n_{0,3}!n_{1,1}!n_{1,2}!n_{1,3}! \prod_{h \in I(\alpha)} l_{h,1}!l_{h,2}!l_{h,3}!} \\ & \quad \partial_{v_0}^{n_{0,1}} \partial_{v_1}^{n_{1,1}} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_{h,1}} (b_{2,k,\alpha_1}(v_0, v_1, u_0)) \\ & \quad \times \partial_{v_0}^{n_{0,2}} \partial_{v_1}^{n_{1,2}} \prod_{h \in I(\alpha)} \partial_{u_h}^{l_{h,2}} (B_j(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S+1)})) \\ & \quad \times \partial_{v_0}^{n_{0,3}} \partial_{v_1}^{n_{1,3}} (\prod_{h \in I(\alpha), h \neq j} \partial_{u_h}^{l_{h,3}}) \partial_{u_j}^{l_{j,3}+1} \phi_{\alpha_2+k-S}(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S)}). \end{aligned}$$

Inside the formula (33), we can rewrite the relations (30) and

$$(34) \quad \begin{aligned} & \partial_{v_0}^{n_0,2} \partial_{v_1}^{n_1,2} \Pi_{h \in I(\alpha)} \partial_{u_h}^{l_h,2} B_j(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S+1)}) \\ &= \partial_{v_0}^{n_0,2} \partial_{v_1}^{n_1,2} \Pi_{h \in I(\alpha_2+k-S+1)} \partial_{u_h}^{l_h,2} B_j(v_0, v_1, (u_h)_{h \in I(\alpha_2+k-S+1)}) \times \Pi_{h \in I(\alpha) \setminus I(\alpha_2+k-S+1)} \delta_{0,l_h,2} \end{aligned}$$

with the factorization (28). \square

2.4 Majorant series and a functional equation with infinitely many variables

Definition 1 We denote $\mathbb{G}[[V_0, V_1, (U_h)_{h \geq 0}, W]]$ the vector space of formal series in the variables $V_0, V_1, (U_h)_{h \geq 0}, W$ of the form

$$(35) \quad \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) = \sum_{\alpha \geq 0} \Psi_\alpha(V_0, V_1, (U_h)_{h \in I(\alpha)}) \frac{W^\alpha}{\alpha!}$$

where $\Psi_\alpha \in \mathbb{C}[[V_0, V_1, (U_h)_{h \in I(\alpha)}]]$ for all $\alpha \geq 0$.

We keep the notations of the previous section and we introduce the following formal series:

$$(36) \quad \begin{aligned} B_{m,k}(V_0, V_1, U_0, W) &= \sum_{\alpha \geq 0} \left(\sum_{n_0, n_1, l_0 \geq 0} b_{m,k,\alpha, n_0, n_1, l_0} \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \frac{U_0^{l_0}}{l_0!} \right) \frac{W^\alpha}{\alpha!}, \\ \tilde{\Omega}(V_0, V_1, (U_h)_{h \geq 0}, W) &= \sum_{\alpha \geq 0} \left(\sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \tilde{\omega}_{\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}} \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \Pi_{h \in I(\alpha)} \frac{U_h^{l_h}}{l_h!} \right) \frac{W^\alpha}{\alpha!} \end{aligned}$$

for $m = 1, 2, 3$, all $k \in \mathcal{S}$, and

$$\begin{aligned} \mathbf{A}_{j,\alpha}(V_0, V_1, (U_h)_{h \in I(\alpha)}) &= \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} A_{j,\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}} \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \Pi_{h \in I(\alpha)} \frac{U_h^{l_h}}{l_h!}, \\ \mathbf{B}_{j,\alpha}(V_0, V_1, (U_h)_{h \in I(\alpha)}) &= \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} B_{j,\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}} \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \Pi_{h \in I(\alpha)} \frac{U_h^{l_h}}{l_h!} \end{aligned}$$

for all $\alpha \geq 0$, all $j \in I(\alpha)$. We also introduce the following linear operators acting on $\mathbb{G}[[V_0, V_1, (U_h)_{h \geq 0}, W]]$. Let

$$\begin{aligned} & \mathbb{D}_{\mathbf{A}} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) \\ &= \sum_{\alpha \geq 0} \left(\sum_{j \in I(\alpha)} \mathbf{A}_{j,\alpha+1}(V_0, V_1, (U_h)_{h \in I(\alpha+1)}) (\partial_{U_j} \Psi_\alpha)(V_0, V_1, (U_h)_{h \in I(\alpha)}) \right) \frac{W^\alpha}{\alpha!}, \\ & \mathbb{D}_{\mathbf{B}} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) \\ &= \sum_{\alpha \geq 0} \left(\sum_{j \in I(\alpha)} \mathbf{B}_{j,\alpha+1}(V_0, V_1, (U_h)_{h \in I(\alpha+1)}) (\partial_{U_j} \Psi_\alpha)(V_0, V_1, (U_h)_{h \in I(\alpha)}) \right) \frac{W^\alpha}{\alpha!} \end{aligned}$$

for all $\Psi \in \mathbb{G}[[V_0, V_1, (U_h)_{h \geq 0}, W]]$. We stress the fact that although these operators act on $\mathbb{G}[[V_0, V_1, (U_h)_{h \geq 0}, W]]$ their image does not have to belong to this space.

Proposition 3 *A formal series*

$$\Psi(V_0, V_1, (U_h)_{h \geq 0}, W) = \sum_{\alpha \geq 0} \left(\sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \psi_{\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}} \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \prod_{h \in I(\alpha)} \frac{U_h^{l_h}}{l_h!} \right) \frac{W^\alpha}{\alpha!}$$

satisfies the following functional equation

$$(37) \quad \begin{aligned} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) = & \sum_{k \in \mathcal{S}} (B_{1,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \partial_{V_0} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) \\ & + B_{1,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \mathbb{D}_{\mathbf{A}} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W)) \\ + & \sum_{k \in \mathcal{S}} (B_{2,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \partial_{V_1} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) \\ & + B_{2,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \mathbb{D}_{\mathbf{B}} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W)) \\ + & \sum_{k \in \mathcal{S}} B_{3,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) \\ & + \tilde{\Omega}(V_0, V_1, (U_h)_{h \geq 0}, W) \end{aligned}$$

if and only if its coefficients $\psi_{\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}}$ satisfy the following recursion

$$\begin{aligned}
(38) \quad \frac{\psi_{\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}}}{\alpha!} &= \sum_{k \in \mathcal{S}} \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ \alpha_2 \geq S-k}} \sum_{\substack{n_{0,1} + n_{0,2} = n_0, n_{1,1} + n_{1,2} = n_1 \\ l_{h,1} + l_{h,2} = l_h, h \in I(\alpha)}} \frac{n_0! n_1! \prod_{h \in I(\alpha)} l_h!}{n_{0,1}! n_{0,2}! n_{1,1}! n_{1,2}! \prod_{h \in I(\alpha)} l_{h,1}! l_{h,2}!} \\
&\frac{b_{1,k, \alpha_1, n_{0,1}, n_{1,1}, l_{0,1}}}{\alpha_1!} \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_{h,1}} \times \frac{\psi_{\alpha_2 + k - S, n_{0,2} + 1, n_{1,2}, (l_{h,2})_{h \in I(\alpha_2 + k - S)}}}{\alpha_2!} \prod_{h \in I(\alpha) \setminus I(\alpha_2 + k - S)} \delta_{0, l_{h,2}} \\
&+ \sum_{j \in I(\alpha_2 + k - S)} \sum_{\substack{n_{0,1} + n_{0,2} + n_{0,3} = n_0, n_{1,1} + n_{1,2} + n_{1,3} = n_1 \\ l_{h,1} + l_{h,2} + l_{h,3} = l_h, h \in I(\alpha)}} \frac{n_0! n_1! \prod_{h \in I(\alpha)} l_h!}{n_{0,1}! n_{0,2}! n_{0,3}! n_{1,1}! n_{1,2}! n_{1,3}! \prod_{h \in I(\alpha)} l_{h,1}! l_{h,2}! l_{h,3}!} \\
&\frac{b_{1,k, \alpha_1, n_{0,1}, n_{1,1}, l_{0,1}}}{\alpha_1!} \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_{h,1}} \times A_{j, \alpha_2 + k - S + 1, n_{0,2}, n_{1,2}, (l_{h,2})_{h \in I(\alpha_2 + k - S + 1)}} \\
&\times \prod_{h \in I(\alpha) \setminus I(\alpha_2 + k - S + 1)} \delta_{0, l_{h,2}} \times \frac{\psi_{\alpha_2 + k - S, n_{0,3}, n_{1,3}, (l_{h,3})_{h \in I(\alpha_2 + k - S), h \neq j}, l_{j,3} + 1}}{\alpha_2!} \times \prod_{h \in I(\alpha) \setminus I(\alpha_2 + k - S)} \delta_{0, l_{h,3}} \\
&+ \sum_{k \in \mathcal{S}} \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ \alpha_2 \geq S-k}} \sum_{\substack{n_{0,1} + n_{0,2} = n_0, n_{1,1} + n_{1,2} = n_1 \\ l_{h,1} + l_{h,2} = l_h, h \in I(\alpha)}} \frac{n_0! n_1! \prod_{h \in I(\alpha)} l_h!}{n_{0,1}! n_{0,2}! n_{1,1}! n_{1,2}! \prod_{h \in I(\alpha)} l_{h,1}! l_{h,2}!} \\
&\frac{b_{2,k, \alpha_1, n_{0,1}, n_{1,1}, l_{0,1}}}{\alpha_1!} \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_{h,1}} \times \frac{\psi_{\alpha_2 + k - S, n_{0,2}, n_{1,2} + 1, (l_{h,2})_{h \in I(\alpha_2 + k - S)}}}{\alpha_2!} \prod_{h \in I(\alpha) \setminus I(\alpha_2 + k - S)} \delta_{0, l_{h,2}} \\
&+ \sum_{j \in I(\alpha_2 + k - S)} \sum_{\substack{n_{0,1} + n_{0,2} + n_{0,3} = n_0, n_{1,1} + n_{1,2} + n_{1,3} = n_1 \\ l_{h,1} + l_{h,2} + l_{h,3} = l_h, h \in I(\alpha)}} \frac{n_0! n_1! \prod_{h \in I(\alpha)} l_h!}{n_{0,1}! n_{0,2}! n_{0,3}! n_{1,1}! n_{1,2}! n_{1,3}! \prod_{h \in I(\alpha)} l_{h,1}! l_{h,2}! l_{h,3}!} \\
&\frac{b_{2,k, \alpha_1, n_{0,1}, n_{1,1}, l_{0,1}}}{\alpha_1!} \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_{h,1}} \times B_{j, \alpha_2 + k - S + 1, n_{0,2}, n_{1,2}, (l_{h,2})_{h \in I(\alpha_2 + k - S + 1)}} \\
&\times \prod_{h \in I(\alpha) \setminus I(\alpha_2 + k - S + 1)} \delta_{0, l_{h,2}} \times \frac{\psi_{\alpha_2 + k - S, n_{0,3}, n_{1,3}, (l_{h,3})_{h \in I(\alpha_2 + k - S), h \neq j}, l_{j,3} + 1}}{\alpha_2!} \times \prod_{h \in I(\alpha) \setminus I(\alpha_2 + k - S)} \delta_{0, l_{h,3}} \\
&+ \sum_{k \in \mathcal{S}} \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ \alpha_2 \geq S-k}} \sum_{\substack{n_{0,1} + n_{0,2} = n_0, n_{1,1} + n_{1,2} = n_1 \\ l_{h,1} + l_{h,2} = l_h, h \in I(\alpha)}} \frac{n_0! n_1! \prod_{h \in I(\alpha)} l_h!}{n_{0,1}! n_{0,2}! n_{1,1}! n_{1,2}! \prod_{h \in I(\alpha)} l_{h,1}! l_{h,2}!} \\
&\frac{b_{3,k, \alpha_1, n_{0,1}, n_{1,1}, l_{0,1}}}{\alpha_1!} \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_{h,1}} \times \frac{\psi_{\alpha_2 + k - S, n_{0,2}, n_{1,2}, (l_{h,2})_{h \in I(\alpha_2 + k - S)}}}{\alpha_2!} \prod_{h \in I(\alpha) \setminus I(\alpha_2 + k - S)} \delta_{0, l_{h,2}} \\
&+ \frac{\tilde{\omega}_{\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}}}{\alpha!}
\end{aligned}$$

for all $\alpha \geq 0$, all $n_0, n_1, l_h \geq 0$ with $h \in I(\alpha)$.

Proof We proceed by identification of the coefficients in the Taylor expansion with respect to the variables $V_0, V_1, (U_h)_{h \in I(\alpha)}$ and W for all $\alpha \geq 0$. By definition, we have that

$$B_{1,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \partial_{V_0} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) = \sum_{\alpha \geq 0} \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ \alpha_2 \geq S-k}} \mathcal{C}_{\alpha_1, \alpha_2}^1 W^\alpha$$

where the coefficients $\mathcal{C}_{\alpha_1, \alpha_2}^1$ can be rewritten, using the Kronecker symbol $\delta_{0,m}$, in the form

$$\begin{aligned} \mathcal{C}_{\alpha_1, \alpha_2}^1 = & \left(\sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \frac{b_{1,k, \alpha_1, n_0, n_1, l_0}}{\alpha_1!} \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_h} \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \prod_{h \in I(\alpha)} \frac{U_h^{l_h}}{l_h!} \right) \\ & \times \left(\sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \frac{\psi_{\alpha_2+k-S, n_0+1, n_1, (l_h)_{h \in I(\alpha_2+k-S)}}}{\alpha_2!} \prod_{h \in I(\alpha) \setminus I(\alpha_2+k-S)} \delta_{0, l_h} \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \prod_{h \in I(\alpha)} \frac{U_h^{l_h}}{l_h!} \right) \end{aligned}$$

Hence,

$$\begin{aligned} (39) \quad \mathcal{C}_{\alpha_1, \alpha_2}^1 = & \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \left(\sum_{\substack{n_{0,1}+n_{0,2}=n_0, n_{1,1}+n_{1,2}=n_1 \\ l_{h,1}+l_{h,2}=l_h, h \in I(\alpha)}} \frac{b_{1,k, \alpha_1, n_{0,1}, n_{1,1}, l_{0,1}}}{\alpha_1! n_{0,1}! n_{1,1}! \prod_{h \in I(\alpha)} l_{h,1}!} \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_{h,1}} \right. \\ & \times \left. \frac{\psi_{\alpha_2+k-S, n_{0,2}+1, n_{1,2}, (l_{h,2})_{h \in I(\alpha_2+k-S)}}}{\alpha_2! n_{0,2}! n_{1,2}! \prod_{h \in I(\alpha)} l_{h,2}!} \prod_{h \in I(\alpha) \setminus I(\alpha_2+k-S)} \delta_{0, l_{h,2}} \right) V_0^{n_0} V_1^{n_1} \prod_{h \in I(\alpha)} U_h^{l_h}. \end{aligned}$$

We also have that

$$B_{1,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \mathbb{D}_{\mathbf{A}} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) = \sum_{\alpha \geq 0} \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ \alpha_2 \geq S-k}} \mathcal{F}_{\alpha_1, \alpha_2}^1 W^\alpha$$

where the coefficients $\mathcal{F}_{\alpha_1, \alpha_2}^1$ can be rewritten in the form

$$\begin{aligned} \mathcal{F}_{\alpha_1, \alpha_2}^1 = & \sum_{j \in I(\alpha_2-S+k)} \left(\sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \frac{b_{1,k, \alpha_1, n_0, n_1, l_0}}{\alpha_1!} \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_h} \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \prod_{h \in I(\alpha)} \frac{U_h^{l_h}}{l_h!} \right) \\ & \times \left(\sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} A_{j, \alpha_2-S+k+1, n_0, n_1, (l_h)_{h \in I(\alpha_2-S+k+1)}} \prod_{h \in I(\alpha) \setminus I(\alpha_2-S+k+1)} \delta_{0, l_h} \right. \\ & \quad \times \left. \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \prod_{h \in I(\alpha)} \frac{U_h^{l_h}}{l_h!} \right) \\ & \times \left(\sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \frac{\psi_{\alpha_2-S+k, n_0, n_1, (l_h)_{h \in I(\alpha_2-S+k), h \neq j}, l_j+1}}{\alpha_2!} \prod_{h \in I(\alpha) \setminus I(\alpha_2-S+k)} \delta_{0, l_h} \right. \\ & \quad \times \left. \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \prod_{h \in I(\alpha)} \frac{U_h^{l_h}}{l_h!} \right) \end{aligned}$$

Therefore,

$$\begin{aligned} (40) \quad \mathcal{F}_{\alpha_1, \alpha_2}^1 = & \sum_{j \in I(\alpha_2-S+k)} \left(\sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \right. \\ & \left(\sum_{\substack{n_{0,1}+n_{0,2}+n_{0,3}=n_0, n_{1,1}+n_{1,2}+n_{1,3}=n_1 \\ l_{h,1}+l_{h,2}+l_{h,3}=l_h, h \in I(\alpha)}} \frac{b_{1,k, \alpha_1, n_{0,1}, n_{1,1}, l_{0,1}}}{\alpha_1! n_{0,1}! n_{1,1}! \prod_{h \in I(\alpha)} l_{h,1}!} \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_{h,1}} \right. \\ & \times \frac{A_{j, \alpha_2-S+k+1, n_{0,2}, n_{1,2}, (l_{h,2})_{h \in I(\alpha_2-S+k+1)}}}{n_{0,2}! n_{1,2}! \prod_{h \in I(\alpha)} l_{h,2}!} \prod_{h \in I(\alpha) \setminus I(\alpha_2-S+k+1)} \delta_{0, l_{h,2}} \\ & \times \left. \frac{\psi_{\alpha_2-S+k, n_{0,3}, n_{1,3}, (l_{h,3})_{h \in I(\alpha_2-S+k), h \neq j}, l_{j,3}+1}}{\alpha_2! n_{0,3}! n_{1,3}! \prod_{h \in I(\alpha)} l_{h,3}!} \prod_{h \in I(\alpha) \setminus I(\alpha_2-S+k)} \delta_{0, l_{h,3}} \right) \\ & \left. \times V_0^{n_0} V_1^{n_1} \prod_{h \in I(\alpha)} U_h^{l_h} \right) \end{aligned}$$

On the other hand, using similar computations we get

$$B_{2,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \partial_{V_1} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) = \sum_{\alpha \geq 0} \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ \alpha_2 \geq S-k}} \mathcal{C}_{\alpha_1, \alpha_2}^2 W^\alpha$$

where

$$(41) \quad \mathcal{C}_{\alpha_1, \alpha_2}^2 = \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \left(\sum_{\substack{n_{0,1} + n_{0,2} = n_0, n_{1,1} + n_{1,2} = n_1 \\ l_{h,1} + l_{h,2} = l_h, h \in I(\alpha)}} \frac{b_{2,k,\alpha_1, n_{0,1}, n_{1,1}, l_{0,1}}}{\alpha_1! n_{0,1}! n_{1,1}! \prod_{h \in I(\alpha)} l_{h,1}!} \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_h, 1} \right. \\ \left. \times \frac{\psi_{\alpha_2+k-S, n_{0,2}, n_{1,2}+1, (l_h, 2)_{h \in I(\alpha_2+k-S)}}}{\alpha_2! n_{0,2}! n_{1,2}! \prod_{h \in I(\alpha)} l_{h,2}!} \prod_{h \in I(\alpha) \setminus I(\alpha_2+k-S)} \delta_{0, l_h, 2} \right) V_0^{n_0} V_1^{n_1} \prod_{h \in I(\alpha)} U_h^{l_h}.$$

We also have that

$$B_{2,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \mathbb{D}_{\mathbf{B}} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) = \sum_{\alpha \geq 0} \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ \alpha_2 \geq S-k}} \mathcal{F}_{\alpha_1, \alpha_2}^2 W^\alpha$$

where

$$(42) \quad \mathcal{F}_{\alpha_1, \alpha_2}^2 = \sum_{j \in I(\alpha_2 - S + k)} \left(\sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \left(\sum_{\substack{n_{0,1} + n_{0,2} + n_{0,3} = n_0, n_{1,1} + n_{1,2} + n_{1,3} = n_1 \\ l_{h,1} + l_{h,2} + l_{h,3} = l_h, h \in I(\alpha)}} \frac{b_{2,k,\alpha_1, n_{0,1}, n_{1,1}, l_{0,1}}}{\alpha_1! n_{0,1}! n_{1,1}! \prod_{h \in I(\alpha)} l_{h,1}!} \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_h, 1} \right. \right. \\ \left. \times \frac{B_{j, \alpha_2 - S + k + 1, n_{0,2}, n_{1,2}, (l_h, 2)_{h \in I(\alpha_2 - S + k + 1)}}}{n_{0,2}! n_{1,2}! \prod_{h \in I(\alpha)} l_{h,2}!} \prod_{h \in I(\alpha) \setminus I(\alpha_2 - S + k + 1)} \delta_{0, l_h, 2} \right. \\ \left. \times \frac{\psi_{\alpha_2 - S + k, n_{0,3}, n_{1,3}, (l_h, 3)_{h \in I(\alpha_2 - S + k), h \neq j}} \cdot l_{j,3} + 1}{\alpha_2! n_{0,3}! n_{1,3}! \prod_{h \in I(\alpha)} l_{h,3}!} \prod_{h \in I(\alpha) \setminus I(\alpha_2 - S + k)} \delta_{0, l_h, 3} \right) \\ \times V_0^{n_0} V_1^{n_1} \prod_{h \in I(\alpha)} U_h^{l_h}$$

and

$$B_{3,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) = \sum_{\alpha \geq 0} \sum_{\substack{\alpha_1 + \alpha_2 = \alpha \\ \alpha_2 \geq S-k}} \mathcal{C}_{\alpha_1, \alpha_2}^3 W^\alpha$$

where

$$(43) \quad \mathcal{C}_{\alpha_1, \alpha_2}^3 = \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \left(\sum_{\substack{n_{0,1} + n_{0,2} = n_0, n_{1,1} + n_{1,2} = n_1 \\ l_{h,1} + l_{h,2} = l_h, h \in I(\alpha)}} \frac{b_{3,k,\alpha_1, n_{0,1}, n_{1,1}, l_{0,1}}}{\alpha_1! n_{0,1}! n_{1,1}! \prod_{h \in I(\alpha)} l_{h,1}!} \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_h, 1} \right. \\ \left. \times \frac{\psi_{\alpha_2+k-S, n_{0,2}, n_{1,2}, (l_h, 2)_{h \in I(\alpha_2+k-S)}}}{\alpha_2! n_{0,2}! n_{1,2}! \prod_{h \in I(\alpha)} l_{h,2}!} \prod_{h \in I(\alpha) \setminus I(\alpha_2+k-S)} \delta_{0, l_h, 2} \right) V_0^{n_0} V_1^{n_1} \prod_{h \in I(\alpha)} U_h^{l_h}.$$

Finally, gathering the expansions (39), (40), (41) and (42) with (43) yields the result. \square

Proposition 4 *The sequences $\varphi_{\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}}$ and $\psi_{\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}}$ satisfy the following inequalities*

$$(44) \quad \varphi_{\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}} \leq \psi_{\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}}$$

for all $\alpha \geq 0$, all $n_0, n_1 \geq 0$, all $l_h \geq 0$, $h \in I(\alpha)$.

Proof For $\alpha = 0$, using the recursions (12) and (38), we get that

$$\varphi_{0,n_0,n_1,(l_h)_{h \in I(0)}} = \tilde{w}_{0,n_0,n_1,(l_h)_{h \in I(0)}} = \psi_{0,n_0,n_1,(l_h)_{h \in I(0)}}$$

for all $n_0, n_1, l_0 \geq 0$. By induction on α and using the inequalities (18) together with the equalities (38), one gets the result. \square

3 Convergent series solutions for a functional equation with infinitely many variables

3.1 Banach spaces of formal series

Let $\rho > 1$ and $\sigma, \bar{V}_0, \bar{V}_1, \bar{W}, \bar{\delta} > 0$ be real numbers. For any given real number $b > 1$, we define the sequences $r_b(\alpha) = \sum_{n=0}^{\alpha} 1/(n+1)^b$ for all $\alpha \geq 0$ and $\bar{U}_h = \bar{\delta}/(h^b + 1)$ for all $h \geq 0$.

Definition 2 Let $\alpha \geq 0$ be an integer. We denote $E_{\rho,\alpha,\bar{V}_0,\bar{V}_1,(\bar{U}_h)_{h \in I(\alpha)}}$ the vector space of formal series

$$\Psi(V_0, V_1, (U_h)_{h \in I(\alpha)}) = \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \psi_{n_0, n_1, (l_h)_{h \in I(\alpha)}} \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \prod_{h \in I(\alpha)} \frac{U_h^{l_h}}{l_h!}$$

that belong to $\mathbb{C}[[V_0, V_1, (U_h)_{h \in I(\alpha)}]]$ such that the series

$$\begin{aligned} & \|\Psi(V_0, V_1, (U_h)_{h \in I(\alpha)})\|_{\rho,\alpha,\bar{V}_0,\bar{V}_1,(\bar{U}_h)_{h \in I(\alpha)}} \\ &= \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \frac{|\psi_{n_0, n_1, (l_h)_{h \in I(\alpha)}}|}{\exp(\sigma r_b(\alpha) \rho)} \frac{\bar{V}_0^{n_0} \bar{V}_1^{n_1} \prod_{h \in I(\alpha)} \bar{U}_h^{l_h}}{(n_0 + n_1 + \sum_{h \in I(\alpha)} l_h + \alpha)!} \end{aligned}$$

is convergent. We denote also $G_{(\rho,\bar{V}_0,\bar{V}_1,(\bar{U}_h)_{h \geq 0}, \bar{W})}$ the vector space of formal series

$$\Psi(V_0, V_1, (U_h)_{h \geq 0}, W) = \sum_{\alpha \geq 0} \Psi_{\alpha}(V_0, V_1, (U_h)_{h \in I(\alpha)}) \frac{W^{\alpha}}{\alpha!}$$

where $\Psi_{\alpha}(V_0, V_1, (U_h)_{h \in I(\alpha)})$ belong to $E_{\rho,\alpha,\bar{V}_0,\bar{V}_1,(\bar{U}_h)_{h \in I(\alpha)}}$ for all $\alpha \geq 0$, such that the series

$$\|\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho,\bar{V}_0,\bar{V}_1,(\bar{U}_h)_{h \geq 0}, \bar{W})} = \sum_{\alpha \geq 0} \|\Psi_{\alpha}\|_{\rho,\alpha,\bar{V}_0,\bar{V}_1,(\bar{U}_h)_{h \in I(\alpha)}} \bar{W}^{\alpha}$$

is convergent. One checks that the space $G_{(\rho,\bar{V}_0,\bar{V}_1,(\bar{U}_h)_{h \geq 0}, \bar{W})}$ equipped with the norm $\|\cdot\|_{(\rho,\bar{V}_0,\bar{V}_1,(\bar{U}_h)_{h \geq 0}, \bar{W})}$ is a Banach space.

In the next two propositions, we study norm estimates for linear operators acting on the Banach spaces $E_{\rho,\alpha,\bar{V}_0,\bar{V}_1,(\bar{U}_h)_{h \in I(\alpha)}}$ constructed above.

Proposition 5 Consider a formal series

$$b(V_0, V_1, (U_h)_{h \in I(\alpha)}) = \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} b_{n_0, n_1, (l_h)_{h \in I(\alpha)}} \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \prod_{h \in I(\alpha)} \frac{U_h^{l_h}}{l_h!}$$

which is absolutely convergent on the polydisc $D(0, \bar{V}_0) \times D(0, \bar{V}_1) \times_{h \in I(\alpha)} D(0, \bar{U}_h)$. We use the notation

$$|b|(\bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}) = \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} |b_{n_0, n_1, (l_h)_{h \in I(\alpha)}}| \frac{\bar{V}_0^{n_0}}{n_0!} \frac{\bar{V}_1^{n_1}}{n_1!} \prod_{h \in I(\alpha)} \frac{\bar{U}_h^{l_h}}{l_h!}.$$

Let $\Psi(V_0, V_1, (U_h)_{h \in I(\alpha)})$ belonging to $E_{\rho, \alpha, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}}$. Then, the following inequality

$$(45) \quad \begin{aligned} & \|b(V_0, V_1, (U_h)_{h \in I(\alpha)}) \Psi(V_0, V_1, (U_h)_{h \in I(\alpha)})\|_{\rho, \alpha, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}} \\ & \leq |b|(\bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}) \|\Psi(V_0, V_1, (U_h)_{h \in I(\alpha)})\|_{\rho, \alpha, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}} \end{aligned}$$

holds.

Proof Let

$$\Psi(V_0, V_1, (U_h)_{h \in I(\alpha)}) = \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \psi_{n_0, n_1, (l_h)_{h \in I(\alpha)}} \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \prod_{h \in I(\alpha)} \frac{U_h^{l_h}}{l_h!}$$

which belongs to $E_{\rho, \alpha, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}}$. By definition, we have that

$$\begin{aligned} & \|b(V_0, V_1, (U_h)_{h \in I(\alpha)}) \Psi(V_0, V_1, (U_h)_{h \in I(\alpha)})\|_{\rho, \alpha, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}} \\ & = \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \left| \sum_{\substack{n_{0,1}+n_{0,2}=n_0, n_{1,1}+n_{1,2}=n_1 \\ l_{h,1}+l_{h,2}=l_h, h \in I(\alpha)}} \frac{n_0! n_1! \prod_{h \in I(\alpha)} l_h!}{n_{0,1}! n_{0,2}! n_{1,1}! n_{1,2}! \prod_{h \in I(\alpha)} l_{h,1}! l_{h,2}!} \right. \\ & \quad \left. b_{n_{0,1}, n_{1,1}, (l_{h,1})_{h \in I(\alpha)}} \psi_{n_{0,2}, n_{1,2}, (l_{h,2})_{h \in I(\alpha)}} \right| \frac{1}{\exp(\sigma r_b(\alpha) \rho)} \frac{\bar{V}_0^{n_0} \bar{V}_1^{n_1} \prod_{h \in I(\alpha)} \bar{U}_h^{l_h}}{(n_0 + n_1 + \sum_{h \in I(\alpha)} l_h + \alpha)!}. \end{aligned}$$

We can give upper bounds for this latter expression

$$(46) \quad \begin{aligned} & \|b(V_0, V_1, (U_h)_{h \in I(\alpha)}) \Psi(V_0, V_1, (U_h)_{h \in I(\alpha)})\|_{\rho, \alpha, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}} \\ & \leq \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \sum_{\substack{n_{0,1}+n_{0,2}=n_0, n_{1,1}+n_{1,2}=n_1 \\ l_{h,1}+l_{h,2}=l_h, h \in I(\alpha)}} \left(\frac{n_0! n_1! \prod_{h \in I(\alpha)} l_h!}{n_{0,2}! n_{1,2}! \prod_{h \in I(\alpha)} l_{h,2}!} \right. \\ & \quad \times \frac{(n_{0,2} + n_{1,2} + \sum_{h \in I(\alpha)} l_{h,2} + \alpha)!}{(n_0 + n_1 + \sum_{h \in I(\alpha)} l_h + \alpha)!} \left. \frac{|b_{n_{0,1}, n_{1,1}, (l_{h,1})_{h \in I(\alpha)}}| \bar{V}_0^{n_{0,1}} \bar{V}_1^{n_{1,1}} \prod_{h \in I(\alpha)} \bar{U}_h^{l_{h,1}}}{n_{0,1}! n_{1,1}! \prod_{h \in I(\alpha)} l_{h,1}!} \right. \\ & \quad \times \left. |\psi_{n_{0,2}, n_{1,2}, (l_{h,2})_{h \in I(\alpha)}}| \frac{1}{\exp(\sigma r_b(\alpha) \rho)} \frac{\bar{V}_0^{n_{0,2}} \bar{V}_1^{n_{1,2}} \prod_{h \in I(\alpha)} \bar{U}_h^{l_{h,2}}}{(n_{0,2} + n_{1,2} + \sum_{h \in I(\alpha)} l_{h,2} + \alpha)!} \right) \end{aligned}$$

Lemma 1 For all integers $\alpha, n_0, n_1 \geq 0$, all $l_h \geq 0$, all $0 \leq n_{0,2} \leq n_0$, all $0 \leq n_{1,2} \leq n_1$, all $0 \leq l_{h,2} \leq l_h$ for $h \in I(\alpha)$, we have that

$$(47) \quad \frac{n_0! n_1! \prod_{h \in I(\alpha)} l_h!}{n_{0,2}! n_{1,2}! \prod_{h \in I(\alpha)} l_{h,2}!} \frac{(n_{0,2} + n_{1,2} + \sum_{h \in I(\alpha)} l_{h,2} + \alpha)!}{(n_0 + n_1 + \sum_{h \in I(\alpha)} l_h + \alpha)!} \leq 1.$$

Proof For any integers $a \leq b$ and $\alpha \geq 0$ one has

$$(48) \quad \frac{(a + \alpha)!}{(b + \alpha)!} \leq \frac{a!}{b!}$$

by using the factorization $(a + \alpha)! = (a + \alpha)(a + \alpha - 1) \cdots (a + 1)a!$. Therefore, one gets the inequality

$$(49) \quad \frac{n_0!n_1!\prod_{h \in I(\alpha)} l_h!}{n_{0,2}!n_{1,2}!\prod_{h \in I(\alpha)} l_{h,2}!} \frac{(n_{0,2} + n_{1,2} + \sum_{h \in I(\alpha)} l_{h,2} + \alpha)!}{(n_0 + n_1 + \sum_{h \in I(\alpha)} l_h + \alpha)!} \\ \leq \frac{n_0!n_1!\prod_{h \in I(\alpha)} l_h!}{n_{0,2}!n_{1,2}!\prod_{h \in I(\alpha)} l_{h,2}!} \frac{(n_{0,2} + n_{1,2} + \sum_{h \in I(\alpha)} l_{h,2})!}{(n_0 + n_1 + \sum_{h \in I(\alpha)} l_h)!}$$

Now, from the identity $(A + B)^{n_0+n_1+\sum_{h \in I(\alpha)} l_h} = (A + B)^{n_0}(A + B)^{n_1} \times \prod_{h \in I(\alpha)} (A + B)^{l_h}$ and the binomial formula, we deduce that

$$\frac{n_0!n_1!\prod_{h \in I(\alpha)} l_h!}{n_{0,1}!n_{0,2}!n_{1,1}!n_{1,2}!\prod_{h \in I(\alpha)} l_{h,1}!l_{h,2}!} \\ \leq \frac{(n_0 + n_1 + \sum_{h \in I(\alpha)} l_h)!}{(n_{0,1} + n_{1,1} + \sum_{h \in I(\alpha)} l_{h,1})!(n_{0,2} + n_{1,2} + \sum_{h \in I(\alpha)} l_{h,2})!}$$

for all $n_{0,1} + n_{0,2} = n_0$, $n_{1,1} + n_{1,2} = n_1$, $l_{h,1} + l_{h,2} = l_h$. Therefore, we deduce that

$$(50) \quad \frac{n_0!n_1!\prod_{h \in I(\alpha)} l_h!}{n_{0,2}!n_{1,2}!\prod_{h \in I(\alpha)} l_{h,2}!} \frac{(n_{0,2} + n_{1,2} + \sum_{h \in I(\alpha)} l_{h,2} + \alpha)!}{(n_0 + n_1 + \sum_{h \in I(\alpha)} l_h + \alpha)!} \\ \leq \frac{n_{0,1}!n_{1,1}!\prod_{h \in I(\alpha)} l_{h,1}!}{(n_{0,1} + n_{1,1} + \sum_{h \in I(\alpha)} l_{h,1})!} \leq 1,$$

and the lemma follows from the inequalities (49), (50). \square

Finally, the inequality (45) follows from (46) and (47). \square

Proposition 6 *Let α, α' be integers such that $\alpha' \geq 0$ and $\alpha' + 1 < \alpha$. Let $j \in I(\alpha')$ and $k \in \{0, 1\}$. We have that*

$$(51) \quad \|\partial_{U_j} \Psi(V_0, V_1, (U_h)_{h \in I(\alpha')})\|_{\rho, \alpha, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}} \\ \leq \frac{\exp(-\sigma \rho \frac{\alpha - \alpha'}{(\alpha + 1)^b})}{\bar{U}_j \prod_{l=1}^{\alpha - \alpha' - 1} (\alpha - l + 1)} \|\Psi(V_0, V_1, (U_h)_{h \in I(\alpha')})\|_{\rho, \alpha', \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)'}}$$

$$(52) \quad \|\partial_{V_k} \Psi(V_0, V_1, (U_h)_{h \in I(\alpha')})\|_{\rho, \alpha, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}} \\ \leq \frac{\exp(-\sigma \rho \frac{\alpha - \alpha'}{(\alpha + 1)^b})}{\bar{V}_k \prod_{l=1}^{\alpha - \alpha' - 1} (\alpha - l + 1)} \|\Psi(V_0, V_1, (U_h)_{h \in I(\alpha')})\|_{\rho, \alpha', \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)'}}$$

and

$$(53) \quad \|\Psi(V_0, V_1, (U_h)_{h \in I(\alpha')})\|_{\rho, \alpha, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}} \\ \leq \frac{\exp(-\sigma \rho \frac{\alpha - \alpha'}{(\alpha + 1)^b})}{\prod_{l=1}^{\alpha - \alpha'} (\alpha - l + 1)} \|\Psi(V_0, V_1, (U_h)_{h \in I(\alpha')})\|_{\rho, \alpha', \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)'}}$$

for all $\Psi(V_0, V_1, (U_h)_{h \in I(\alpha')}) \in E_{\rho, \alpha', \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)'}}$.

Proof Let $\Psi(V_0, V_1, (U_h)_{h \in I(\alpha')}) \in E_{\rho, \alpha', \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha')}}$ that we write in the form

$$\begin{aligned} \Psi(V_0, V_1, (U_h)_{h \in I(\alpha')}) &= \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \psi_{n_0, n_1, (l_h)_{h \in I(\alpha')}} \prod_{h \in I(\alpha) \setminus I(\alpha')} \delta_{0, l_h} \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \prod_{h \in I(\alpha)} \frac{U_h^{l_h}}{l_h!}. \end{aligned}$$

By definition, we get that

$$\begin{aligned} & \|\partial_{U_j} \Psi(V_0, V_1, (U_h)_{h \in I(\alpha')})\|_{\rho, \alpha, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}} \\ &= \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \frac{|\psi_{n_0, n_1, (l_h)_{h \in I(\alpha')}, h \neq j, l_j + 1} \prod_{h \in I(\alpha) \setminus I(\alpha')} \delta_{0, l_h}|}{\exp(\sigma r_b(\alpha) \rho)} \frac{\bar{V}_0^{n_0} \bar{V}_1^{n_1} \prod_{h \in I(\alpha)} \bar{U}_h^{l_h}}{(n_0 + n_1 + \sum_{h \in I(\alpha)} l_h + \alpha)!}. \end{aligned}$$

We give upper bounds for this latter expression,

$$\begin{aligned} (54) \quad & \|\partial_{U_j} \Psi(V_0, V_1, (U_h)_{h \in I(\alpha')})\|_{\rho, \alpha, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}} \\ &= \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha')} \left(\frac{(n_0 + n_1 + \sum_{h \in I(\alpha'), h \neq j} l_h + l_j + 1 + \alpha)!}{(n_0 + n_1 + \sum_{h \in I(\alpha')} l_h + \alpha)!} \frac{1}{\bar{U}_j \exp(\sigma \rho (r_b(\alpha) - r_b(\alpha')))}) \right) \\ & \quad \times \frac{|\psi_{n_0, n_1, (l_h)_{h \in I(\alpha')}, h \neq j, l_j + 1}|}{\exp(\sigma r_b(\alpha') \rho)} \frac{\bar{V}_0^{n_0} \bar{V}_1^{n_1} \prod_{h \in I(\alpha'), h \neq j} \bar{U}_h^{l_h} \bar{U}_j^{l_j + 1}}{(n_0 + n_1 + \sum_{h \in I(\alpha'), h \neq j} l_h + l_j + 1 + \alpha)!} \end{aligned}$$

Lemma 2 *We have*

$$\begin{aligned} (55) \quad & \frac{(n_0 + n_1 + \sum_{h \in I(\alpha'), h \neq j} l_h + l_j + 1 + \alpha)!}{(n_0 + n_1 + \sum_{h \in I(\alpha')} l_h + \alpha)!} \frac{1}{\exp(\sigma \rho (r_b(\alpha) - r_b(\alpha')))}) \\ & \leq \frac{\exp(-\sigma \rho \frac{\alpha - \alpha'}{(\alpha + 1)^b})}{\prod_{l=1}^{\alpha - \alpha' - 1} (\alpha - l + 1)} \end{aligned}$$

Proof We notice that

$$r_b(\alpha) - r_b(\alpha') = \sum_{n=\alpha'+1}^{\alpha} \frac{1}{(n+1)^b} \geq \frac{\alpha - \alpha'}{(\alpha + 1)^b}$$

and, with the help of (48), that for all integers $a \geq 0$,

$$\frac{(a + 1 + \alpha')!}{(a + \alpha)!} \leq \frac{1}{\prod_{l=1}^{\alpha - \alpha' - 1} (\alpha - l + 1)}.$$

The lemma follows. \square

We get that the inequality (51) follows from (54) together with (55). Finally, using similar arguments, one gets also the inequalities (52) and (53). \square

In the next two propositions, we study norm estimates for linear operators acting on the Banach space $G_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$.

Proposition 7 *Let a formal series $b(V_0, V_1, U_0, W) \in \mathbb{C}[[V_0, V_1, U_0, W]]$ be absolutely convergent on the polydisc $D(0, \bar{V}_0) \times D(0, \bar{V}_1) \times D(0, \bar{U}_0) \times D(0, \bar{W})$. Let $\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)$ belonging to $G_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$. Then, the product $b(V_0, V_1, U_0, W)\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)$ belongs to $G_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$ and the following inequality*

$$(56) \quad \begin{aligned} & \|b(V_0, V_1, U_0, W)\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ & \leq |b|(\bar{V}_0, \bar{V}_1, \bar{U}_0, \bar{W})\|\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \end{aligned}$$

holds.

Proof Let

$$\begin{aligned} b(V_0, V_1, U_0, W) &= \sum_{\alpha \geq 0} b_\alpha(V_0, V_1, U_0) \frac{W^\alpha}{\alpha!}, \\ \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) &= \sum_{\alpha \geq 0} \Psi_\alpha(V_0, V_1, (U_h)_{h \in I(\alpha)}) \frac{W^\alpha}{\alpha!}. \end{aligned}$$

By definition, we get

$$(57) \quad \begin{aligned} & \|b(V_0, V_1, U_0, W)\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ &= \sum_{\alpha \geq 0} \left\| \sum_{\alpha_1 + \alpha_2 = \alpha} \alpha! \frac{b_{\alpha_1}(V_0, V_1, U_0)}{\alpha_1!} \frac{\Psi_{\alpha_2}(V_0, V_1, (U_h)_{h \in I(\alpha_2)})}{\alpha_2!} \right\|_{\rho, \alpha, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}} \bar{W}^\alpha. \end{aligned}$$

Lemma 3 *We have*

$$(58) \quad \begin{aligned} & \|b_{\alpha_1}(V_0, V_1, U_0)\Psi_{\alpha_2}(V_0, V_1, (U_h)_{h \in I(\alpha_2)})\|_{\rho, \alpha, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}} \\ & \leq \frac{\alpha_2!}{\alpha!} |b_{\alpha_1}|(\bar{V}_0, \bar{V}_1, \bar{U}_0) \|\Psi_{\alpha_2}(V_0, V_1, (U_h)_{h \in I(\alpha_2)})\|_{\rho, \alpha_2, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2)}}. \end{aligned}$$

Proof We can write

$$b_{\alpha_1}(V_0, V_1, U_0) = \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} b_{\alpha_1, n_0, n_1, l_0} \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_h} \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \prod_{h \in I(\alpha)} \frac{U_h^{l_h}}{l_h!}$$

and

$$\begin{aligned} & \Psi_{\alpha_2}(V_0, V_1, (U_h)_{h \in I(\alpha_2)}) \\ &= \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \psi_{\alpha_2, n_0, n_1, (l_h)_{h \in I(\alpha_2)}} \prod_{h \in I(\alpha) \setminus I(\alpha_2)} \delta_{0, l_h} \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \prod_{h \in I(\alpha)} \frac{U_h^{l_h}}{l_h!}. \end{aligned}$$

By remembering (45) of Proposition 5, we deduce that

$$(59) \quad \begin{aligned} & \|b_{\alpha_1}(V_0, V_1, U_0)\Psi_{\alpha_2}(V_0, V_1, (U_h)_{h \in I(\alpha_2)})\|_{\rho, \alpha, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}} \leq \\ & |b_{\alpha_1}|(\bar{V}_0, \bar{V}_1, \bar{U}_0) \left(\sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha_2)} \frac{|\psi_{\alpha_2, n_0, n_1, (l_h)_{h \in I(\alpha_2)}}|}{\exp(\sigma r_b(\alpha)\rho)} \frac{\bar{V}_0^{n_0} \bar{V}_1^{n_1} \prod_{h \in I(\alpha_2)} \bar{U}_h^{l_h}}{(n_0 + n_1 + \sum_{h \in I(\alpha_2)} l_h + \alpha)!} \right). \end{aligned}$$

Lemma 4 *We have*

$$(60) \quad \frac{1}{(n_0 + n_1 + \sum_{h \in I(\alpha_2)} l_h + \alpha)!} \leq \frac{\alpha_2!}{\alpha!} \frac{1}{(n_0 + n_1 + \sum_{h \in I(\alpha_2)} l_h + \alpha_2)!}.$$

Proof We write

$$\frac{1}{(n_0 + n_1 + \sum_{h \in I(\alpha_2)} l_h + \alpha)!} = \frac{(n_0 + n_1 + \sum_{h \in I(\alpha_2)} l_h + \alpha_2)!}{(n_0 + n_1 + \sum_{h \in I(\alpha_2)} l_h + \alpha)!} \frac{1}{(n_0 + n_1 + \sum_{h \in I(\alpha_2)} l_h + \alpha_2)!}$$

and we use the inequality

$$\frac{(a + \alpha_2)!}{(a + \alpha)!} \leq \frac{\alpha_2!}{\alpha!}$$

for all $\alpha = \alpha_1 + \alpha_2$ and all $a \in \mathbb{N}$ which follows from (48). This yields the lemma. \square

Using the fact that $\exp(\sigma r_b(\alpha)\rho) \geq \exp(\sigma r_b(\alpha_2)\rho)$ and gathering the inequalities (59) and (60) yields (58). \square

Finally, using (57) with (58), one gets

$$(61) \quad \begin{aligned} & \|b(V_0, V_1, U_0, W)\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ & \leq \sum_{\alpha \geq 0} \left(\sum_{\alpha_1 + \alpha_2 = \alpha} \frac{|b_{\alpha_1}(\bar{V}_0, \bar{V}_1, \bar{U}_0)|}{\alpha_1!} \|\Psi_{\alpha_2}(V_0, V_1, (U_h)_{h \in I(\alpha_2)})\|_{\rho, \alpha_2, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2)}} \right) \bar{W}^\alpha \end{aligned}$$

from which the inequality (56) follows. \square

Proposition 8 1) *Let $S, k \geq 0$ be integers such that*

$$(62) \quad S \geq k + 1 + \max(b(d_{1,k} + 2) + 3, d + 1 + b(d + d_{1,k} + 1)).$$

Then, there exists a constant $C_{8.1} > 0$ (which is independent of $\rho > 1$) such that

$$(63) \quad \begin{aligned} & \|B_{1,k}(V_0, V_1, U_0, W)\partial_W^{-S+k}\mathbb{D}_A\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ & \leq C_{8.1}\bar{W}^{S-k} \|\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \end{aligned}$$

for all $\Psi(V_0, V_1, (U_h)_{h \geq 0}, W) \in G_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$.

2) *Let $S, k \geq 0$ be integers such that*

$$(64) \quad S \geq k + 3 + b(2 + d_{2,k}).$$

Then, there exists a constant $C_{8.2} > 0$ (which is independent of $\rho > 1$) such that

$$(65) \quad \begin{aligned} & \|B_{2,k}(V_0, V_1, U_0, W)\partial_W^{-S+k}\mathbb{D}_B\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ & \leq C_{8.2}\bar{W}^{S-k} \|\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \end{aligned}$$

for all $\Psi(V_0, V_1, (U_h)_{h \geq 0}, W) \in G_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$.

Proof 1) We show the first inequality (63). We expand

$$B_{1,k}(V_0, V_1, U_0, W) = \sum_{\alpha \geq 0} B_{1,k,\alpha}(V_0, V_1, U_0) \frac{W^\alpha}{\alpha!}.$$

By definition, we have

$$(66) \quad \begin{aligned} & \|B_{1,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \mathbb{D}_{\mathbf{A}} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ &= \sum_{\alpha \geq 0} \left\| \sum_{\alpha_1 + \alpha_2 = \alpha, \alpha_2 \geq S-k} \alpha! \frac{B_{1,k,\alpha_1}(V_0, V_1, U_0)}{\alpha_1!} \right. \\ & \times \left(\sum_{j \in I(\alpha_2 - S + k)} \frac{\mathbf{A}_{j, \alpha_2 - S + k + 1}(V_0, V_1, (U_h)_{h \in I(\alpha_2 - S + k + 1)})}{\alpha_2!} \right. \\ & \quad \left. \left. \times (\partial_{U_j} \Psi_{\alpha_2 - S + k})(V_0, V_1, (U_h)_{h \in I(\alpha_2 - S + k)}) \right) \right\|_{\rho, \alpha, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}} \bar{W}^\alpha. \end{aligned}$$

Now, using Lemma 3, we deduce that

$$(67) \quad \begin{aligned} & \|B_{1,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \mathbb{D}_{\mathbf{A}} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ & \leq \sum_{\alpha \geq 0} \left(\sum_{\alpha_1 + \alpha_2 = \alpha, \alpha_2 \geq S-k} \frac{|B_{1,k,\alpha_1}(\bar{V}_0, \bar{V}_1, \bar{U}_0)|}{\alpha_1!} \left\| \sum_{j \in I(\alpha_2 - S + k)} \mathbf{A}_{j, \alpha_2 - S + k + 1}(V_0, V_1, (U_h)_{h \in I(\alpha_2 - S + k + 1)}) \right. \right. \\ & \quad \left. \left. \times (\partial_{U_j} \Psi_{\alpha_2 - S + k})(V_0, V_1, (U_h)_{h \in I(\alpha_2 - S + k)}) \right\|_{\rho, \alpha_2, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2)}} \right) \bar{W}^\alpha \end{aligned}$$

In the next lemma, we give estimates for the coefficients of the series $\mathbf{A}_{j,\alpha}$ and $|B_{1,k,\alpha}|$.

Lemma 5 1) *The coefficients of the Taylor series of $\mathbf{A}_{j, \alpha_2 - S + k + 1}(V_0, V_1, (U_h)_{h \in I(\alpha_2 - S + k + 1)})$*

$$\begin{aligned} & \mathbf{A}_{j, \alpha_2 - S + k + 1}(V_0, V_1, (U_h)_{h \in I(\alpha_2 - S + k + 1)}) = \\ & \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha_2 - S + k + 1)} A_{j, \alpha_2 - S + k + 1, n_0, n_1, (l_h)_{h \in I(\alpha_2 - S + k + 1)}} \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \prod_{h \in I(\alpha_2 - S + k + 1)} \frac{U_h^{l_h}}{l_h!} \end{aligned}$$

satisfy the next estimates. There exist constants $a, \delta > 0$, with $\delta > \bar{\delta}$, $a_p > 0$, $0 \leq p \leq d$ such that

$$(68) \quad \begin{aligned} & \frac{A_{j, \alpha_2 - S + k + 1, n_0, n_1, (l_h)_{h \in I(\alpha_2 - S + k + 1)}}}{n_0! n_1! \prod_{h \in I(\alpha_2 - S + k + 1)} l_h!} \\ & \leq \frac{a \nu (\alpha_2 - S + k + 1)^2 (\rho + \delta) + (d + 1) \max_{0 \leq p \leq d} a_p (\rho + \delta)^d \mathcal{P}_d(\alpha_2 - S + k)}{\delta^{n_0 + n_1 + \sum_{h \in I(\alpha_2 - S + k + 1)} l_h}} \end{aligned}$$

for all $\alpha_2 \geq S - k$, all $j \in I(\alpha_2 - S + k)$, all $n_0, n_1, l_h \geq 0$, $h \in I(\alpha_2 - S + k + 1)$ where \mathcal{P}_d is defined in (73).

2) *The coefficients of the Taylor series of $|B_{1,k,\alpha_1}(\bar{V}_0, \bar{V}_1, \bar{U}_0)|$*

$$|B_{1,k,\alpha_1}(\bar{V}_0, \bar{V}_1, \bar{U}_0)| = \sum_{n_0, n_1, l_0 \geq 0} b_{1,k,\alpha_1, n_0, n_1, l_0} \frac{\bar{V}_0^{n_0}}{n_0!} \frac{\bar{V}_1^{n_1}}{n_1!} \frac{\bar{U}_0^{l_0}}{l_0!}$$

satisfy the following inequalities. There exist constants $\delta > \bar{\delta}$, $D_{1,k}, \hat{D}_{1,k} > 0$ with

$$(69) \quad \frac{b_{1,k,\alpha_1, n_0, n_1, l_0}}{n_0! n_1! l_0!} \leq \frac{D_{1,k} (\rho + \delta)^{d_{1,k}} \alpha_1! \hat{D}_{1,k}^{\alpha_1}}{\delta^{n_0 + n_1 + l_0}}$$

for all $\alpha_1 \geq 0$, all $n_0, n_1, l_0 \geq 0$.

Proof We first treat the estimates for $\mathbf{A}_{j,\alpha}$. From the Cauchy formula in several variables, one can write

$$(70) \quad \frac{\partial_{v_0}^{n_0} \partial_{v_1}^{n_1} \prod_{h \in I(\alpha_2 - S + k + 1)} \partial_{u_h}^{l_h} A_j(v_0, v_1, (u_h)_{h \in I(\alpha_2 - S + k + 1)})}{n_0! n_1! \prod_{h \in I(\alpha_2 - S + k + 1)} l_h!}$$

$$= \left(\frac{1}{2i\pi}\right)^{\alpha_2 - S + k + 4} \int_{C(v_0, \delta)} \int_{C(v_1, \delta)} \prod_{h \in I(\alpha_2 - S + k + 1)} \int_{C(u_h, \delta)} A_j(\chi_0, \chi_1, (\xi_h)_{h \in I(\alpha_2 - S + k + 1)})$$

$$\times \frac{d\chi_0 d\chi_1 \prod_{h \in I(\alpha_2 - S + k + 1)} d\xi_h}{(\chi_0 - v_0)^{n_0 + 1} (\chi_1 - v_1)^{n_1 + 1} \prod_{h \in I(\alpha_2 - S + k + 1)} (\xi_h - u_h)^{l_h + 1}}$$

for all $|v_0| < R$, $|v_1| < R$, $|u_h| < \rho$, $h \in I(\alpha_2 - S + k + 1)$ and $j \in I(\alpha_2 - S + k)$ where R is introduced in Section 2.2. The integration is made along positively oriented circles with radius $\delta > 0$, $C(v_0, \delta)$, $C(v_1, \delta)$ and $C(u_h, \delta)$ for $h \in I(\alpha_2 - S + k + 1)$. We choose the real number $\delta > \bar{\delta}$ in such a way that $R + \delta < R'$ where R' is defined in Section 2.1 and $\bar{\delta}$ at the beginning of Section 3.1. Now, since the functions $a(\chi_0, \chi_1)$ and $a_p(\chi_0, \chi_1)$ are holomorphic on $D(0, R')^2$, the number $\nu > 0$ (see (6)) can be chosen large enough such that there exist real numbers $a, a_p > 0$, for $0 \leq p \leq d$, with

$$(71) \quad \sup_{|\chi_0| < R + \delta, |\chi_1| < R + \delta} \left| \frac{\partial_{\chi_1}^{l_1} a(\chi_0, \chi_1)}{l_1! \nu^{l_1}} \right| \leq a, \quad \sup_{|\chi_0| < R + \delta, |\chi_1| < R + \delta} \left| \frac{\partial_{\chi_1}^{l_0} a_p(\chi_0, \chi_1)}{l_0! \nu^{l_0}} \right| \leq a_p$$

for all $l_0, l_1 \geq 0$. We recall also that for any integers $k, n \geq 1$, the number of tuples $(b_1, \dots, b_k) \in \mathbb{N}^k$ such that $b_1 + \dots + b_k = n$ is $(n + k - 1)! / ((k - 1)! n!)$. From these latter statements and the definition of A_j given by (16), we deduce that

$$(72) \quad |A_j(\chi_0, \chi_1, (\xi_h)_{h \in I(\alpha_2 - S + k + 1)})| \leq a\nu(j + 1)^2(\rho + \delta) + (d + 1) \max_{0 \leq p \leq d} a_p(\rho + \delta)^d \mathcal{P}_d(j)$$

(since $\rho > 1$), where

$$(73) \quad \mathcal{P}_d(j) = \frac{(j + d)!}{j!} = \prod_{l=1}^d (j + l)$$

is a polynomial of degree d in j with positive coefficients, for all $|\chi_0| < R + \delta$, $|\chi_1| < R + \delta$, $|\xi_h| < \rho + \delta$, $h \in I(\alpha_2 - S + k + 1)$ and $j \in I(\alpha_2 - S + k)$. Gathering (70) and (72) yields (68).

Again, from the Cauchy formula in several variables, one can write

$$(74) \quad \frac{\partial_{v_0}^{n_0} \partial_{v_1}^{n_1} \partial_{u_0}^{l_0} b_{1,k,\alpha_1}(v_0, v_1, u_0)}{n_0! n_1! l_0!} = \left(\frac{1}{2i\pi}\right)^3 \int_{C(v_0, \delta)} \int_{C(v_1, \delta)} \int_{C(u_0, \delta)} b_{1,k,\alpha_1}(\chi_0, \chi_1, \xi_0)$$

$$\times \frac{d\chi_0 d\chi_1 d\xi_0}{(\chi_0 - v_0)^{n_0 + 1} (\chi_1 - v_1)^{n_1 + 1} (\xi_0 - u_0)^{l_0 + 1}}$$

for all $|v_0| < R$, $|v_1| < R$, $|u_0| < \rho$. Again, one chooses the real number $\delta > \bar{\delta}$ in such a way that $R + \delta < R'$. By construction of b_{1,k,α_1} in Section 2.2, we know that there exist two constants $D_{1,k}, \hat{D}_{1,k} > 0$ such that

$$(75) \quad |b_{1,k,\alpha_1}(\chi_0, \chi_1, \xi_0)| \leq D_{1,k}(\rho + \delta)^{d_{1,k}} \alpha_1! \hat{D}_{1,k}^{\alpha_1}$$

for all $\alpha_1 \geq 0$, all $|\chi_0| < R + \delta$, $|\chi_1| < R + \delta$, $|\xi_0| < \rho + \delta$. Gathering (74) and (75) yields (69). \square

From (69), we deduce that

$$(76) \quad \frac{|B_{1,k,\alpha_1}|(\bar{V}_0, \bar{V}_1, \bar{U}_0)}{\alpha_1!} \leq \frac{D_{1,k}(\rho + \delta)^{d_{1,k}} \hat{D}_{1,k}^{\alpha_1}}{(1 - \frac{\bar{V}_0}{\delta})(1 - \frac{\bar{V}_1}{\delta})(1 - \frac{\bar{U}_0}{\delta})}.$$

On the other hand, from the Proposition 5, we deduce that

$$(77) \quad \begin{aligned} & \|\mathbf{A}_{j,\alpha_2-S+k+1}(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k+1)}) \\ & \quad \times (\partial_{U_j} \Psi_{\alpha_2-S+k})(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k)})\|_{\rho, \alpha_2, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2)}} \\ & \leq |\mathbf{A}_{j,\alpha_2-S+k+1}|(\bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2-S+k+1)}) \\ & \quad \times \|(\partial_{U_j} \Psi_{\alpha_2-S+k})(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k)})\|_{\rho, \alpha_2, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2)}}. \end{aligned}$$

From (68), we deduce that

$$(78) \quad \begin{aligned} & |\mathbf{A}_{j,\alpha_2-S+k+1}|(\bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2-S+k+1)}) \\ & \leq \frac{a\nu(\alpha_2 - S + k + 1)^2(\rho + \delta) + (d + 1) \max_{0 \leq p \leq d} a_p(\rho + \delta)^d \mathcal{P}_d(\alpha_2 - S + k)}{(1 - \frac{\bar{V}_0}{\delta})(1 - \frac{\bar{V}_1}{\delta}) \prod_{h \in I(\alpha_2-S+k+1)} (1 - \frac{\bar{U}_h}{\delta})} \end{aligned}$$

for all $j \in I(\alpha_2 - S + k)$. Now, from the definition of $\bar{U}_h = \bar{\delta}/(h^b + 1)$, where $b > 1$, we know that there exists $\kappa > 0$ such that

$$(79) \quad \prod_{h \in I(\alpha)} (1 - \frac{\bar{U}_h}{\delta}) \geq \kappa$$

for all $\alpha \geq 0$. From Proposition 6, we have that

$$(80) \quad \begin{aligned} & \|(\partial_{U_j} \Psi_{\alpha_2-S+k})(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k)})\|_{\rho, \alpha_2, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2)}} \leq \frac{\exp(-\sigma \rho \frac{S-k}{(\alpha_2+1)^b})}{\bar{U}_j \prod_{l=1}^{S-k-1} (\alpha_2 - l + 1)} \\ & \quad \times \|\Psi_{\alpha_2-S+k}(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k)})\|_{\rho, \alpha_2-S+k, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2-S+k)}}. \end{aligned}$$

Collecting the estimates (78), (79) and (80), we get from (77) that

$$(81) \quad \begin{aligned} & \left\| \sum_{j \in I(\alpha_2-S+k)} \mathbf{A}_{j,\alpha_2-S+k+1}(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k+1)}) \right. \\ & \quad \times (\partial_{U_j} \Psi_{\alpha_2-S+k})(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k)})\|_{\rho, \alpha_2, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2)}} \\ & \leq \mathcal{A}_{\rho, \alpha_2} \|\Psi_{\alpha_2-S+k}(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k)})\|_{\rho, \alpha_2-S+k, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2-S+k)}} \end{aligned}$$

where

$$\begin{aligned} \mathcal{A}_{\rho, \alpha_2} = & \frac{a\nu(\alpha_2 - S + k + 1)^2(\rho + \delta) + (d + 1) \max_{0 \leq p \leq d} a_p(\rho + \delta)^d \mathcal{P}_d(\alpha_2 - S + k)}{(1 - \frac{\bar{V}_0}{\delta})(1 - \frac{\bar{V}_1}{\delta}) \kappa} \\ & \times \frac{\exp(-\sigma \rho \frac{S-k}{(\alpha_2+1)^b})}{\prod_{l=1}^{S-k-1} (\alpha_2 - l + 1)} \frac{1}{\bar{\delta}} (\alpha_2 - S + k + 1) ((\alpha_2 - S + k)^b + 1). \end{aligned}$$

Now, we recall the following classical estimates. Let $\delta, m_1, m_2 > 0$ positive real numbers, then

$$(82) \quad \sup_{x \geq 0} (x + \delta)^{m_1} \exp(-m_2 x) \leq \left(\frac{m_1}{m_2}\right)^{m_1} \exp(-m_1) \exp(\delta m_2)$$

holds. Hence,

$$(83) \quad (\rho + \delta)^{d_{1,k}} \mathcal{A}_{\rho, \alpha_2} \leq \left(\frac{a\nu(\alpha_2 - S + k + 1)^2(\alpha_2 + 1)^{b(1+d_{1,k})} \left(\frac{\exp(-1)(1+d_{1,k})}{\sigma(S-k)} \right)^{1+d_{1,k}} \exp(\delta\sigma(S-k))}{(1 - \frac{\bar{V}_0}{\delta})(1 - \frac{\bar{V}_1}{\delta})\kappa} \right. \\ \left. + \frac{(d+1) \max_{0 \leq p \leq d} a_p \mathcal{P}_d(\alpha_2 - S + k)(\alpha_2 + 1)^{b(d+d_{1,k})} \left(\frac{(d+d_{1,k}) \exp(-1)}{\sigma(S-k)} \right)^{d+d_{1,k}} \exp(\delta\sigma(S-k))}{(1 - \frac{\bar{V}_0}{\delta})(1 - \frac{\bar{V}_1}{\delta})\kappa} \right) \\ \times \frac{(\alpha_2 - S + k + 1)((\alpha_2 - S + k)^b + 1)}{\bar{\delta} \prod_{l=1}^{S-k-1} (\alpha_2 - l + 1)}.$$

Under the assumptions (62), one gets a constant $\tilde{C}_{8.1} > 0$ (depending on $a, \max_{0 \leq p \leq d} a_p, \delta, \bar{\delta}, b, d, d_{1,k}, \sigma, \nu, S, k, \kappa, \bar{V}_0, \bar{V}_1$) such that

$$(84) \quad (\rho + \delta)^{d_{1,k}} \mathcal{A}_{\rho, \alpha_2} \leq \tilde{C}_{8.1}$$

for all $\rho \geq 0$, all $\alpha_2 \geq S - k$. Finally, gathering (67), (76), (81), (84), one gets that

$$(85) \quad \|B_{1,k}(V_0, V_1, U_0, W) \partial_{\bar{W}}^{-S+k} \mathbb{D}_{\mathbf{A}} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ \leq \sum_{\alpha \geq 0} \left(\sum_{\alpha_1 + \alpha_2 = \alpha, \alpha_2 \geq S-k} \frac{D_{1,k}}{(1 - \frac{\bar{V}_0}{\delta})(1 - \frac{\bar{V}_1}{\delta})(1 - \frac{\bar{U}_0}{\delta})} \hat{D}_{1,k}^{\alpha_1} \right) \\ \times \tilde{C}_{8.1} \|\Psi_{\alpha_2 - S + k}(V_0, V_1, (U_h)_{h \in I(\alpha_2 - S + k)})\|_{\rho, \alpha_2 - S + k, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2 - S + k)}} \bar{W}^{\alpha_1 + \alpha_2 - S + k} \bar{W}^{S-k} \\ = \frac{\tilde{C}_{8.1} D_{1,k}}{(1 - \frac{\bar{V}_0}{\delta})(1 - \frac{\bar{V}_1}{\delta})(1 - \frac{\bar{U}_0}{\delta})(1 - \hat{D}_{1,k} \bar{W})} \\ \times \bar{W}^{S-k} \|\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$$

provided that $\bar{V}_0 < \delta, \bar{V}_1 < \delta, \bar{U}_0 < \delta$ and $\bar{W} < 1/\hat{D}_{1,k}$, which yields (63).

2) Now, we turn towards the estimates (65) which will follow from the same arguments as in 1). Using Lemma 3, we get that

$$(86) \quad \|B_{2,k}(V_0, V_1, U_0, W) \partial_{\bar{W}}^{-S+k} \mathbb{D}_{\mathbf{B}} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ \leq \sum_{\alpha \geq 0} \left(\sum_{\alpha_1 + \alpha_2 = \alpha, \alpha_2 \geq S-k} \frac{|B_{2,k, \alpha_1}(\bar{V}_0, \bar{V}_1, \bar{U}_0)|}{\alpha_1!} \right) \sum_{j \in I(\alpha_2 - S + k)} \mathbf{B}_{j, \alpha_2 - S + k + 1}(V_0, V_1, (U_h)_{h \in I(\alpha_2 - S + k + 1)}) \\ \times (\partial_{U_j} \Psi_{\alpha_2 - S + k})(V_0, V_1, (U_h)_{h \in I(\alpha_2 - S + k)})\|_{\rho, \alpha_2, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2)}} \bar{W}^\alpha$$

In the next lemma, we give estimates for the coefficients of the series $\mathbf{B}_{j, \alpha}$ and $|B_{2,k, \alpha}|$.

Lemma 6 1) *The coefficients of the Taylor series of $\mathbf{B}_{j, \alpha_2 - S + k + 1}(V_0, V_1, (U_h)_{h \in I(\alpha_2 - S + k + 1)})$*

$$\mathbf{B}_{j, \alpha_2 - S + k + 1}(V_0, V_1, (U_h)_{h \in I(\alpha_2 - S + k + 1)}) = \\ \sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha_2 - S + k + 1)} B_{j, \alpha_2 - S + k + 1, n_0, n_1, (l_h)_{h \in I(\alpha_2 - S + k + 1)}} \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \prod_{h \in I(\alpha_2 - S + k + 1)} \frac{U_h^{l_h}}{l_h!}$$

satisfy the next estimates. There exist a constant $\delta > 0$, with $\delta > \bar{\delta}$ such that

$$(87) \quad \frac{B_{j,\alpha_2-S+k+1,n_0,n_1,(l_h)_{h \in I(\alpha_2-S+k+1)}}}{n_0!n_1!\prod_{h \in I(\alpha_2-S+k+1)}l_h!} \leq \frac{\nu(\alpha_2-S+k+1)(\rho+\delta)}{\delta^{n_0+n_1+\sum_{h \in I(\alpha_2-S+k+1)}l_h}}$$

for all $\alpha_2 \geq S-k$, all $j \in I(\alpha_2-S+k)$, all $n_0, n_1, l_h \geq 0$, $h \in I(\alpha_2-S+k+1)$.

2) The coefficients of the Taylor series of $|B_{2,k,\alpha_1}|(\bar{V}_0, \bar{V}_1, \bar{U}_0)$

$$|B_{2,k,\alpha_1}|(\bar{V}_0, \bar{V}_1, \bar{U}_0) = \sum_{n_0, n_1, l_0 \geq 0} b_{2,k,\alpha_1,n_0,n_1,l_0} \frac{\bar{V}_0^{n_0}}{n_0!} \frac{\bar{V}_1^{n_1}}{n_1!} \frac{\bar{U}_0^{l_0}}{l_0!}$$

satisfy the following inequalities. There exist constants $\delta > \bar{\delta}$, $D_{2,k}, \hat{D}_{2,k} > 0$ with

$$(88) \quad \frac{b_{2,k,\alpha_1,n_0,n_1,l_0}}{n_0!n_1!l_0!} \leq \frac{D_{2,k}(\rho+\delta)^{d_{2,k}}\alpha_1!\hat{D}_{2,k}^{\alpha_1}}{\delta^{n_0+n_1+l_0}}$$

for all $\alpha_1 \geq 0$, all $n_0, n_1, l_0 \geq 0$.

Proof 1) From the Cauchy formula in several variables, one can check that

$$(89) \quad \frac{\partial_{v_0}^{n_0} \partial_{v_1}^{n_1} \prod_{h \in I(\alpha_2-S+k+1)} \partial_{u_h}^{l_h} B_j(v_0, v_1, (u_h)_{h \in I(\alpha_2-S+k+1)})}{n_0!n_1!\prod_{h \in I(\alpha_2-S+k+1)}l_h!} \\ = \left(\frac{1}{2i\pi}\right)^{\alpha_2-S+k+4} \int_{C(v_0,\delta)} \int_{C(v_1,\delta)} \prod_{h \in I(\alpha_2-S+k+1)} \int_{C(u_h,\delta)} B_j(\chi_0, \chi_1, (\xi_h)_{h \in I(\alpha_2-S+k+1)}) \\ \times \frac{d\chi_0 d\chi_1 \prod_{h \in I(\alpha_2-S+k+1)} d\xi_h}{(\chi_0 - v_0)^{n_0+1} (\chi_1 - v_1)^{n_1+1} \prod_{h \in I(\alpha_2-S+k+1)} (\xi_h - u_h)^{l_h+1}}$$

for all $|v_0| < R$, $|v_1| < R$, $|u_h| < \rho$, $h \in I(\alpha_2-S+k+1)$ and $j \in I(\alpha_2-S+k)$. We choose the real number $\delta > \bar{\delta}$ in such a way that $R+\delta < R'$. From the definition given in (17), we get that

$$(90) \quad |B_j(\chi_0, \chi_1, (\xi_h)_{h \in I(\alpha_2-S+k+1)})| \leq \nu(j+1)(\rho+\delta)$$

for all $|\chi_0| < R+\delta$, $|\chi_1| < R+\delta$, $|\xi_h| < \rho+\delta$, $h \in I(\alpha_2-S+k+1)$ and $j \in I(\alpha_2-S+k)$. Gathering (89) and (90) yields (87).

2) The proof is exactly the same as 2) in Lemma 5. \square

From (88), we deduce that

$$(91) \quad \frac{|B_{2,k,\alpha_1}|(\bar{V}_0, \bar{V}_1, \bar{U}_0)}{\alpha_1!} \leq \frac{D_{2,k}(\rho+\delta)^{d_{2,k}}\hat{D}_{2,k}^{\alpha_1}}{(1-\frac{\bar{V}_0}{\delta})(1-\frac{\bar{V}_1}{\delta})(1-\frac{\bar{U}_0}{\delta})}$$

Using Propositions 5,6, we deduce that

$$(92) \quad \left\| \sum_{j \in I(\alpha_2-S+k)} \mathbf{B}_{j,\alpha_2-S+k+1}(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k+1)}) \right. \\ \left. \times (\partial_{U_j} \Psi_{\alpha_2-S+k})(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k)}) \right\|_{\rho,\alpha_2,\bar{V}_0,\bar{V}_1,(\bar{U}_h)_{h \in I(\alpha_2)}} \\ \leq \mathcal{B}_{\rho,\alpha_2} \left\| \Psi_{\alpha_2-S+k}(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k)}) \right\|_{\rho,\alpha_2-S+k,\bar{V}_0,\bar{V}_1,(\bar{U}_h)_{h \in I(\alpha_2-S+k)}}$$

where

$$\mathcal{B}_{\rho, \alpha_2} = \frac{\nu(\alpha_2 - S + k + 1)(\rho + \delta)}{(1 - \frac{\bar{V}_0}{\delta})(1 - \frac{\bar{V}_1}{\delta})\kappa} \times \frac{\exp(-\sigma\rho\frac{S-k}{(\alpha_2+1)^b})}{\prod_{l=1}^{S-k-1}(\alpha_2 - l + 1)} \frac{1}{\bar{\delta}} (\alpha_2 - S + k + 1)((\alpha_2 - S + k)^b + 1)$$

and where κ is introduced in (79). Using the estimates (82), we get

$$(93) \quad (\rho + \delta)^{d_{2,k}} \mathcal{B}_{\rho, \alpha_2} \leq \left(\frac{\nu(\alpha_2 - S + k + 1)(\alpha_2 + 1)^{b(1+d_{2,k})} \left(\frac{\exp(-1)(1+d_{2,k})}{\sigma(S-k)}\right)^{1+d_{2,k}} \exp(\delta\sigma(S-k))}{(1 - \frac{\bar{V}_0}{\delta})(1 - \frac{\bar{V}_1}{\delta})\kappa} \right) \times \frac{(\alpha_2 - S + k + 1)((\alpha_2 - S + k)^b + 1)}{\bar{\delta} \prod_{l=1}^{S-k-1}(\alpha_2 - l + 1)}.$$

Under the assumptions (64), one gets a constant $\tilde{C}_{8.2} > 0$ (depending on $\delta, \bar{\delta}, b, d_{2,k}, \sigma, \nu, S, k, \kappa, \bar{V}_0, \bar{V}_1$) such that

$$(94) \quad (\rho + \delta)^{d_{2,k}} \mathcal{B}_{\rho, \alpha_2} \leq \tilde{C}_{8.2}$$

for all $\rho \geq 0$, all $\alpha_2 \geq S - k$. Finally, gathering (86), (91), (92) and (94), we get (65). \square

Proposition 9 1) Let $S, k \geq 0$ be integers such that

$$(95) \quad S \geq k + 1 + b \max(d_{1,k}, d_{2,k}).$$

Then, for $m \in \{0, 1\}$, there exists a constant $C_9 > 0$ (which is independent of $\rho > 1$) such that

$$(96) \quad \|B_{m+1,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \partial_{V_m} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ \leq C_9 \bar{W}^{S-k} \|\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$$

for all $\Psi(V_0, V_1, (U_h)_{h \geq 0}, W) \in G_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$.

2) Let $S, k \geq 0$ be integers such that

$$(97) \quad S \geq k + b d_{3,k}.$$

Then, there exists a constant $C_{9.1} > 0$ (which is independent of $\rho > 1$) such that

$$(98) \quad \|B_{3,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ \leq C_{9.1} \bar{W}^{S-k} \|\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$$

for all $\Psi(V_0, V_1, (U_h)_{h \geq 0}, W) \in G_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$.

Proof 1) We expand

$$B_{m+1,k}(V_0, V_1, U_0, W) = \sum_{\alpha \geq 0} B_{m+1,k,\alpha}(V_0, V_1, U_0) \frac{W^\alpha}{\alpha!}.$$

By definition, we have

$$(99) \quad \begin{aligned} & \|B_{m+1,k}(V_0, V_1, U_0, W) \partial_{\bar{W}}^{-S+k} \partial_{V_m} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) \|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ &= \sum_{\alpha \geq 0} \left\| \sum_{\alpha_1 + \alpha_2 = \alpha, \alpha_2 \geq S-k} \alpha! \frac{B_{m+1,k,\alpha_1}(V_0, V_1, U_0)}{\alpha_1!} \right. \\ & \quad \times \left. \frac{\partial_{V_m} \Psi_{\alpha_2-S+k}(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k)})}{\alpha_2!} \right\|_{\rho, \alpha, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}} \bar{W}^\alpha. \end{aligned}$$

Now, using Lemma 3, we deduce that

$$(100) \quad \begin{aligned} & \|B_{m+1,k}(V_0, V_1, U_0, W) \partial_{\bar{W}}^{-S+k} \partial_{V_m} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) \|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ & \leq \sum_{\alpha \geq 0} \left(\sum_{\alpha_1 + \alpha_2 = \alpha, \alpha_2 \geq S-k} \frac{|B_{m+1,k,\alpha_1}(\bar{V}_0, \bar{V}_1, \bar{U}_0)|}{\alpha_1!} \right. \\ & \quad \times \left. \|(\partial_{V_m} \Psi_{\alpha_2-S+k})(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k)})\|_{\rho, \alpha_2, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2)}} \right) \bar{W}^\alpha \end{aligned}$$

From Proposition 6, we know that

$$(101) \quad \begin{aligned} & \|(\partial_{V_m} \Psi_{\alpha_2-S+k})(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k)})\|_{\rho, \alpha_2, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2)}} \leq \frac{\exp(-\sigma \rho \frac{S-k}{(\alpha_2+1)^b})}{\bar{V}_m \prod_{l=1}^{S-k-1} (\alpha_2 - l + 1)} \\ & \quad \times \|\Psi_{\alpha_2-S+k}(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k)})\|_{\rho, \alpha_2-S+k, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2-S+k)}}. \end{aligned}$$

From (76), (91), (100) and (101), we get that

$$(102) \quad \begin{aligned} & \|B_{m+1,k}(V_0, V_1, U_0, W) \partial_{\bar{W}}^{-S+k} \partial_{V_m} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) \|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ & \leq \sum_{\alpha \geq 0} \left(\sum_{\alpha_1 + \alpha_2 = \alpha, \alpha_2 \geq S-k} \hat{D}_{m+1,k}^{\alpha_1} \right. \\ & \quad \times \left. \mathcal{C}_{\rho, \alpha_2} \|\Psi_{\alpha_2-S+k}(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k)})\|_{\rho, \alpha_2-S+k, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2-S+k)}} \right) \bar{W}^\alpha \end{aligned}$$

where

$$\mathcal{C}_{\rho, \alpha_2} = \frac{D_{m+1,k}(\rho + \delta)^{d_{m+1,k}}}{(1 - \frac{\bar{V}_0}{\delta})(1 - \frac{\bar{V}_1}{\delta})(1 - \frac{\bar{U}_0}{\delta})} \times \frac{\exp(-\sigma \rho \frac{S-k}{(\alpha_2+1)^b})}{\bar{V}_m \prod_{l=1}^{S-k-1} (\alpha_2 - l + 1)}.$$

Using the estimates (82), we deduce that

$$\mathcal{C}_{\rho, \alpha_2} \leq \frac{D_{m+1,k} \left(\frac{d_{m+1,k} \exp(-1)}{\sigma(S-k)} \right)^{d_{m+1,k}} \exp(\delta \sigma(S-k))}{(1 - \frac{\bar{V}_0}{\delta})(1 - \frac{\bar{V}_1}{\delta})(1 - \frac{\bar{U}_0}{\delta})} \times \frac{(\alpha_2 + 1)^{bd_{m+1,k}}}{\bar{V}_m \prod_{l=1}^{S-k-1} (\alpha_2 - l + 1)}.$$

Under the assumption (95), we get a constant $\tilde{C}_9 > 0$ (depending on $D_{m+1,k}, d_{m+1,k}, S, k, \delta, \sigma, \bar{V}_0, \bar{V}_1, \bar{U}_0, b$) such that

$$(103) \quad \mathcal{C}_{\rho, \alpha_2} \leq \tilde{C}_9$$

for all $\rho > 1$, all $\alpha_2 \geq S - k$. Finally, collecting (102) and (103), we get

$$\begin{aligned}
(104) \quad & \|B_{m+1,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \partial_{V_m} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\
& \leq \sum_{\alpha \geq 0} \left(\sum_{\alpha_1 + \alpha_2 = \alpha, \alpha_2 \geq S-k} \hat{D}_{m+1,k}^{\alpha_1} \right. \\
& \quad \times \tilde{C}_9 \|\Psi_{\alpha_2-S+k}(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k)})\|_{\rho, \alpha_2-S+k, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2-S+k)}} \bar{W}^{\alpha_1 + \alpha_2 - S+k} \bar{W}^{S-k} \\
& \quad \left. = \frac{\tilde{C}_9}{1 - \hat{D}_{m+1,k} \bar{W}} \bar{W}^{S-k} \|\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \right)
\end{aligned}$$

which yields (96).

2) We expand

$$B_{3,k}(V_0, V_1, U_0, W) = \sum_{\alpha \geq 0} B_{3,k,\alpha}(V_0, V_1, U_0) \frac{W^\alpha}{\alpha!}.$$

By definition, we have

$$\begin{aligned}
(105) \quad & \|B_{3,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\
& = \sum_{\alpha \geq 0} \left\| \sum_{\alpha_1 + \alpha_2 = \alpha, \alpha_2 \geq S-k} \alpha! \frac{B_{3,k,\alpha_1}(V_0, V_1, U_0)}{\alpha_1!} \right. \\
& \quad \times \left. \frac{\Psi_{\alpha_2-S+k}(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k)})}{\alpha_2!} \right\|_{\rho, \alpha, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha)}} \bar{W}^\alpha.
\end{aligned}$$

Now, using Lemma 3, we deduce that

$$\begin{aligned}
(106) \quad & \|B_{3,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\
& = \sum_{\alpha \geq 0} \left(\sum_{\alpha_1 + \alpha_2 = \alpha, \alpha_2 \geq S-k} \frac{|B_{3,k,\alpha_1}(\bar{V}_0, \bar{V}_1, \bar{U}_0)|}{\alpha_1!} \right. \\
& \quad \times \left. \|\Psi_{\alpha_2-S+k}(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k)})\|_{\rho, \alpha_2, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2)}} \bar{W}^\alpha \right).
\end{aligned}$$

From Proposition 6, we know that

$$\begin{aligned}
(107) \quad & \|\Psi_{\alpha_2-S+k}(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k)})\|_{\rho, \alpha_2, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2)}} \leq \frac{\exp(-\sigma \rho \frac{S-k}{(\alpha_2+1)^b})}{\prod_{l=1}^{S-k} (\alpha_2 - l + 1)} \\
& \quad \times \|\Psi_{\alpha_2-S+k}(V_0, V_1, (U_h)_{h \in I(\alpha_2-S+k)})\|_{\rho, \alpha_2-S+k, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2-S+k)}}.
\end{aligned}$$

On the other hand, the coefficients of the Taylor series of $|B_{3,k,\alpha_1}(\bar{V}_0, \bar{V}_1, \bar{U}_0)|$

$$|B_{3,k,\alpha_1}(\bar{V}_0, \bar{V}_1, \bar{U}_0)| = \sum_{n_0, n_1, l_0 \geq 0} b_{3,k,\alpha_1, n_0, n_1, l_0} \frac{\bar{V}_0^{n_0} \bar{V}_1^{n_1} \bar{U}_0^{l_0}}{n_0! n_1! l_0!}$$

satisfy the following inequalities. There exist constants $\delta > \bar{\delta}$, $D_{3,k}, \hat{D}_{3,k} > 0$ with

$$(108) \quad \frac{b_{3,k,\alpha_1, n_0, n_1, l_0}}{n_0! n_1! l_0!} \leq \frac{D_{3,k} (\rho + \delta)^{d_{3,k}} \alpha_1! \hat{D}_{3,k}^{\alpha_1}}{\delta^{n_0 + n_1 + l_0}}$$

for all $\alpha_1 \geq 0$, all $n_0, n_1, l_0 \geq 0$. The proof copies 2) from Lemma 5. From (108), we deduce that

$$(109) \quad \frac{|B_{3,k,\alpha_1}|(\bar{V}_0, \bar{V}_1, \bar{U}_0)}{\alpha_1!} \leq \frac{D_{3,k}(\rho + \delta)^{d_{3,k}} \hat{D}_{3,k}^{\alpha_1}}{(1 - \frac{\bar{V}_0}{\delta})(1 - \frac{\bar{V}_1}{\delta})(1 - \frac{\bar{U}_0}{\delta})}.$$

From (109), (106) and (107), we get that

$$(110) \quad \|B_{3,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ \leq \sum_{\alpha \geq 0} \left(\sum_{\alpha_1 + \alpha_2 = \alpha, \alpha_2 \geq S-k} \hat{D}_{3,k}^{\alpha_1} \right) \\ \times \mathcal{D}_{\rho, \alpha_2} \|\Psi_{\alpha_2 - S+k}(V_0, V_1, (U_h)_{h \in I(\alpha_2 - S+k)})\|_{\rho, \alpha_2 - S+k, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2 - S+k)}} \bar{W}^\alpha$$

where

$$\mathcal{D}_{\rho, \alpha_2} = \frac{D_{3,k}(\rho + \delta)^{d_{3,k}}}{(1 - \frac{\bar{V}_0}{\delta})(1 - \frac{\bar{V}_1}{\delta})(1 - \frac{\bar{U}_0}{\delta})} \times \frac{\exp(-\sigma \rho \frac{S-k}{(\alpha_2+1)^b})}{\prod_{l=1}^{S-k} (\alpha_2 - l + 1)}.$$

Using the estimates (82), we deduce that

$$\mathcal{D}_{\rho, \alpha_2} \leq \frac{D_{3,k}(\frac{d_{3,k} \exp(-1)}{\sigma(S-k)})^{d_{3,k}} \exp(\delta \sigma(S-k))}{(1 - \frac{\bar{V}_0}{\delta})(1 - \frac{\bar{V}_1}{\delta})(1 - \frac{\bar{U}_0}{\delta})} \times \frac{(\alpha_2 + 1)^{bd_{3,k}}}{\prod_{l=1}^{S-k} (\alpha_2 - l + 1)}.$$

Under the assumption (97), we get a constant $\tilde{C}_{9.1} > 0$ (depending on $D_{3,k}, d_{3,k}, S, k, \delta, \sigma, \bar{V}_0, \bar{V}_1, \bar{U}_0, b$) such that

$$(111) \quad \mathcal{D}_{\rho, \alpha_2} \leq \tilde{C}_{9.1}$$

for all $\rho > 1$, all $\alpha_2 \geq S - k$. Finally, collecting (110) and (111), we get

$$(112) \quad \|B_{3,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ \leq \sum_{\alpha \geq 0} \left(\sum_{\alpha_1 + \alpha_2 = \alpha, \alpha_2 \geq S-k} \hat{D}_{3,k}^{\alpha_1} \right) \\ \times \tilde{C}_{9.1} \|\Psi_{\alpha_2 - S+k}(V_0, V_1, (U_h)_{h \in I(\alpha_2 - S+k)})\|_{\rho, \alpha_2 - S+k, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \in I(\alpha_2 - S+k)}} \bar{W}^{\alpha_1 + \alpha_2 - S+k} \bar{W}^{S-k} \\ = \frac{\tilde{C}_{9.1}}{1 - \hat{D}_{3,k} \bar{W}} \bar{W}^{S-k} \|\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$$

which yields (98).

□

3.2 A functional partial differential equation in the Banach spaces of infinitely many variables $G_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$.

In the next proposition, we solve a functional fixed point equation within the Banach spaces of formal series introduced in the previous subsection.

Proposition 10 *We make the following assumptions*

$$(113) \quad S \geq k + 1 + \max(b(d_{1,k} + 2) + 3, d + 1 + b(d + d_{1,k} + 1)) \quad , \quad S \geq k + 3 + b(2 + d_{2,k}), \\ S \geq k + 1 + b \max(d_{1,k}, d_{2,k}) \quad , \quad S \geq k + bd_{3,k}$$

for all $k \in \mathcal{S}$. Then, for given $\bar{V}_0, \bar{V}_1, \bar{\delta} > 0$, there exists $\bar{W} > 0$ (independent of $\rho > 1$) such that, for all $\tilde{I}(V_0, V_1, (U_h)_{h \geq 0}, W) \in G_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$, the functional equation

$$(114) \quad \begin{aligned} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) &= \sum_{k \in \mathcal{S}} B_{1,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \partial_{V_0} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) \\ &\quad + B_{1,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \mathbb{D}_{\mathbf{A}} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) \\ &\quad + \sum_{k \in \mathcal{S}} B_{2,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \partial_{V_1} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) \\ &\quad + B_{2,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \mathbb{D}_{\mathbf{B}} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) \\ &\quad + \sum_{k \in \mathcal{S}} B_{3,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) \\ &\quad + \tilde{I}(V_0, V_1, (U_h)_{h \geq 0}, W) \end{aligned}$$

has a unique solution $\Psi(V_0, V_1, (U_h)_{h \geq 0}, W) \in G_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$. Moreover, we have that

$$(115) \quad \|\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \leq 2 \|\tilde{I}(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}.$$

Proof We consider the map \mathfrak{M} from the space $\mathbb{G}[[V_0, V_1, (U_h)_{h \geq 0}, W]]$ of formal series (introduced in Definition 1) into itself defined as follows:

$$(116) \quad \begin{aligned} \mathfrak{M}(\Delta(V_0, V_1, (U_h)_{h \geq 0}, W)) &= \sum_{k \in \mathcal{S}} B_{1,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \partial_{V_0} \Delta(V_0, V_1, (U_h)_{h \geq 0}, W) \\ &\quad + B_{1,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \mathbb{D}_{\mathbf{A}} \Delta(V_0, V_1, (U_h)_{h \geq 0}, W) \\ &\quad + \sum_{k \in \mathcal{S}} B_{2,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \partial_{V_1} \Delta(V_0, V_1, (U_h)_{h \geq 0}, W) \\ &\quad + B_{2,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \mathbb{D}_{\mathbf{B}} \Delta(V_0, V_1, (U_h)_{h \geq 0}, W) \\ &\quad + \sum_{k \in \mathcal{S}} B_{3,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \Delta(V_0, V_1, (U_h)_{h \geq 0}, W) \end{aligned}$$

for all $\Delta(V_0, V_1, (U_h)_{h \geq 0}, W) \in \mathbb{G}[[V_0, V_1, (U_h)_{h \geq 0}, W]]$.

In order to prove the proposition, we need the following lemma.

Lemma 7 *Let id the identity map $x \mapsto x$ from $\mathbb{G}[[V_0, V_1, (U_h)_{h \geq 0}, W]]$ into itself. Then, for a well chosen $\bar{W} > 0$, the map $\text{id} - \mathfrak{M}$ defines an invertible map such that $(\text{id} - \mathfrak{M})^{-1}$ is defined from $G_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$ into itself. Moreover, we have that*

$$(117) \quad \begin{aligned} \|(\text{id} - \mathfrak{M})^{-1}(\Xi(V_0, V_1, (U_h)_{h \geq 0}, W))\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ \leq 2 \|\Xi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \end{aligned}$$

for all $\Xi(V_0, V_1, (U_h)_{h \geq 0}, W) \in G_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$.

Proof Taking care of the constraints (113), we get from Propositions 8 and 9 a constant $C_{10} > 0$ (depending on the constants introduced above and also on the aforementioned propositions: $a, \max_{0 \leq p \leq d} a_p, \delta, \bar{\delta}, b, d, \max_{k \in \mathcal{S}} d_{1,k}$,

$\max_{k \in \mathcal{S}} D_{1,k}$, $\max_{k \in \mathcal{S}} d_{2,k}$, $\max_{k \in \mathcal{S}} D_{2,k}$, $\max_{k \in \mathcal{S}} d_{3,k}$, $\max_{k \in \mathcal{S}} D_{3,k}$, $\sigma, \nu, S, \mathcal{S}, \kappa, \bar{V}_0, \bar{V}_1$ but independent of $\rho > 1$) such that

$$\begin{aligned} & \|\mathfrak{M}(\Delta(V_0, V_1, (U_h)_{h \geq 0}, W))\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ & \leq C_{10} \left(\sum_{k \in \mathcal{S}} \bar{W}^{S-k} \right) \|\Delta(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \end{aligned}$$

for all $\Delta(V_0, V_1, (U_h)_{h \geq 0}, W) \in G_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$ with $0 \leq \bar{W} \leq \min_{m \in \{0,1,2\}, k \in \mathcal{S}} 1/(2\hat{D}_{m+1,k})$. Since $S > k$ for all $k \in \mathcal{S}$, we can choose $\bar{W} > 0$ such that

$$C_{10} \sum_{k \in \mathcal{S}} \bar{W}^{S-k} < \frac{1}{2}$$

together with $\bar{W} \leq \min_{m \in \{0,1,2\}, k \in \mathcal{S}} 1/(2\hat{D}_{m+1,k})$. We deduce that

$$\begin{aligned} & \|\mathfrak{M}(\Delta(V_0, V_1, (U_h)_{h \geq 0}, W))\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \\ & \leq \frac{1}{2} \|\Delta(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \end{aligned}$$

for all $\Delta(V_0, V_1, (U_h)_{h \geq 0}, W) \in G_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$. This yields the estimates (117). \square

Finally, let $\tilde{I}(V_0, V_1, (U_h)_{h \geq 0}, W) \in G_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$ for $\bar{W} > 0$ chosen as in Lemma 7. We define

$$\Psi(V_0, V_1, (U_h)_{h \geq 0}, W) = (\text{id} - \mathfrak{M})^{-1}(\tilde{I}(V_0, V_1, (U_h)_{h \geq 0}, W)).$$

By construction, $\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)$ belongs to $G_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$ and solves the equation (114) with the estimates (115). \square

4 Analytic solutions with growth estimates of linear partial differential equations in \mathbb{C}^3 .

We are now in position to state the main result of our work.

Theorem 1 *Let $b_{m,k}(t, z, u_0, w)$ be the functions defined in (11) for $m = 1, 2, 3$ and $k \in \mathcal{S}$. Let us assume that there exists $b > 1$ such that*

$$(118) \quad \begin{aligned} S \geq k + 1 + \max(b(d_{1,k} + 2) + 3, d + 1 + b(d + d_{1,k} + 1)) \quad , \quad S \geq k + 3 + b(2 + d_{2,k}), \\ S \geq k + 1 + b \max(d_{1,k}, d_{2,k}) \quad , \quad S \geq k + b d_{3,k} \end{aligned}$$

for all $k \in \mathcal{S}$. For all $0 \leq j \leq S - 1$, we consider functions $\omega_j(t, z)$ which are assumed to be holomorphic and bounded on the product $D(0, R')^2$.

Then, there exist constants $\sigma, \bar{W}, C_{12} > 0$ such that the problem

$$(119) \quad \begin{aligned} \partial_w^S Y(t, z, w) &= \sum_{k \in \mathcal{S}} (b_{1,k}(t, z, X(t, z), w) \partial_t \partial_w^k Y(t, z, w) \\ &+ b_{2,k}(t, z, X(t, z), w) \partial_z \partial_w^k Y(t, z, w) + b_{3,k}(t, z, X(t, z), w) \partial_w^k Y(t, z, w)) \end{aligned}$$

with initial data

$$(120) \quad (\partial_w^j Y)(t, z, 0) = \omega_j(t, z) \quad , \quad 0 \leq j \leq S-1,$$

has a solution $Y(t, z, w)$ which is holomorphic on $\text{Int}(K) \times D(0, \bar{W}/2)$ and which fulfills the following estimates

$$(121) \quad \sup_{(t,z) \in \text{Int}(K), w \in D(0, \bar{W}/2)} |Y(t, z, w)| \leq C_{12} \exp(\sigma \zeta(b) \rho) + \sum_{j=0}^{S-1} \sup_{(t,z) \in \text{Int}(K)} |\omega_j(t, z)| \frac{(\bar{W}/2)^j}{j!}$$

where $\zeta(b) = \sum_{n \geq 0} 1/(n+1)^b$, for any compact set $K \subset D(0, R)^2 \setminus \Theta$ with non-empty interior $\text{Int}(K)$ for some $R < R'$ and any $\rho > 1$ which satisfies (6). We stress that the constants $\sigma, \bar{W}, C_{12} > 0$ do not depend neither on K nor on $\rho > 1$.

Proof By convention, we will put $\omega_j(t, z) \equiv 0$ for all $j \geq S$. On the other hand, we specialize the functions $\tilde{\omega}_\alpha$ which were introduced in (8) in order that

$$(122) \quad \begin{aligned} \tilde{\omega}_\alpha(v_0, v_1, (u_h)_{h \in I(\alpha)}) &= \hat{\omega}_\alpha(v_0, v_1, u_0) \\ &= \sum_{k \in \mathcal{S}} \sum_{\alpha_1 + \alpha_2 = \alpha} \alpha! \left(\frac{b_{1,k,\alpha_1}(v_0, v_1, u_0)}{\alpha_1!} \frac{\partial_{v_0} \omega_{\alpha_2+k}(v_0, v_1)}{\alpha_2!} + \frac{b_{2,k,\alpha_1}(v_0, v_1, u_0)}{\alpha_1!} \frac{\partial_{v_1} \omega_{\alpha_2+k}(v_0, v_1)}{\alpha_2!} \right. \\ &\quad \left. + \frac{b_{3,k,\alpha_1}(v_0, v_1, u_0)}{\alpha_1!} \frac{\omega_{\alpha_2+k}(v_0, v_1)}{\alpha_2!} \right). \end{aligned}$$

By construction and using the definition (15), we can write with the help of the Kronecker symbol,

$$(123) \quad \tilde{\omega}_{\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}} = \hat{\omega}_{\alpha, n_0, n_1, l_0} \times \prod_{h \in I(\alpha) \setminus \{0\}} \delta_{0, l_h}$$

where

$$\hat{\omega}_{\alpha, n_0, n_1, l_0} = \sup_{|v_0| < R, |v_1| < R, |u_0| < \rho} |\partial_{v_0}^{n_0} \partial_{v_1}^{n_1} \partial_{u_0}^{l_0} \hat{\omega}_\alpha(v_0, v_1, u_0)|.$$

Lemma 8 *There exist $\tilde{V}_0, \tilde{V}_1, \tilde{W} > 0$ such that the formal series*

$$(124) \quad \begin{aligned} \tilde{\Omega}(V_0, V_1, (U_h)_{h \geq 0}, W) \\ = \sum_{\alpha \geq 0} \left(\sum_{n_0, n_1, l_h \geq 0, h \in I(\alpha)} \tilde{\omega}_{\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}} \frac{V_0^{n_0}}{n_0!} \frac{V_1^{n_1}}{n_1!} \prod_{h \in I(\alpha)} \frac{U_h^{l_h}}{l_h!} \right) \frac{W^\alpha}{\alpha!} \end{aligned}$$

belongs to $G_{(\rho, \tilde{V}_0, \tilde{V}_1, (\bar{U}_h)_{h \geq 0}, \tilde{W})}$. Moreover, there exists a constant $C_{11} > 0$ (independent of ρ) such that

$$(125) \quad \|\tilde{\Omega}(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \tilde{V}_0, \tilde{V}_1, (\bar{U}_h)_{h \geq 0}, \tilde{W})} \leq C_{11}.$$

Proof Let $k \in \mathcal{S}$. Due to the estimates (10) for the functions $b_{m,k,\alpha}(t, z, u_0)$, we get couples of constants $D_{1,k}, \hat{D}_{1,k} > 0$, $D_{2,k}, \hat{D}_{2,k} > 0$ and $D_{3,k}, \hat{D}_{3,k} > 0$ such that

$$(126) \quad \begin{aligned} |b_{1,k,\alpha_1}(\chi_0, \chi_1, \xi_0)| &\leq D_{1,k}(\rho + \delta)^{d_{1,k}} \alpha_1! (\hat{D}_{1,k})^{\alpha_1}, \\ |b_{2,k,\alpha_1}(\chi_0, \chi_1, \xi_0)| &\leq D_{2,k}(\rho + \delta)^{d_{2,k}} \alpha_1! (\hat{D}_{2,k})^{\alpha_1}, \quad |b_{3,k,\alpha_1}(\chi_0, \chi_1, \xi_0)| \leq D_{3,k}(\rho + \delta)^{d_{3,k}} \alpha_1! (\hat{D}_{3,k})^{\alpha_1} \end{aligned}$$

for all $\alpha_1 \geq 0$, all $|\chi_0| < R + \delta < R'$, $|\chi_1| < R + \delta < R'$, $|\xi_0| < \rho + \delta$. Moreover, we also get couples of constants $E_{1,k}, \hat{E}_{1,k} > 0$, $E_{2,k}, \hat{E}_{2,k} > 0$ and $E_{3,k}, \hat{E}_{3,k} > 0$ such that

$$(127) \quad |\partial_{\chi_0} \omega_{\alpha_2+k}(\chi_0, \chi_1)| \leq E_{1,k} \alpha_2! (\hat{E}_{1,k})^{\alpha_2},$$

$$|\partial_{\chi_1} \omega_{\alpha_2+k}(\chi_0, \chi_1)| \leq E_{2,k} \alpha_2! (\hat{E}_{2,k})^{\alpha_2} \quad , \quad |\omega_{\alpha_2+k}(\chi_0, \chi_1)| \leq E_{3,k} \alpha_2! (\hat{E}_{3,k})^{\alpha_2}$$

for all $\alpha_2 \geq 0$, all $|\chi_0| < R + \delta < R'$, $|\chi_1| < R + \delta < R'$. From (126) and (127) we deduce

$$(128) \quad |\hat{\omega}_\alpha(\chi_0, \chi_1, \xi_0)|$$

$$\leq \sum_{k \in \mathcal{S}} \sum_{\alpha_1 + \alpha_2 = \alpha} \alpha! (D_{1,k} E_{1,k} (\rho + \delta)^{d_{1,k}} (\hat{D}_{1,k})^{\alpha_1} (\hat{E}_{1,k})^{\alpha_2} + D_{2,k} E_{2,k} (\rho + \delta)^{d_{2,k}} (\hat{D}_{2,k})^{\alpha_1} (\hat{E}_{2,k})^{\alpha_2}$$

$$+ D_{3,k} E_{3,k} (\rho + \delta)^{d_{3,k}} (\hat{D}_{3,k})^{\alpha_1} (\hat{E}_{3,k})^{\alpha_2})$$

for all $\alpha \geq 0$, all $|\chi_0| < R + \delta < R'$, $|\chi_1| < R + \delta < R'$, $|\xi_0| < \rho + \delta$. From the Cauchy formula in several variables, one can write

$$\frac{\partial_{v_0}^{n_0} \partial_{v_1}^{n_1} \partial_{u_0}^{l_0} \hat{\omega}_\alpha(v_0, v_1, u_0)}{n_0! n_1! l_0!} = \left(\frac{1}{2i\pi}\right)^3 \int_{C(v_0, \delta)} \int_{C(v_1, \delta)} \int_{C(u_0, \delta)} \hat{\omega}_\alpha(\chi_0, \chi_1, \xi_0)$$

$$\times \frac{d\chi_0 d\chi_1 d\xi_0}{(\chi_0 - v_0)^{n_0+1} (\chi_1 - v_1)^{n_1+1} (\xi_0 - u_0)^{l_0+1}}$$

for all $|v_0| < R$, $|v_1| < R$, $|u_0| < \rho$. We deduce that

$$(129) \quad \frac{\hat{\omega}_{\alpha, n_0, n_1, l_0}}{n_0! n_1! l_0!} \leq \frac{1}{\delta^{n_0 + n_1 + l_0}}$$

$$\times \sum_{k \in \mathcal{S}} \sum_{\alpha_1 + \alpha_2 = \alpha} \alpha! (D_{1,k} E_{1,k} (\rho + \delta)^{d_{1,k}} (\hat{D}_{1,k})^{\alpha_1} (\hat{E}_{1,k})^{\alpha_2} + D_{2,k} E_{2,k} (\rho + \delta)^{d_{2,k}} (\hat{D}_{2,k})^{\alpha_1} (\hat{E}_{2,k})^{\alpha_2}$$

$$+ D_{3,k} E_{3,k} (\rho + \delta)^{d_{3,k}} (\hat{D}_{3,k})^{\alpha_1} (\hat{E}_{3,k})^{\alpha_2})$$

for all $\alpha \geq 0$, all $n_0, n_1, l_0 \geq 0$. Using (123), we get that

$$(130) \quad \|\tilde{\Omega}(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \tilde{V}_0, \tilde{V}_1, (\bar{U}_h)_{h \geq 0}, \tilde{W})}$$

$$= \sum_{\alpha \geq 0} \left(\sum_{n_0, n_1, l_0 \geq 0} \frac{|\hat{\omega}_{\alpha, n_0, n_1, l_0}|}{\exp(\sigma r_b(\alpha) \rho)} \frac{\tilde{V}_0^{n_0} \tilde{V}_1^{n_1} \bar{U}_0^{l_0}}{(n_0 + n_1 + l_0 + \alpha)!} \right) \tilde{W}^\alpha.$$

From (129), (82) and with the help of the classical estimates

$$(n_0 + n_1 + l_0 + \alpha)! \geq n_0! n_1! l_0! \alpha!,$$

for all $n_0, n_1, l_0, \alpha \geq 0$, we get a constant $C_{11,1} > 0$ (depending on $D_{1,k}, d_{1,k}, E_{1,k}, D_{2,k}, d_{2,k}, E_{2,k}, D_{3,k}, d_{3,k}, E_{3,k}$ for all $k \in \mathcal{S}, \sigma, \delta$) such that

$$(131) \quad \|\tilde{\Omega}(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \tilde{V}_0, \tilde{V}_1, (\bar{U}_h)_{h \geq 0}, \tilde{W})}$$

$$\leq \frac{C_{11,1}}{(1 - \frac{\tilde{V}_0}{\delta})(1 - \frac{\tilde{V}_1}{\delta})(1 - \frac{\bar{U}_0}{\delta})} \sum_{k \in \mathcal{S}} \frac{1}{(1 - \hat{D}_{1,k} \tilde{W})(1 - \hat{E}_{1,k} \tilde{W})}$$

$$+ \frac{1}{(1 - \hat{D}_{2,k} \tilde{W})(1 - \hat{E}_{2,k} \tilde{W})} + \frac{1}{(1 - \hat{D}_{3,k} \tilde{W})(1 - \hat{E}_{3,k} \tilde{W})}.$$

We choose

$$(132) \quad 0 < \tilde{W} < \min_{k \in \mathcal{S}} (1/(2\hat{D}_{1,k}), 1/(2\hat{D}_{2,k}), 1/(2\hat{D}_{3,k}), 1/(2\hat{E}_{1,k}), 1/(2\hat{E}_{2,k}), 1/(2\hat{E}_{3,k})),$$

$$0 < \tilde{V}_0 < \delta/2, 0 < \tilde{V}_1 < \delta/2, 0 < \bar{U}_0 < \delta/2.$$

From (131) we deduce the inequality (125). \square

Under the assumption (118), we get from Proposition 10 four constants $0 < \bar{V}_0 < \tilde{V}_0$, $0 < \bar{V}_1 < \tilde{V}_1$, $0 < \bar{U}_0$ and $0 < \bar{W} < \tilde{W}$ (independent of ρ) such that the functional equation

$$(133) \quad \Psi(V_0, V_1, (U_h)_{h \geq 0}, W) = \sum_{k \in \mathcal{S}} (B_{1,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \partial_{V_0} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W))$$

$$+ B_{1,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \mathbb{D}_{\mathbf{A}} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W))$$

$$+ \sum_{k \in \mathcal{S}} (B_{2,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \partial_{V_1} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W))$$

$$+ B_{2,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \mathbb{D}_{\mathbf{B}} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W))$$

$$+ \sum_{k \in \mathcal{S}} B_{3,k}(V_0, V_1, U_0, W) \partial_W^{-S+k} \Psi(V_0, V_1, (U_h)_{h \geq 0}, W)$$

$$+ \tilde{\Omega}(V_0, V_1, (U_h)_{h \geq 0}, W)$$

has a unique solution $\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)$ belonging to $G_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$ which satisfies moreover the estimates

$$(134) \quad \|\Psi(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})} \leq 2 \|\tilde{\Omega}(V_0, V_1, (U_h)_{h \geq 0}, W)\|_{(\rho, \bar{V}_0, \bar{V}_1, (\bar{U}_h)_{h \geq 0}, \bar{W})}$$

$$\leq 2C_{11}.$$

Now, from Proposition 4, we know that the sequence $\varphi_{\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}}$ introduced in (14) satisfies the inequality

$$(135) \quad \varphi_{\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}} \leq \psi_{\alpha, n_0, n_1, (l_h)_{h \in I(\alpha)}}$$

for all $\alpha \geq 0$, all $n_0, n_1, l_h \geq 0$, for $h \in I(\alpha)$. Gathering (134) and (135), and from the definition of the Banach spaces in Section 3.1, we get, in particular, for $n_0 = n_1 = l_h = 0$, for all $h \in I(\alpha)$, all $\alpha \geq 0$, that

$$(136) \quad \sup_{|v_0| < R, |v_1| < R, |u_h| < \rho, h \in I(\alpha)} |\phi_{\alpha}(v_0, v_1, (u_h)_{h \in I(\alpha)})| \leq \psi_{\alpha, 0, 0, (0)_{h \in I(\alpha)}}$$

$$\leq 2C_{11} \exp(\sigma r_b(\alpha) \rho) \left(\frac{1}{\bar{W}}\right)^{\alpha} \alpha! \leq 2C_{11} \exp(\sigma \zeta(b) \rho) \left(\frac{1}{\bar{W}}\right)^{\alpha} \alpha!$$

for all $\alpha \geq 0$ and where $\zeta(b) = \sum_{n \geq 0} 1/(n+1)^b$. From (136), we get that the formal series $U(t, z, w)$ introduced in (7) actually defines a holomorphic function (denoted again by $U(t, z, w)$) on $\text{Int}(K) \times D(0, \bar{W}/2)$ for which the estimates

$$(137) \quad \sup_{(t, z) \in \text{Int}(K), w \in D(0, \bar{W}/2)} |U(t, z, w)| \leq 4C_{11} \exp(\sigma \zeta(b) \rho)$$

hold and which satisfies the equation (13) on $\text{Int}(K) \times D(0, \bar{W}/2)$.

Finally, we define the function

$$Y(t, z, w) = \partial_w^{-S} U(t, z, w) + \sum_{j=0}^{S-1} \omega_j(t, z) \frac{w^j}{j!}.$$

By construction, $Y(t, z, w)$ defines a holomorphic function on $\text{Int}(K) \times D(0, \bar{W}/2)$ with bounds estimates

$$(138) \quad \sup_{(t,z) \in \text{Int}(K), w \in D(0, \bar{W}/2)} |Y(t, z, w)| \leq 4 \left(\frac{\bar{W}}{2}\right)^S C_{11} \exp(\sigma \zeta(b) \rho) \\ + \sum_{j=0}^{S-1} \sup_{(t,z) \in \text{Int}(K)} |\omega_j(t, z)| \frac{(\bar{W}/2)^j}{j!}$$

and solves the problem (119), (120). This yields the result. \square

References

- [1] S. Alinhac, *Problèmes de Cauchy pour des opérateurs singuliers*. (French) Bull. Soc. Math. France 102 (1974), 289–315.
- [2] A. Bove, J. Lewis, C. Parenti, *Propagation of singularities for fuchsian operators*. Lecture Notes in Mathematics, 984. Springer-Verlag, Berlin, 1983. ii+161 pp.
- [3] O. Costin, H. Parkh and Y. Takei, *Borel summability of the heat equation with variable coefficients*, J. Diff. Eq. Volume 252, 4, pp. 3076–3092 (2012).
- [4] L. Debnath, *Nonlinear partial differential equations for scientists and engineers*. Birkhäuser Boston, Inc., Boston, MA, 1997. 593 pp.
- [5] S. Dineen, *Complex analysis on infinite dimensional spaces*, Springer, 1999.
- [6] L. Evans, *Partial differential equations*. Second edition. Graduate Studies in Mathematics, 19. American Mathematical Society, Providence, RI, 2010. 749 pp.
- [7] R. Gérard, H. Tahara, *Singular nonlinear partial differential equations*. Aspects of Mathematics. Friedr. Vieweg and Sohn, Braunschweig, 1996. viii+269 pp.
- [8] Y. Hamada, *The singularities of the solutions of the Cauchy problem*. Publ. Res. Inst. Math. Sci. 5 1969 21–40.
- [9] Y. Hamada, J. Leray, C. Wagschal, *Systèmes d'équations aux dérivées partielles à caractéristiques multiples: problème de Cauchy ramifié; hyperbolicité partielle*. J. Math. Pures Appl. (9) 55 (1976), no. 3, 297–352.
- [10] K. Igari, *On the branching of singularities in complex domains*. Proc. Japan Acad. Ser. A Math. Sci. 70 (1994), no. 5, 128–130.
- [11] M. Kametani, *On multi-valued analytic solutions of first order nonlinear Cauchy problems*. Publ. Res. Inst. Math. Sci. 27 (1991), no. 1, 1–131.

- [12] T. Kobayashi, *Singular solutions and prolongation of holomorphic solutions to nonlinear differential equations*. Publ. Res. Inst. Math. Sci. 34 (1998), no. 1, 43–63.
- [13] T. Kobayashi, *Propagation of singularities for a first order semilinear system in C^{n+1}* . Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 12 (1985), no. 2, 173–189.
- [14] A. Lastra, S. Malek, J. Sanz, *On q -asymptotics for linear q -difference-differential equations with Fuchsian and irregular singularities*. J. Differential Equations 252 (2012), no. 10, 5185–5216.
- [15] J. Leray, *Problème de Cauchy. I. Uniformisation de la solution du problème linéaire analytique de Cauchy près de la variété qui porte les données de Cauchy*. Bull. Soc. Math. France 85 1957 389–429.
- [16] J. Lope, H. Tahara, *On the analytic continuation of solutions to nonlinear partial differential equations*. J. Math. Pures Appl. (9) 81 (2002), no. 9, 811–826.
- [17] S. Malek, *On the summability of formal solutions of linear partial differential equations*. J. Dyn. Control Syst. 11 (2005), no. 3, 389–403.
- [18] S. Malek, *On functional linear partial differential equations in Gevrey spaces of holomorphic functions*. Ann. Fac. Sci. Toulouse Math. (6) 16 (2007), no. 2, 285–302.
- [19] S. Malek, *On Gevrey functional solutions of partial differential equations with Fuchsian and irregular singularities*. J. Dyn. Control Syst. 15 (2009), no. 2, 277–305.
- [20] S. Malek, C. Stenger, *Complex singularity analysis of holomorphic solutions of linear partial differential equations*, Adv. Dyn. Syst. Appl. vol 6. no. 2 (2011).
- [21] W. Malfliet, *Solitary wave solutions of nonlinear wave equations*. Amer. J. Phys. 60 (1992), no. 7, 650–654.
- [22] T. Mandai, *The method of Frobenius to Fuchsian partial differential equations*. J. Math. Soc. Japan 52 (2000), no. 3, 645–672.
- [23] Y. Ohta, J. Satsuma, D. Takahashi, D. Tokihiro, *An elementary introduction to Sato theory*. Recent developments in soliton theory. Progr. Theoret. Phys. Suppl. No. 94 (1988), 210–241.
- [24] S. Ouchi, *An integral representation of singular solutions of linear partial differential equations in the complex domain*. J. Fac. Sci. Univ. Tokyo Sect. IA Math. 27 (1980), no. 1, 37–85.
- [25] S. Ouchi, *The behaviour near the characteristic surface of singular solutions of linear partial differential equations in the complex domain*. Proc. Japan Acad. Ser. A Math. Sci. 65 (1989), no. 4, 102–105.
- [26] B. Sternin, V. Shatalov, *Differential equations on complex manifolds*. Mathematics and its Applications, 276. Kluwer Academic Publishers Group, Dordrecht, 1994. xii+504 pp.
- [27] H. Tahara, *Coupling of two partial differential equations and its applications*, Publ. RIMS. Kyoto Univ. 43 (2007).
- [28] H. Tahara, *On the singularities of solutions of nonlinear partial differential equations in the complex domain. II*. Differential equations and asymptotic theory in mathematical physics, 343–354, Ser. Anal., 2, World Sci. Publ., Hackensack, NJ, 2004.

- [29] H. Tahara, *On the singularities of solutions of nonlinear partial differential equations in the complex domain*. Microlocal analysis and complex Fourier analysis, 273–283, World Sci. Publ., River Edge, NJ, 2002.
- [30] Y. Tsuno, *On the prolongation of local holomorphic solutions of nonlinear partial differential equations*. J. Math. Soc. Japan 27 (1975), no. 3, 454–466.
- [31] C. Wagschal, *Sur le problème de Cauchy ramifié*. J. Math. Pures Appl. (9) 53 (1974), 147–163.