

SCHRÖDINGER OPERATORS WITH δ AND δ' -INTERACTIONS ON LIPSCHITZ SURFACES AND CHROMATIC NUMBERS OF ASSOCIATED PARTITIONS

JUSSI BEHRNDT, PAVEL EXNER, AND VLADIMIR LOTOREICHIK

ABSTRACT. We investigate Schrödinger operators with δ and δ' -interactions supported on hypersurfaces, which separate the Euclidean space into finitely many bounded and unbounded Lipschitz domains. It turns out that the combinatorial properties of the partition and the spectral properties of the corresponding operators are related. As the main result we prove an operator inequality for the Schrödinger operators with δ and δ' -interactions which is based on an optimal colouring and involves the chromatic number of the partition. This inequality implies various relations for the spectra of the Schrödinger operators and, in particular, it allows to transform known results for Schrödinger operators with δ -interactions to Schrödinger operators with δ' -interactions.

1. INTRODUCTION

Schrödinger operators with singular δ -type interactions supported on discrete sets, curves and surfaces are used for the description of quantum mechanical systems with a certain degree of idealization. The spectral properties of Schrödinger operators with δ and δ' -interactions were investigated in numerous mathematical and physical articles in the recent past; we mention only [BN11, KM10, MS12, O10] for interactions on point sets, [CK11, EI01, EK08, EN03, EP12, K12, KV07] on curves, and [AKMN13, BLL13, EF09, EK03] for interactions on surfaces. For a survey and further references we refer the reader to [E08] and to the standard monograph [AGHH].

In this paper we investigate attractive δ and δ' -interactions supported on general hypersurfaces, which separate the Euclidean space \mathbb{R}^d into finitely many bounded and unbounded Lipschitz domains. We establish a connection between the combinatorial properties of these so-called Lipschitz partitions and the relation of the Schrödinger operators with δ and δ' -interactions to each other. More precisely, suppose that the Euclidean space \mathbb{R}^d , $d \geq 2$, is split into a finite number of Lipschitz domains Ω_k , $k = 1, \dots, n$, and let Σ be the union of the boundaries of all Ω_k . The chromatic number χ of the partition is defined as the minimal number of colours, which is sufficient to colour all domains Ω_k in such a way that any two neighbouring domains have distinct colours. In the two dimensional case the famous four colour theorem states that $\chi \leq 4$ for any Lipschitz partition of the plane. In the following the strengths of the δ and δ' -interactions are assumed to be constant along their support Σ , which simplifies the explanation of our results. Let $\alpha \in \mathbb{R}$, $\beta \in \mathbb{R} \setminus \{0\}$ and define the quadratic forms

$$\mathfrak{a}_{\delta, \alpha}[f] := \|\nabla f\|_{L^2(\mathbb{R}^d; \mathbb{C}^d)}^2 - \alpha \|f|_{\Sigma}\|_{L^2(\Sigma)}^2, \quad \text{dom } \mathfrak{a}_{\delta, \alpha} = H^1(\mathbb{R}^d),$$

and

$$\mathfrak{a}_{\delta',\beta}[f] := \sum_{k=1}^n \|\nabla f_k\|_{L^2(\Omega_k; \mathbb{C}^d)}^2 - \sum_{k=1}^{n-1} \sum_{l=k+1}^n \beta^{-1} \|f_k|_{\Sigma_{kl}} - f_l|_{\Sigma_{kl}}\|_{L^2(\Sigma_{kl})}^2,$$

$$\text{dom } \mathfrak{a}_{\delta',\beta} = \bigoplus_{k=1}^n H^1(\Omega_k),$$

where $f_k = f|_{\Omega_k}$ and $\Sigma_{kl} = \partial\Omega_k \cap \partial\Omega_l$, $k \neq l$. It turns out that $\mathfrak{a}_{\delta,\alpha}$ and $\mathfrak{a}_{\delta',\beta}$ are densely defined, closed, symmetric forms in the Hilbert space $L^2(\mathbb{R}^d)$ which are semibounded from below, and hence $\mathfrak{a}_{\delta,\alpha}$ and $\mathfrak{a}_{\delta',\beta}$ induce self-adjoint operators $-\Delta_{\delta,\alpha}$ and $-\Delta_{\delta',\beta}$ in $L^2(\mathbb{R}^d)$. It will be shown in Theorem 3.3 that these operators act as minus Laplacians and the functions in their domains satisfy appropriate δ and δ' -boundary conditions on Σ .

Our main result, Theorem 3.6, is an inequality for the quadratic forms $\mathfrak{a}_{\delta,\alpha}$ and $\mathfrak{a}_{\delta',\beta}$, or equivalently, for the Schrödinger operators $-\Delta_{\delta,\alpha}$ and $-\Delta_{\delta',\beta}$ with δ -interaction of strength α and δ' -interaction of strength β , respectively. Namely, if α , β and the chromatic number χ of the partition satisfy

$$(1.1) \quad 0 < \beta \leq \frac{4}{\alpha} \sin^2(\pi/\chi)$$

then it will be shown that there exists a unitary operator U in $L^2(\mathbb{R}^d)$ such that

$$(1.2) \quad U^{-1}(-\Delta_{\delta',\beta})U \leq -\Delta_{\delta,\alpha}$$

holds. The operator U can be constructed explicitly as soon as the optimal colouring of the partition is provided. The value $4 \sin^2(\pi/\chi)$ in (1.1) pops up as the square of the edge length of the equilateral polygon with χ vertices, which is circumscribed in the unit circle on the complex plane. We also discuss the sharpness of Theorem 3.6 for some cases. First of all it is shown in Example 3.10 that the assumption (1.1) is sharp if $\chi = 2$. In Section 3.4 we then discuss the case $\chi = 3$. It turns out that the weaker assumption $0 < \beta \leq \frac{4}{\alpha}$ (corresponding to $\chi = 2$ in (1.1)) is not sufficient for the existence of a unitary operator U such that (1.2) holds for every partition with $\chi = 3$. This fact will be shown explicitly by considering a symmetric star-graph with three leads as the support of the δ and δ' -interaction.

The inequality (1.2) is particularly useful since it implies various relations of the spectra of $-\Delta_{\delta,\alpha}$ and $-\Delta_{\delta',\beta}$, and it allows to transform known results for Schrödinger operators with δ -interactions to Schrödinger operators with δ' -interactions. We apply our main theorem and its consequences to Lipschitz partitions with compact boundary and so-called locally deformed partitions, where also unbounded Lipschitz domains with unbounded boundaries appear. In these situations we are able to determine or to describe the essential spectra of $-\Delta_{\delta,\alpha}$ and $-\Delta_{\delta',\beta}$, and we derive some consequences on the spectral properties of $-\Delta_{\delta',\beta}$. In particular, it turns out that $-\Delta_{\delta',\beta}$ has a non-empty discrete spectrum if the same holds for $-\Delta_{\delta,\alpha}$, and hence we conclude results on the existence of deformation-induced bound states of $-\Delta_{\delta',\beta}$ from the corresponding results in [EI01, EK03] for the δ -case. We mention that various results on the spectral properties of Schrödinger operators with δ -interactions supported by locally deformed or weakly straight lines and hyperplanes or under more general assumptions of asymptotic flatness exist in the mathematical literature, see, e.g. [CK11, EI01, EK03, EK05, LLP10].

The structure of the paper is as follows. In Section 2 some preliminary facts on the ordering of quadratic forms, Lipschitz partitions, and Sobolev spaces on Lipschitz domains are provided. The quadratic forms $\mathfrak{a}_{\delta,\alpha}$ and $\mathfrak{a}_{\delta',\beta}$, and the corresponding Schrödinger operators $-\Delta_{\delta,\alpha}$ and $-\Delta_{\delta',\beta}$ are introduced and studied in Section 3. This section contains also the main result, Theorem 3.6, and some examples. The more computational aspects in the example of a symmetric star graph with three leads were outsourced in an appendix. The essential spectra and bound states of Schrödinger operators with δ and δ' -interactions on Lipschitz partitions with compact boundary and locally deformed Lipschitz partitions are studied in Section 4.

Acknowledgements. The authors gratefully acknowledge financial support by the Austrian Science Fund (FWF), project P 25162-N26, Czech Science Foundation (GAČR), project P203/11/0701, and the Austria-Czech Republic cooperation grant CZ01/2013.

2. PRELIMINARIES

In this paper we use mainly standard facts from operator theory in Hilbert spaces and basic properties of Sobolev spaces on Lipschitz domains. In this section we briefly recall and define some notions on semibounded sesquilinear forms, Lipschitz partitions and Sobolev spaces.

2.1. Ordering of sesquilinear forms. The self-adjoint operators in this paper are introduced with the help of closed, densely defined, semibounded, symmetric sesquilinear forms via the first representation theorem [K, VI Theorem 2.1]. For a comprehensive introduction into the theory of forms we refer the reader to [K, Chapter VI], [BS87, Chapter 10], and [BEH08, Chapter 4.6].

First we recall the ordering of forms and associated self-adjoint operators.

Definition 2.1. *Let \mathfrak{a}_1 and \mathfrak{a}_2 be closed, densely defined, symmetric sesquilinear forms in a Hilbert space \mathcal{H} and assume that \mathfrak{a}_1 and \mathfrak{a}_2 are bounded from below. Then we shall write $\mathfrak{a}_2 \leq \mathfrak{a}_1$ if*

$$\text{dom } \mathfrak{a}_1 \subset \text{dom } \mathfrak{a}_2 \quad \text{and} \quad \mathfrak{a}_2[f] \leq \mathfrak{a}_1[f] \quad \text{for all } f \in \text{dom } \mathfrak{a}_1.$$

If H_1 and H_2 denote the self-adjoint operators associated with \mathfrak{a}_1 and \mathfrak{a}_2 in \mathcal{H} , respectively, then we write $H_2 \leq H_1$ if and only if $\mathfrak{a}_2 \leq \mathfrak{a}_1$.

We note that by [K, VI Theorem 2.21] two self-adjoint operators H_1 and H_2 which are semibounded from below by ν_1 and ν_2 , respectively, satisfy $H_2 \leq H_1$ if and only if for some, and hence for all, $\nu < \min\{\nu_1, \nu_2\}$

$$(H_2 - \nu)^{-1} - (H_1 - \nu)^{-1} \geq 0.$$

The essential spectrum of a self-adjoint operator H is denoted by $\sigma_{\text{ess}}(H)$. If $\sigma_{\text{ess}}(H) = \emptyset$ we set $\min \sigma_{\text{ess}}(H) = +\infty$ in the following definition.

Definition 2.2. *Let H be a self-adjoint operator in an infinite dimensional Hilbert space and assume that H is bounded from below. We set*

$$N(H) := \#\{\lambda \in (-\infty, \min \sigma_{\text{ess}}(H)) : \lambda \in \sigma_{\text{p}}(H)\} \in \mathbb{N}_0 \cup \{\infty\}$$

and denote by $\{\lambda_k(H)\}_{k=1}^\infty$ the sequence of eigenvalues of H lying below $\min \sigma_{\text{ess}}(H)$, enumerated in non-decreasing order and repeated with multiplicities. In the case $N(H) < \infty$ this sequence is extended by setting $\lambda_{N(H)+k}(H) = \min \sigma_{\text{ess}}(H)$, $k \in \mathbb{N}$.

The statements in the next theorem are consequences of the min-max principle, see, e.g. [BS87, 10.2 Theorem 4] and [RS-IV, Theorem XIII.2].

Theorem 2.3. *Let \mathfrak{a}_1 and \mathfrak{a}_2 be closed, densely defined, symmetric sesquilinear forms in \mathcal{H} which are bounded from below and let H_1 and H_2 be the corresponding self-adjoint operators. Assume that $\mathfrak{a}_2 \leq \mathfrak{a}_1$, or equivalently that $H_2 \leq H_1$. Let $\{\lambda_k(H_i)\}_{k=1}^\infty$ and $N(H_i)$, $i = 1, 2$, be as in Definition 2.2. Then the following statements hold:*

- (i) $\lambda_k(H_2) \leq \lambda_k(H_1)$ for all $k \in \mathbb{N}$;
- (ii) $\min \sigma_{\text{ess}}(H_2) \leq \min \sigma_{\text{ess}}(H_1)$;
- (iii) If $\min \sigma_{\text{ess}}(H_1) = \min \sigma_{\text{ess}}(H_2)$ then $N(H_1) \leq N(H_2)$.

2.2. Lipschitz partitions of Euclidean spaces. In this short subsection we introduce the notion of finite Lipschitz partitions and discuss a combinatorial property of these partitions. For the definition and basic properties of Lipschitz domains we refer the reader to [St, VI.3].

Definition 2.4. *A finite family of Lipschitz domains $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ is called a Lipschitz partition of \mathbb{R}^d , $d \geq 2$, if*

$$\mathbb{R}^d = \bigcup_{k=1}^n \bar{\Omega}_k \quad \text{and} \quad \Omega_k \cap \Omega_l = \emptyset, \quad k, l = 1, 2, \dots, n, \quad k \neq l.$$

The union $\bigcup_{k=1}^n \partial\Omega_k =: \Sigma$ is the boundary of the Lipschitz partition \mathcal{P} . For $k \neq l$ we set $\Sigma_{kl} := \partial\Omega_k \cap \partial\Omega_l$ and we say that Ω_k and Ω_l , $k \neq l$, are neighbouring domains if $\sigma_k(\Sigma_{kl}) > 0$, where σ_k denotes the Lebesgue measure on $\partial\Omega_k$.

The chromatic number of a Lipschitz partition is defined with the help of colouring mappings.

Definition 2.5. *Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a Lipschitz partition of \mathbb{R}^d , $d \geq 2$, with $\Sigma_{kl} = \partial\Omega_k \cap \partial\Omega_l$, $k \neq l$. Then a mapping $\varphi: \{1, 2, \dots, n\} \rightarrow \{0, 1, \dots, m-1\}$ is called an m -colouring for \mathcal{P} if*

$$\sigma_k(\Sigma_{kl}) > 0 \quad \implies \quad \varphi(k) \neq \varphi(l)$$

for all $k, l = 1, 2, \dots, n$, $k \neq l$. The chromatic number χ of the Lipschitz partition \mathcal{P} is defined as

$$\chi := \min\{m \in \mathbb{N}: \exists m\text{-colouring mapping for } \mathcal{P}\}.$$

Thus the chromatic number χ of a Lipschitz partition $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ of \mathbb{R}^d is the minimal number of colours, which is sufficient to colour all domains Ω_k such that any two neighbouring domains have different colours; recall that Ω_k and Ω_l are regarded as neighbouring domains only if the Lebesgue measure of $\Sigma_{kl} = \partial\Omega_k \cap \partial\Omega_l$ is positive. As a famous example we mention the four colour theorem which states that the chromatic number of any Lipschitz partition \mathcal{P} of \mathbb{R}^2 is $\chi \leq 4$.

2.3. Sobolev spaces on arbitrary Lipschitz domains. For a Lipschitz domain Ω we denote the standard L^2 -based Sobolev spaces on Ω and $\partial\Omega$ by $H^s(\Omega)$, $s \in \mathbb{R}$, and $H^t(\partial\Omega)$, $t \in [-1, 1]$, respectively. For the definition and general properties of Sobolev spaces on Lipschitz domains and their boundaries we refer the reader to

[McL] and [St]. Recall that for a Lipschitz domain $\Omega \subset \mathbb{R}^d$ there exists an *extension operator*

$$E: L^2(\Omega) \rightarrow L^2(\mathbb{R}^d)$$

satisfying the following conditions:

- (i) $(Ef) \upharpoonright \Omega = f$ for all $f \in L^2(\Omega)$;
- (ii) $E(H^k(\Omega)) \subset H^k(\mathbb{R}^d)$ for all $k \in \mathbb{N}_0$;
- (iii) $E: H^k(\Omega) \rightarrow H^k(\mathbb{R}^d)$ is continuous for all $k \in \mathbb{N}_0$.

The useful estimate on the trace in the next lemma is essentially a consequence of the continuity of the trace map and the above mentioned properties of the extension operator. For the convenience of the reader we provide a short proof.

Lemma 2.6. *Let $\Omega \subset \mathbb{R}^d$ be a bounded or unbounded Lipschitz domain. Then for any $\varepsilon > 0$ there exists a constant $C(\varepsilon) > 0$ such that*

$$\|f|_{\partial\Omega}\|_{L^2(\partial\Omega)}^2 \leq \varepsilon \|\nabla f\|_{L^2(\Omega; \mathbb{C}^d)}^2 + C(\varepsilon) \|f\|_{L^2(\Omega)}^2$$

holds for all $f \in H^1(\Omega)$.

Proof. Let $f \in H^1(\Omega)$, fix some $s \in (\frac{1}{2}, 1)$ and let $Ef \in H^1(\mathbb{R}^d)$ be the extension of f . The continuity of the trace [M87, Ne] and the properties of the extension operator imply that there exists $c > 0$ such that

$$\|f|_{\partial\Omega}\|_{L^2(\partial\Omega)} \leq c \|f\|_{H^s(\Omega)} \leq c \|Ef\|_{H^s(\mathbb{R}^d)}.$$

Hence for $\varepsilon > 0$ there exists a constant $C_1(\varepsilon) > 0$ such that

$$\|f|_{\partial\Omega}\|_{L^2(\partial\Omega)} \leq c \|Ef\|_{H^s(\mathbb{R}^d)} \leq \varepsilon \|Ef\|_{H^1(\mathbb{R}^d)} + C_1(\varepsilon) \|Ef\|_{L^2(\mathbb{R}^d)},$$

see, e.g. [HT, Theorem 3.30] or [W00, Satz 11.18 (e)]. As E is continuous (see property (iii) for $k = 0$ and $k = 1$) we conclude that for $\varepsilon > 0$ there exists $C_2(\varepsilon) > 0$ such that

$$\|f|_{\partial\Omega}\|_{L^2(\partial\Omega)} \leq \varepsilon \|f\|_{H^1(\Omega)} + C_2(\varepsilon) \|f\|_{L^2(\Omega)}.$$

Thus for $\varepsilon > 0$ there exists a constant $C_3(\varepsilon) > 0$ such that

$$\|f|_{\partial\Omega}\|_{L^2(\partial\Omega)}^2 \leq \varepsilon \|f\|_{H^1(\Omega)}^2 + C_3(\varepsilon) \|f\|_{L^2(\Omega)}^2$$

and hence the assertion follows from $\|f\|_{H^1(\Omega)}^2 = \|\nabla f\|_{L^2(\Omega; \mathbb{C}^d)}^2 + \|f\|_{L^2(\Omega)}^2$. \square

For our purposes it is convenient to define the Laplacian and the Neumann trace in a weak sense in L^2 .

Definition 2.7. *Let Ω be a Lipschitz domain and let $u \in H^1(\Omega)$.*

- (i) *If there exists $f \in L^2(\Omega)$ such that*

$$(\nabla u, \nabla v)_{L^2(\Omega; \mathbb{C}^d)} = (f, v)_{L^2(\Omega)} \quad \text{for all } v \in H_0^1(\Omega)$$

then we define $-\Delta u := f$ and say that $\Delta u \in L^2(\Omega)$.

- (ii) *If $\Delta u \in L^2(\Omega)$ and there exists $b \in L^2(\partial\Omega)$ such that*

$$(\nabla u, \nabla v)_{L^2(\Omega; \mathbb{C}^d)} - (-\Delta u, v)_{L^2(\Omega)} = (b, v|_{\partial\Omega})_{L^2(\partial\Omega)} \quad \text{for all } v \in H^1(\Omega)$$

then we define $\partial_\nu u|_{\partial\Omega} := b$ and say that $\partial_\nu u|_{\partial\Omega} \in L^2(\partial\Omega)$.

We note that Δu and $\partial_\nu u|_{\partial\Omega}$ in the above definition (if they exist) are unique since $H_0^1(\Omega)$ is dense in $L^2(\Omega)$ and the space $\{v|_{\partial\Omega} : v \in H^1(\Omega)\}$ is dense in $L^2(\partial\Omega)$, respectively; cf. [McL, Theorem 3.37].

Definition 2.8. Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a Lipschitz partition of \mathbb{R}^d , $d \geq 2$, with boundary Σ . Let $u \in H^1(\mathbb{R}^d)$, denote the restrictions of u onto Ω_k by u_k and assume that $\Delta u_k \in L^2(\Omega_k)$ for all $k = 1, 2, \dots, n$. If there exists $b \in L^2(\Sigma)$ such that

$$(\nabla u, \nabla v)_{L^2(\mathbb{R}^d; \mathbb{C}^d)} - \left(\bigoplus_{k=1}^n (-\Delta u_k), v \right)_{L^2(\mathbb{R}^d)} = (b, v|_\Sigma)_{L^2(\Sigma)} \quad \text{for all } v \in H^1(\mathbb{R}^d)$$

then we define $\partial_{\mathcal{P}}u|_\Sigma := b$ and say that $\partial_{\mathcal{P}}u|_\Sigma \in L^2(\Sigma)$.

Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a Lipschitz partition with boundary Σ and let $u \in H^1(\mathbb{R}^d)$. As $\{v|_\Sigma : v \in H^1(\mathbb{R}^d)\}$ is dense in $L^2(\Sigma)$ it follows that $\partial_{\mathcal{P}}u|_\Sigma$ (if it exists) is unique.

Remark 2.9. Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a Lipschitz partition of \mathbb{R}^d and assume that $\partial_{\mathcal{P}}u|_\Sigma \in L^2(\Sigma)$ exists for some $u \in H^1(\mathbb{R}^d)$ in the sense of Definition 2.8. Let Ω_k and Ω_l be neighbouring domains and assume that the Neumann traces $\partial_{\nu_k} u_k|_{\partial\Omega_k} \in L^2(\partial\Omega_k)$ and $\partial_{\nu_l} u_l|_{\partial\Omega_l} \in L^2(\partial\Omega_l)$ exist in the sense of Definition 2.7 (ii). Let Γ be a bounded open subset of Σ_{kl} which is part of a Lipschitz dissection in the sense of [McL, page 99] such that $\bar{\Gamma} \cap \Sigma_{km} = \emptyset$ for $m = 1, \dots, n$ with $m \neq l, k$. Then it follows that

$$(2.1) \quad \partial_{\mathcal{P}}u|_\Gamma = \partial_{\nu_k} u_k|_\Gamma + \partial_{\nu_l} u_l|_\Gamma$$

holds. In particular, if $\mathcal{P} = \{\Omega_1, \Omega_2\}$ with $\Omega_2 = \mathbb{R}^d \setminus \bar{\Omega}_1$, boundary $\Sigma = \partial\Omega$ and the Neumann traces exist then

$$\partial_{\mathcal{P}}u|_\Sigma = \partial_{\nu_1} u_1|_\Sigma + \partial_{\nu_2} u_2|_\Sigma.$$

3. SCHRÖDINGER OPERATORS WITH δ AND δ' -INTERACTIONS ASSOCIATED WITH LIPSCHITZ PARTITIONS

In this section we define and study self-adjoint Schrödinger operators with δ and δ' -interactions supported on the boundary Σ of a Lipschitz partition $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ of \mathbb{R}^d , $d \geq 2$. As the main result we prove an operator inequality between the δ and δ' -operator, which implies a certain ordering of their spectra. The key assumption for this inequality is expressed in terms of the chromatic number of the Lipschitz partition.

3.1. Free and Neumann Laplacians. Let in the following $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a Lipschitz partition of \mathbb{R}^d with the boundary Σ . The functions $f \in L^2(\mathbb{R}^d)$ will be decomposed in the form

$$f = \bigoplus_{k=1}^n f_k, \quad f_k := f|_{\Omega_k} \in L^2(\Omega_k), \quad k = 1, 2, \dots, n.$$

The *free Laplacian* $-\Delta_{\text{free}}$ and the *Neumann Laplacian* $-\Delta_{\text{N}}$ with Neumann boundary conditions on Σ are defined as the self-adjoint operators in $L^2(\mathbb{R}^d)$ associated with the sesquilinear forms

$$(3.1) \quad \begin{aligned} \mathfrak{a}_{\text{free}}[f, g] &:= (\nabla f, \nabla g)_{L^2(\mathbb{R}^d; \mathbb{C}^d)}, & \text{dom } \mathfrak{a}_{\text{free}} &:= H^1(\mathbb{R}^d), \\ \mathfrak{a}_{\text{N}}[f, g] &:= \sum_{k=1}^n (\nabla f_k, \nabla g_k)_{L^2(\Omega_k; \mathbb{C}^d)}, & \text{dom } \mathfrak{a}_{\text{N}} &:= \bigoplus_{k=1}^n H^1(\Omega_k), \end{aligned}$$

which are symmetric, closed and semibounded from below, see, e.g. [EE, §VII.1.1-2]. Note that $\text{dom}(-\Delta_{\text{free}}) = H^2(\mathbb{R}^d)$ but the functions in $\text{dom}(-\Delta_{\text{N}})$ have only local H^2 -regularity, that is, $\text{dom}(-\Delta_{\text{N}}) \subset H_{\text{loc}}^2(\mathbb{R}^d \setminus \Sigma)$.

3.2. Definition of Schrödinger operators with δ and δ' -interactions via sesquilinear forms. In this subsection we define Schrödinger operators with δ and δ' -interactions supported on possibly non-compact boundaries of Lipschitz partitions with the help of corresponding sesquilinear forms; cf. [BEKS94] for the case of δ -interactions and [BLL13] for the case of δ' -interactions on smooth hypersurfaces. The domains of these operators are characterized and, in particular, the boundary conditions are given explicitly. For the special case of smooth domains with compact boundaries the present description reduces to the one in [BLL13], where a different approach via extension theory of symmetric operators and boundary triple techniques from [BL07] was used. We also refer to [AKMN13, AGS87, S88] for an approach via separation of variables in the case of spherically symmetric supports of interactions.

Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a Lipschitz partition of \mathbb{R}^d with the boundary Σ , let $\alpha, \beta : \Sigma \rightarrow \mathbb{R}$ be such that $\alpha, \beta^{-1} \in L^\infty(\Sigma)$ and define the symmetric sesquilinear forms $\mathbf{a}_{\delta, \alpha}$ and $\mathbf{a}_{\delta', \beta}$ by

$$(3.2) \quad \mathbf{a}_{\delta, \alpha}[f, g] := (\nabla f, \nabla g)_{L^2(\mathbb{R}^d; \mathbb{C}^d)} - (\alpha f|_\Sigma, g|_\Sigma)_{L^2(\Sigma)}, \quad \text{dom } \mathbf{a}_{\delta, \alpha} = H^1(\mathbb{R}^d),$$

and

$$(3.3) \quad \begin{aligned} \mathbf{a}_{\delta', \beta}[f, g] &:= \sum_{k=1}^n (\nabla f_k, \nabla g_k)_{L^2(\Omega_k; \mathbb{C}^d)} \\ &\quad - \sum_{k=1}^{n-1} \sum_{l=k+1}^n (\beta_{kl}^{-1}(f_k|_{\Sigma_{kl}} - f_l|_{\Sigma_{kl}}), g_k|_{\Sigma_{kl}} - g_l|_{\Sigma_{kl}})_{L^2(\Sigma_{kl})}, \\ \text{dom } \mathbf{a}_{\delta', \beta} &= \bigoplus_{k=1}^n H^1(\Omega_k), \end{aligned}$$

respectively; here $\Sigma_{kl} = \partial\Omega_k \cap \partial\Omega_l$ for $k, l = 1, 2, \dots, n$, $k \neq l$, and β_{kl} denotes the restrictions of β to Σ_{kl} . The traces $f_k|_{\Sigma_{kl}}$ are understood as restrictions of the trace $f_k|_{\partial\Omega_k}$ onto Σ_{kl} . Note that $\sigma_k(\Sigma_{kl}) = \sigma_l(\Sigma_{kl}) = 0$ if the domains Ω_k and Ω_l are not neighbouring and that

$$L^2(\Sigma) = \bigoplus_{k=1}^{n-1} \bigoplus_{l=k+1}^n L^2(\Sigma_{kl}) \quad \text{and} \quad L^2(\partial\Omega_k) = \bigoplus_{l=1, l \neq k}^n L^2(\Sigma_{kl}).$$

Proposition 3.1. *The symmetric sesquilinear forms $\mathbf{a}_{\delta, \alpha}$ and $\mathbf{a}_{\delta', \beta}$ are closed and semibounded from below.*

Proof. We verify the assertion for $\mathbf{a}_{\delta, \alpha}$ first. For this note that $\mathbf{a}_{\delta, \alpha} = \mathbf{a}_{\text{free}} + \mathbf{a}'$, where \mathbf{a}_{free} is as in (3.1) and

$$\mathbf{a}'[f, g] := -(\alpha f, g)_{L^2(\Sigma)}, \quad \text{dom } \mathbf{a}' = H^1(\mathbb{R}^d).$$

We show that \mathbf{a}' is bounded with respect to \mathbf{a}_{free} with form bound < 1 . In fact, for $f = \bigoplus_{k=1}^n f_k \in H^1(\mathbb{R}^d)$ we have

$$(3.4) \quad |\mathbf{a}'[f]| \leq \|\alpha\|_\infty \|f|_\Sigma\|_{L^2(\Sigma)}^2 = \|\alpha\|_\infty \frac{1}{2} \sum_{k=1}^n \|f_k|_{\partial\Omega_k}\|_{L^2(\partial\Omega_k)}^2.$$

According to Lemma 2.6 for any $\varepsilon > 0$ and $k = 1, 2, \dots, n$, there exists $C_k(\varepsilon) > 0$ such that

$$(3.5) \quad \|f_k|_{\partial\Omega_k}\|_{L^2(\partial\Omega_k)}^2 \leq \varepsilon \|\nabla f_k\|_{L^2(\Omega_k; \mathbb{C}^d)}^2 + C_k(\varepsilon) \|f_k\|_{L^2(\Omega_k)}^2.$$

Therefore (3.4) yields

$$\begin{aligned} |\mathbf{a}'[f]| &\leq \|\alpha\|_\infty \frac{\varepsilon}{2} \sum_{k=1}^n \|\nabla f_k\|_{L^2(\Omega_k; \mathbb{C}^d)}^2 + \|\alpha\|_\infty \frac{1}{2} \sum_{k=1}^n C_k(\varepsilon) \|f_k\|_{L^2(\Omega_k)}^2 \\ &\leq \|\alpha\|_\infty \frac{\varepsilon}{2} \mathbf{a}_{\text{free}}[f] + \|\alpha\|_\infty \frac{\max_k C_k(\varepsilon)}{2} \|f\|_{L^2(\mathbb{R}^d)}^2 \end{aligned}$$

for all $f \in \text{dom } \mathbf{a}' = \text{dom } \mathbf{a}_{\text{free}}$. Thus, for sufficiently small ε the form \mathbf{a}' is form bounded with respect to the form \mathbf{a}_{free} with form bound < 1 . Then by [K, VI Theorem 1.33] the form $\mathbf{a}_{\delta, \alpha}$ is closed and semibounded from below.

Next we prove the statement for $\mathbf{a}_{\delta', \beta}$. As above we have $\mathbf{a}_{\delta', \beta} = \mathbf{a}_N + \mathbf{a}''$, where \mathbf{a}_N is as in (3.1) and

$$\begin{aligned} \mathbf{a}''[f, g] &:= - \sum_{k=1}^{n-1} \sum_{l=k+1}^n (\beta_{kl}^{-1} (f_k|_{\Sigma_{kl}} - f_l|_{\Sigma_{kl}}), g_k|_{\Sigma_{kl}} - g_l|_{\Sigma_{kl}})_{L^2(\Sigma_{kl})}, \\ \text{dom } \mathbf{a}'' &= \bigoplus_{k=1}^n H^1(\Omega_k). \end{aligned}$$

We show that \mathbf{a}'' is bounded with respect to \mathbf{a}_N with form bound < 1 . In fact,

$$\begin{aligned} |\mathbf{a}''[f]| &\leq \|\beta^{-1}\|_\infty \sum_{k=1}^{n-1} \sum_{l=k+1}^n \|f_k|_{\Sigma_{kl}} - f_l|_{\Sigma_{kl}}\|_{L^2(\Sigma_{kl})}^2 \\ &\leq 2\|\beta^{-1}\|_\infty \sum_{k=1}^{n-1} \sum_{l=k+1}^n \left(\|f_k|_{\Sigma_{kl}}\|_{L^2(\Sigma_{kl})}^2 + \|f_l|_{\Sigma_{kl}}\|_{L^2(\Sigma_{kl})}^2 \right) \\ &= 2\|\beta^{-1}\|_\infty \sum_{k=1}^n \|f_k|_{\partial\Omega_k}\|_{L^2(\partial\Omega_k)}^2 \end{aligned}$$

and with the help of (3.5) (see Lemma 2.6) we conclude that for any $\varepsilon > 0$ and $k = 1, 2, \dots, n$, there exists $C_k(\varepsilon) > 0$ such that

$$\begin{aligned} |\mathbf{a}''[f]| &\leq 2\varepsilon \|\beta^{-1}\|_\infty \sum_{k=1}^n \|\nabla f_k\|_{L^2(\Omega_k; \mathbb{C}^d)}^2 + 2\|\beta^{-1}\|_\infty \sum_{k=1}^n C_k(\varepsilon) \|f_k\|_{L^2(\Omega_k)}^2 \\ &\leq 2\varepsilon \|\beta^{-1}\|_\infty \mathbf{a}_N[f] + 2\|\beta^{-1}\|_\infty \max_k C_k(\varepsilon) \|f\|_{L^2(\mathbb{R}^d)}^2 \end{aligned}$$

for all $f \in \text{dom } \mathbf{a}'' = \text{dom } \mathbf{a}_N$. Hence for $\varepsilon > 0$ sufficiently small \mathbf{a}'' is bounded with respect to \mathbf{a}_N with form bound < 1 . As above it follows from [K, VI Theorem 1.33] that $\mathbf{a}_{\delta', \beta}$ is closed and semibounded from below. \square

It follows from Proposition 3.1 and the first representation theorem [K, VI Theorem 2.1] that there are unique self-adjoint operators $-\Delta_{\delta, \alpha}$ and $-\Delta_{\delta', \beta}$ in $L^2(\mathbb{R}^d)$ associated with the sesquilinear forms $\mathbf{a}_{\delta, \alpha}$ and $\mathbf{a}_{\delta', \beta}$, respectively, such that

$$(-\Delta_{\delta, \alpha} f, g)_{L^2(\mathbb{R}^d)} = \mathbf{a}_{\delta, \alpha}[f, g] \quad \text{and} \quad (-\Delta_{\delta', \beta} f, g)_{L^2(\mathbb{R}^d)} = \mathbf{a}_{\delta', \beta}[f, g]$$

for $f \in \text{dom } (-\Delta_{\delta, \alpha}) \subset \text{dom } \mathbf{a}_{\delta, \alpha}$, $g \in \text{dom } \mathbf{a}_{\delta, \alpha}$, and $f \in \text{dom } (-\Delta_{\delta', \beta}) \subset \text{dom } \mathbf{a}_{\delta', \beta}$, $g \in \text{dom } \mathbf{a}_{\delta', \beta}$, respectively.

Definition 3.2. *The selfadjoint operator $-\Delta_{\delta, \alpha}$ ($-\Delta_{\delta', \beta}$) in $L^2(\mathbb{R}^d)$ is called Schrödinger operator with δ -interaction of strength α (δ' -interaction of strength β , respectively) supported on Σ .*

Observe that by the definition of the forms $\mathbf{a}_{\delta,\alpha}$ and $\mathbf{a}_{\delta',\beta}$ the δ -interaction is strong if α is big, and the δ' -interaction is strong if β is small.

In the next theorem we characterize the action and domain of $-\Delta_{\delta,\alpha}$ and $-\Delta_{\delta',\beta}$.

Theorem 3.3. *Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a Lipschitz partition of \mathbb{R}^d with the boundary Σ , let $\alpha, \beta : \Sigma \rightarrow \mathbb{R}$ be such that $\alpha, \beta^{-1} \in L^\infty(\Sigma)$ and let $-\Delta_{\delta,\alpha}$ and $-\Delta_{\delta',\beta}$ be the self-adjoint operators associated with $\mathbf{a}_{\delta,\alpha}$ and $\mathbf{a}_{\delta',\beta}$, respectively. Then the following holds.*

- (i) $-\Delta_{\delta,\alpha}f = \oplus_{k=1}^n(-\Delta f_k)$ and $f = \oplus_{k=1}^n f_k \in \text{dom}(-\Delta_{\delta,\alpha})$ if and only if
 - (a) $f \in H^1(\mathbb{R}^d)$,
 - (b) $\Delta f_k \in L^2(\Omega_k)$ for all $k = 1, 2, \dots, n$,
 - (c) $\partial_{\mathcal{P}}f|_{\Sigma} \in L^2(\Sigma)$ exists in the sense of Definition 2.8 and

$$\partial_{\mathcal{P}}f|_{\Sigma} = \alpha f|_{\Sigma}.$$
- (ii) $-\Delta_{\delta',\beta}f = \oplus_{k=1}^n(-\Delta f_k)$ and $f = \oplus_{k=1}^n f_k \in \text{dom}(-\Delta_{\delta',\beta})$ if and only if
 - (a') $f_k \in H^1(\Omega_k)$ for all $k = 1, 2, \dots, n$,
 - (b') $\Delta f_k \in L^2(\Omega_k)$ for all $k = 1, 2, \dots, n$,
 - (c') $\partial_{\nu_k} f_k|_{\partial\Omega_k} \in L^2(\partial\Omega_k)$ for all $k = 1, 2, \dots, n$ in the sense of Definition 2.7 (ii) and

$$f_k|_{\partial\Omega_k} - \oplus_{l \neq k} f_l|_{\Sigma_{kl}} = \beta_k \partial_{\nu_k} f_k|_{\partial\Omega_k}, \quad k = 1, 2, \dots, n.$$

Proof. The proof of items (i) and (ii) consists of three steps each. First we show that $-\Delta_{\delta,\alpha}$ and $-\Delta_{\delta',\beta}$ act as minus Laplacians on each Ω_k . In the second step we verify that $f \in \text{dom}(-\Delta_{\delta,\alpha})$ ($f \in \text{dom}(-\Delta_{\delta',\beta})$) satisfies the conditions (a)–(c) ((a')–(c')), respectively, and in the last step we prove the converse implication.

(i) **Step I.** Let $f \in \text{dom}(-\Delta_{\delta,\alpha})$, $g_k \in H_0^1(\Omega_k)$ for some $k = 1, 2, \dots, n$, and extend g_k by zero to $\tilde{g}_k \in H^1(\mathbb{R}^d) = \text{dom } \mathbf{a}_{\delta,\alpha}$. From $\tilde{g}_k|_{\Sigma} = 0$ and the first representation theorem we obtain

$$\begin{aligned} ((-\Delta_{\delta,\alpha}f)_k, g_k)_{L^2(\Omega_k)} &= (-\Delta_{\delta,\alpha}f, \tilde{g}_k)_{L^2(\mathbb{R}^d)} = \mathbf{a}_{\delta,\alpha}[f, \tilde{g}_k] \\ &= (\nabla f, \nabla \tilde{g}_k)_{L^2(\mathbb{R}^d; \mathbb{C}^d)} - (\alpha f|_{\Sigma}, \tilde{g}_k|_{\Sigma})_{L^2(\Sigma)} \\ &= (\nabla f, \nabla \tilde{g}_k)_{L^2(\mathbb{R}^d; \mathbb{C}^d)} = (\nabla f_k, \nabla g_k)_{L^2(\Omega_k; \mathbb{C}^d)}. \end{aligned}$$

Therefore, by Definition 2.7 (i) we have $(-\Delta_{\delta,\alpha}f)_k = -\Delta f_k \in L^2(\Omega_k)$ for all $k = 1, 2, \dots, n$, that is,

$$-\Delta_{\delta,\alpha}f = \oplus_{k=1}^n(-\Delta f_k) \in L^2(\mathbb{R}^d).$$

Step II. Let f be a function in $\text{dom}(-\Delta_{\delta,\alpha})$. Then f satisfies condition (a) since $\text{dom}(-\Delta_{\delta,\alpha}) \subset \text{dom } \mathbf{a}_{\delta,\alpha} = H^1(\mathbb{R}^d)$. Condition (b) is satisfied as we have shown in Step I. Hence it remains to check condition (c). For this let $h \in \text{dom } \mathbf{a}_{\delta,\alpha}$. From Step I and the first representation theorem we conclude

$$\begin{aligned} (\oplus_{k=1}^n(-\Delta f_k), h)_{L^2(\mathbb{R}^d)} &= (-\Delta_{\delta,\alpha}f, h)_{L^2(\mathbb{R}^d)} = \mathbf{a}_{\delta,\alpha}[f, h] \\ &= (\nabla f, \nabla h)_{L^2(\mathbb{R}^d; \mathbb{C}^d)} - (\alpha f|_{\Sigma}, h|_{\Sigma})_{L^2(\Sigma)} \end{aligned}$$

which yields

$$(\nabla f, \nabla h)_{L^2(\mathbb{R}^d; \mathbb{C}^d)} - (\oplus_{k=1}^n(-\Delta f_k), h)_{L^2(\mathbb{R}^d)} = (\alpha f|_{\Sigma}, h|_{\Sigma})_{L^2(\Sigma)}.$$

Hence, by Definition 2.8 we have $\partial_{\mathcal{P}}f|_{\Sigma} \in L^2(\Sigma)$ and $\partial_{\mathcal{P}}f|_{\Sigma} = \alpha f|_{\Sigma}$, that is, f satisfies condition (c).

Step III. Assume that f satisfies the conditions (a)-(c) and let $h \in \text{dom } \mathbf{a}_{\delta, \alpha}$. By condition (a) we have $f \in \text{dom } \mathbf{a}_{\delta, \alpha}$ and hence

$$\mathbf{a}_{\delta, \alpha}[f, h] = (\nabla f, \nabla h)_{L^2(\mathbb{R}^d; \mathbb{C}^d)} - (\alpha f|_{\Sigma}, h|_{\Sigma})_{L^2(\Sigma)}.$$

The conditions (b) and (c) together with Definition 2.8 imply that

$$(\nabla f, \nabla h)_{L^2(\mathbb{R}^d; \mathbb{C}^d)} - \left(\bigoplus_{k=1}^n (-\Delta f_k), h \right)_{L^2(\mathbb{R}^d)} = (\partial_{\mathcal{P}}f|_{\Sigma}, h|_{\Sigma})_{L^2(\Sigma)} = (\alpha f|_{\Sigma}, h|_{\Sigma})_{L^2(\Sigma)}$$

and hence $\mathbf{a}_{\delta, \alpha}[f, h] = \left(\bigoplus_{k=1}^n (-\Delta f_k), h \right)_{L^2(\mathbb{R}^d)}$ for all $h \in \text{dom } \mathbf{a}_{\delta, \alpha}$. The first representation theorem yields $f \in \text{dom } (-\Delta_{\delta, \alpha})$.

(ii) **Step I.** The same reasoning as in (i) Step I yields $-\Delta_{\delta', \beta}f = \bigoplus_{k=1}^n (-\Delta f_k) \in L^2(\mathbb{R}^d)$ for all $f \in \text{dom } (-\Delta_{\delta', \beta})$.

Step II. Let f be a function in $\text{dom } (-\Delta_{\delta', \beta})$. Then f satisfies condition (a') since $\text{dom } (-\Delta_{\delta', \beta}) \subset \text{dom } \mathbf{a}_{\delta', \beta}$ and condition (b') holds by Step I. We check condition (c'). For this let $h_k \in H^1(\Omega_k)$ for some $k = 1, 2, \dots, n$, and let $\tilde{h}_k \in \text{dom } \mathbf{a}_{\delta', \beta}$ be its extension by zero. From Step I and the first representation theorem we conclude

$$(3.6) \quad \begin{aligned} & (-\Delta f_k, h_k)_{L^2(\Omega_k)} = (-\Delta_{\delta', \beta}f, \tilde{h}_k)_{L^2(\mathbb{R}^d)} = \mathbf{a}_{\delta', \beta}[f, \tilde{h}_k] \\ & = (\nabla f_k, \nabla h_k)_{L^2(\Omega_k; \mathbb{C}^d)} - \sum_{l=1, l \neq k}^n (\beta_{kl}^{-1}(f_k|_{\Sigma_{kl}} - f_l|_{\Sigma_{kl}}), h_k|_{\Sigma_{kl}})_{L^2(\Sigma_{kl})} \end{aligned}$$

where we used that $\tilde{h}_k|_{\Sigma_{pq}} = 0$ if $k \neq p, q$. For $k = 1, \dots, n$ we set

$$b_k := \bigoplus_{l=1, l \neq k}^n (\beta_{kl}^{-1}(f_k|_{\Sigma_{kl}} - f_l|_{\Sigma_{kl}})) \in \bigoplus_{l=1, l \neq k}^n L^2(\Sigma_{kl}) = L^2(\partial\Omega_k).$$

From (3.6) we then obtain

$$(\nabla f_k, \nabla h_k)_{L^2(\Omega_k; \mathbb{C}^d)} - (-\Delta f_k, h_k)_{L^2(\Omega_k)} = (b_k, h_k|_{\partial\Omega_k})_{L^2(\partial\Omega_k)}$$

for all $h_k \in H^1(\Omega_k)$ and $k = 1, \dots, n$. Hence, $\partial_{\nu_k}f_k|_{\partial\Omega_k} \in L^2(\partial\Omega_k)$ exists in the sense of Definition 2.7 (ii) and the boundary condition

$$\beta_k \partial_{\nu_k}f_k|_{\partial\Omega_k} = \beta_k b_k = f_k|_{\partial\Omega_k} - \bigoplus_{l=1, l \neq k}^n f_l|_{\Sigma_{kl}}, \quad k = 1, \dots, n,$$

holds, that is, condition (c') is valid for all $f \in \text{dom } (-\Delta_{\delta', \beta})$.

Step III. Assume that f satisfies conditions (a')-(c'), and let $h \in \text{dom } \mathbf{a}_{\delta', \beta}$. Fix some $k = 1, \dots, n$ and let \tilde{h}_k be the extension of $h_k \in H^1(\Omega_k)$ by zero. By condition (a') $f \in \text{dom } \mathbf{a}_{\delta', \beta}$ and hence

$$\mathbf{a}_{\delta', \beta}[f, \tilde{h}_k] = (\nabla f_k, \nabla h_k)_{L^2(\Omega_k; \mathbb{C}^d)} - \sum_{l=1, l \neq k}^n (\beta_{kl}^{-1}(f_k|_{\Sigma_{kl}} - f_l|_{\Sigma_{kl}}), h_k|_{\Sigma_{kl}})_{L^2(\Sigma_{kl})}.$$

On the other hand Definition 2.7 (ii) and conditions (b') and (c') imply

$$\begin{aligned} & (\nabla f_k, \nabla h_k)_{L^2(\Omega_k; \mathbb{C}^d)} - (-\Delta f_k, h_k)_{L^2(\Omega_k)} \\ & = (\beta_k^{-1}(f_k|_{\partial\Omega_k} - \bigoplus_{l \neq k} f_l|_{\Sigma_{kl}}), h_k|_{\partial\Omega_k})_{L^2(\partial\Omega_k)} \end{aligned}$$

and hence $\mathbf{a}_{\delta',\beta}[f, \tilde{h}_k] = (-\Delta f_k, h_k)_{L^2(\Omega_k)}$ for $k = 1, 2, \dots, n$. Summing up we conclude

$$\mathbf{a}_{\delta',\beta}[f, h] = \sum_{k=1}^n \mathbf{a}_{\delta',\beta}[f, \tilde{h}_k] = \sum_{k=1}^n (-\Delta f_k, h_k)_{L^2(\Omega_k)} = (\oplus_{k=1}^n (-\Delta f_k), h)_{L^2(\mathbb{R}^d)}$$

for any $h \in \text{dom } \mathbf{a}_{\delta',\beta}$. This implies $f \in \text{dom } (-\Delta_{\delta',\beta})$. \square

We remark that the condition $\partial_{\mathcal{P}} f|_{\Sigma} = \alpha f|_{\Sigma}$ for the functions in $\text{dom } (-\Delta_{\delta,\alpha})$ in Theorem 3.3 (i)-(c) reflects the classical δ -jump boundary conditions on common boundaries of the domains in the partition; cf. [AGHH, I. Theorem 3.1.1] and Remark 2.9. Similarly the condition

$$f_k|_{\partial\Omega_k} - \oplus_{l \neq k} f_l|_{\Sigma_{kl}} = \beta_k \partial_{\nu_k} f_k|_{\partial\Omega_k}, \quad k = 1, 2, \dots, n,$$

for the functions in $\text{dom } (-\Delta_{\delta',\beta})$ in Theorem 3.3 (ii)-(c') corresponds to the classical δ' -jump boundary conditions; cf. [AGHH, I. equation (4.5)]. Note also that our sign choice for α and β in the definition of the forms in (3.2)-(3.3) and the associated operators is opposite with respect to [AGHH].

Observe that for a function f in $\text{dom } (-\Delta_{\delta,\alpha})$ or $\text{dom } (-\Delta_{\delta',\beta})$ it follows from Theorem 3.3 (i)-(b), (ii)-(b') and elliptic regularity that

$$f_k \in H_{\text{loc}}^2(\Omega_k), \quad k = 1, \dots, n.$$

It is not surprising that additional assumptions on the smoothness of the boundary (or parts of the boundary) and the coefficients α, β^{-1} lead to H^2 -regularity of f up to the boundary (or parts of it, respectively). We first recall a result from [BLL13] for a particular smooth partition and turn to a more general situation in the lemma below.

Proposition 3.4. *Let Ω be a bounded domain with C^∞ -boundary and consider the partition $\mathcal{P} = \{\Omega, \mathbb{R}^d \setminus \bar{\Omega}\}$ with boundary $\Sigma = \partial\Omega$. Then the following holds.*

- (i) *If $\alpha \in L^\infty(\Sigma)$ then both domains $\text{dom } (-\Delta_{\delta,\alpha})$ and $\text{dom } (-\Delta_{\delta',\beta})$ are contained in $H^{3/2}(\Omega) \oplus H^{3/2}(\mathbb{R}^d \setminus \bar{\Omega})$.*
- (ii) *If $\alpha \in W^{1,\infty}(\Sigma)$ then both domains $\text{dom } (-\Delta_{\delta,\alpha})$ and $\text{dom } (-\Delta_{\delta',\beta})$ are contained in $H^2(\Omega) \oplus H^2(\mathbb{R}^d \setminus \bar{\Omega})$.*

In the next lemma we establish local H^2 -regularity up to parts of the boundary Σ of a Lipschitz partition $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ under the assumption that the corresponding part of the boundary and $\alpha, \beta^{-1} \in L^\infty(\Sigma)$ are locally $C^{1,1}$ and C^1 , respectively. This observation, which is essentially a consequence of the boundary conditions in Theorem 3.3 (i)-(c), (ii)-(c') and [McL, Theorem 4.18, Theorem 4.20], will be used in the proof of Theorem 4.7.

Lemma 3.5. *Let Ω_k and Ω_l be neighbouring domains of a Lipschitz partition $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ and let Γ be a bounded open subset of $\Sigma_{kl} = \partial\Omega_k \cap \partial\Omega_l$ which is $C^{1,1}$ and part of a Lipschitz dissection of $\partial\Omega_k$ in the sense of [McL, page 99]. Assume that $\Gamma \cap \Sigma_{km} = \emptyset$ for $m = 1, \dots, n$ with $m \neq l, k$. Then for any relatively open subset γ of Γ with $\bar{\gamma} \subset \Gamma$ there exists an open set G_{kl} of $\Omega_k \cup \Omega_l \cup \Gamma$ such that $\bar{\gamma} \subset G_{kl}$ and the following holds for $j = k, l$.*

- (i) *If $\alpha|_{\Gamma} \in C^1(\Gamma)$ and $f \in \text{dom } (-\Delta_{\delta,\alpha})$ then $f_j \in H^2(\Omega_j \cap G_{kl})$;*
- (ii) *If $\beta^{-1}|_{\Gamma} \in C^1(\Gamma)$ and $f \in \text{dom } (-\Delta_{\delta',\beta})$ then $f_j \in H^2(\Omega_j \cap G_{kl})$.*

Proof. (i) For $f \in \text{dom}(-\Delta_{\delta,\alpha})$ we have $f_k \in H^1(\Omega_k)$ and hence $f_k|_\Gamma \in H^{1/2}(\Gamma)$. Therefore $\alpha|_\Gamma \in C^1(\Gamma)$ yields $\alpha f_k|_\Gamma \in H^{1/2}(\Gamma)$ by [McL, Theorem 3.20]. The boundary condition in Theorem 3.3 (i)-(c) and its local form in (2.1) (interpreted in $H^{-1/2}(\Gamma)$ if the Neumann traces $\partial_{\nu_k} u_k|_\Gamma$ and $\partial_{\nu_l} u_l|_\Gamma$ do not exist in $L^2(\Gamma)$) together with [McL, Theorem 4.20] implies the statement.

(ii) As in the proof of (i) we have $f_k|_\Gamma \in H^{1/2}(\Gamma)$ for $f \in \text{dom}(-\Delta_{\delta',\beta})$ and the assumption $\beta^{-1}|_\Gamma \in C^1(\Gamma)$ together with Theorem 3.3 (ii)-(c') and [McL, Theorem 3.20] implies $\beta^{-1}(f_k|_\Gamma - f_l|_\Gamma) = \partial_{\nu_k} f_k|_\Gamma \in H^{1/2}(\Gamma)$. Now the assertion follows from [McL, Theorem 4.18 (ii)]. \square

3.3. An operator inequality for Schrödinger operators with δ and δ' -interactions. Let again $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a Lipschitz partition of \mathbb{R}^d with boundary Σ , and let $\alpha, \beta : \Sigma \rightarrow \mathbb{R}$ be such that $\alpha, \beta^{-1} \in L^\infty(\Sigma)$. In Theorem 3.6 below we prove an operator inequality for the Schrödinger operators $-\Delta_{\delta,\alpha}$ and $-\Delta_{\delta',\beta}$ which is intimately related with the chromatic number χ of the partition \mathcal{P} .

Theorem 3.6. *Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a Lipschitz partition of \mathbb{R}^d with boundary Σ and chromatic number χ . Let $\alpha, \beta : \Sigma \rightarrow \mathbb{R}$ be such that $\alpha, \beta^{-1} \in L^\infty(\Sigma)$ and assume that*

$$(3.7) \quad 0 < \beta \leq \frac{4}{\alpha} \sin^2(\pi/\chi).$$

Then there exists a unitary operator $U : L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$ such that the self-adjoint operators $-\Delta_{\delta,\alpha}$ and $-\Delta_{\delta',\beta}$ satisfy the inequality

$$U^{-1}(-\Delta_{\delta',\beta})U \leq -\Delta_{\delta,\alpha}.$$

Proof. By the definition of the chromatic number (Definition 2.5) there exists an optimal colouring mapping

$$\phi : \{1, 2, \dots, n\} \rightarrow \{0, 1, \dots, \chi - 1\}$$

such that for any $k, l = 1, 2, \dots, n$, $k \neq l$, we have

$$\sigma_k(\Sigma_{kl}) > 0 \implies \phi(k) \neq \phi(l).$$

Next, we define n complex numbers $\mathcal{Z} := \{z_k\}_{k=1}^n$ on the unit circle by

$$z_k := \exp\left(\frac{2\pi\phi(k)}{\chi}i\right), \quad k = 1, 2, \dots, n.$$

Among the z_k there are only χ distinct numbers. The points z_k , $k = 1, \dots, n$, on the unit circle form the vertices of an equilateral polygon with χ edges. The square of the length of these edges is

$$(3.8) \quad 2 - 2 \cos(2\pi/\chi) = 4 \sin^2(\pi/\chi).$$

Observe that for any $k, l = 1, 2, \dots, n$, $k \neq l$, with $\sigma_k(\Sigma_{kl}) > 0$ we have

$$\begin{aligned} |z_k - z_l|^2 &= \left[\cos\left(\frac{2\pi\phi(k)}{\chi}\right) - \cos\left(\frac{2\pi\phi(l)}{\chi}\right) \right]^2 + \left[\sin\left(\frac{2\pi\phi(k)}{\chi}\right) - \sin\left(\frac{2\pi\phi(l)}{\chi}\right) \right]^2 \\ &= 2 - 2 \cos\left(\frac{2\pi\phi(k)}{\chi}\right) \cos\left(\frac{2\pi\phi(l)}{\chi}\right) - 2 \sin\left(\frac{2\pi\phi(k)}{\chi}\right) \sin\left(\frac{2\pi\phi(l)}{\chi}\right) \\ &= 2 - 2 \cos\left(\frac{2\pi}{\chi}(\phi(k) - \phi(l))\right) \geq 2 - 2 \cos(2\pi/\chi), \end{aligned}$$

where we used standard trigonometric identities in the third equality and

$$\phi(k) - \phi(l) \in \{-(\chi - 1), \dots, \chi - 1\} \setminus \{0\}$$

in the last estimate. Together with (3.8) we find $4 \sin^2(\pi/\chi) \leq |z_k - z_l|^2$ and hence by the assumption (3.7)

$$(3.9) \quad 0 < \alpha \leq \frac{4}{\beta} \sin^2(\pi/\chi) \leq \alpha_{\mathcal{Z}},$$

where $\alpha_{\mathcal{Z}}(x) = |z_k - z_l|^2 \beta_{kl}^{-1}(x)$, $x \in \Sigma_{kl}$, and $k, l = 1, 2, \dots, n$, $k \neq l$.

Define a unitary mapping $U_{\mathcal{Z}}: L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$ by

$$(U_{\mathcal{Z}}f)(x) := z_k f_k(x), \quad x \in \Omega_k, \quad k = 1, \dots, n,$$

and a corresponding sesquilinear form $\tilde{\mathbf{a}}_{\delta', \beta}$ by

$$\tilde{\mathbf{a}}_{\delta', \beta}[f, g] := \mathbf{a}_{\delta', \beta}[U_{\mathcal{Z}}f, U_{\mathcal{Z}}g], \quad \text{dom } \tilde{\mathbf{a}}_{\delta', \beta} = \text{dom } \mathbf{a}_{\delta', \beta}.$$

Observe that $\tilde{\mathbf{a}}_{\delta', \beta}$ is a closed, densely defined, symmetric form which is semi-bounded from below, and that the selfadjoint operator associated with $\tilde{\mathbf{a}}_{\delta', \beta}$ is

$$U_{\mathcal{Z}}^{-1}(-\Delta_{\delta', \beta})U_{\mathcal{Z}}, \quad \text{dom}(U_{\mathcal{Z}}^{-1}(-\Delta_{\delta', \beta})U_{\mathcal{Z}}) = U_{\mathcal{Z}}^{-1}(\text{dom}(-\Delta_{\delta', \beta})).$$

We claim that the inequality $\tilde{\mathbf{a}}_{\delta', \beta} \leq \mathbf{a}_{\delta, \alpha_{\mathcal{Z}}}$ holds. In fact,

$$\text{dom } \mathbf{a}_{\delta, \alpha_{\mathcal{Z}}} = H^1(\mathbb{R}^d) \subset \bigoplus_{k=1}^n H^1(\Omega_k) = \text{dom } \tilde{\mathbf{a}}_{\delta', \beta}$$

is clear and for $f \in \text{dom } \mathbf{a}_{\delta, \alpha_{\mathcal{Z}}}$ we have $f|_{\Sigma_{kl}} = f_k|_{\Sigma_{kl}} = f_l|_{\Sigma_{kl}}$. Therefore we obtain

$$\begin{aligned} \tilde{\mathbf{a}}_{\delta', \beta}[f] &= \mathbf{a}_{\delta', \beta}[U_{\mathcal{Z}}f] \\ &= \sum_{k=1}^n \|z_k \nabla f_k\|_{L^2(\Omega_k; \mathbb{C}^d)}^2 - \sum_{k=1}^{n-1} \sum_{l=k+1}^n |z_k - z_l|^2 (\beta_{kl}^{-1} f|_{\Sigma_{kl}}, f|_{\Sigma_{kl}})_{L^2(\Sigma_{kl})} \\ &= \|\nabla f\|_{L^2(\mathbb{R}^d; \mathbb{C}^d)}^2 - (\alpha_{\mathcal{Z}} f|_{\Sigma}, f|_{\Sigma})_{L^2(\Sigma)} = \mathbf{a}_{\delta, \alpha_{\mathcal{Z}}}[f] \end{aligned}$$

for all $f \in \text{dom } \mathbf{a}_{\delta, \alpha_{\mathcal{Z}}}$, and hence $\tilde{\mathbf{a}}_{\delta', \beta} \leq \mathbf{a}_{\delta, \alpha_{\mathcal{Z}}}$. Moreover, as $\alpha \leq \alpha_{\mathcal{Z}}$ by (3.9) we also have $\mathbf{a}_{\delta, \alpha_{\mathcal{Z}}} \leq \mathbf{a}_{\delta, \alpha}$. This implies

$$\tilde{\mathbf{a}}_{\delta', \beta} \leq \mathbf{a}_{\delta, \alpha}$$

and hence $U_{\mathcal{Z}}^{-1}(-\Delta_{\delta', \beta})U_{\mathcal{Z}} \leq -\Delta_{\delta, \alpha}$. \square

As an immediate consequence of Theorem 3.6 and Theorem 2.3 we obtain the following corollary on the relation of the spectra of $-\Delta_{\delta, \alpha}$ and $-\Delta_{\delta', \beta}$.

Corollary 3.7. *Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a Lipschitz partition of \mathbb{R}^d with boundary Σ . Let $\alpha, \beta: \Sigma \rightarrow \mathbb{R}$ be such that $\alpha, \beta^{-1} \in L^\infty(\Sigma)$ and assume that*

$$0 < \beta \leq \frac{4}{\alpha} \sin^2(\pi/\chi).$$

Denote by $\{\lambda_k(-\Delta_{\delta, \alpha})\}_{k=1}^\infty$ and $\{\lambda_k(-\Delta_{\delta', \beta})\}_{k=1}^\infty$ the eigenvalues of the operators $-\Delta_{\delta, \alpha}$ and $-\Delta_{\delta', \beta}$, respectively, below the bottom of their essential spectra, enumerated in non-decreasing order and repeated with multiplicities, and let $N(-\Delta_{\delta, \alpha})$ and $N(-\Delta_{\delta', \beta})$ be their total numbers as in Definition 2.2. Then the following holds.

- (i) $\lambda_k(-\Delta_{\delta', \beta}) \leq \lambda_k(-\Delta_{\delta, \alpha})$ for all $k \in \mathbb{N}$;
- (ii) $\min \sigma_{\text{ess}}(-\Delta_{\delta', \beta}) \leq \min \sigma_{\text{ess}}(-\Delta_{\delta, \alpha})$;
- (iii) If $\min \sigma_{\text{ess}}(-\Delta_{\delta, \alpha}) = \min \sigma_{\text{ess}}(-\Delta_{\delta', \beta})$ then $N(-\Delta_{\delta, \alpha}) \leq N(-\Delta_{\delta', \beta})$.

According to the four colour theorem the chromatic number of a Lipschitz partition \mathcal{P} of \mathbb{R}^2 is $\chi \leq 4$; cf. [AH77, AHK77] or [MT01, §8.2]. This implies the following corollary in the case $d = 2$.

Corollary 3.8. *Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a Lipschitz partition of \mathbb{R}^2 with boundary Σ and chromatic number χ . Let $\alpha, \beta: \Sigma \rightarrow \mathbb{R}$ be such that $\alpha, \beta^{-1} \in L^\infty(\Sigma)$ and assume that*

$$0 < \beta \leq \frac{2}{\alpha}.$$

Then there exists a unitary operator $U: L^2(\mathbb{R}^2) \rightarrow L^2(\mathbb{R}^2)$ such that the self-adjoint operators $-\Delta_{\delta, \alpha}$ and $-\Delta_{\delta', \beta}$ satisfy the inequality

$$U^{-1}(-\Delta_{\delta', \beta})U \leq -\Delta_{\delta, \alpha},$$

and hence the assertions in Corollary 3.7 hold.

For the case of a Lipschitz partition with chromatic number $\chi = 2$ Theorem 3.6 reads as follows.

Corollary 3.9. *Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a Lipschitz partition of \mathbb{R}^d with boundary Σ and chromatic number $\chi = 2$. Let $\alpha, \beta: \Sigma \rightarrow \mathbb{R}$ be such that $\alpha, \beta^{-1} \in L^\infty(\Sigma)$ and assume that*

$$0 < \beta \leq \frac{4}{\alpha}.$$

Then there exists a unitary operator $U: L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$ such that the self-adjoint operators $-\Delta_{\delta, \alpha}$ and $-\Delta_{\delta', \beta}$ satisfy the operator inequality

$$U^{-1}(-\Delta_{\delta', \beta})U \leq -\Delta_{\delta, \alpha},$$

and hence the assertions in Corollary 3.7 hold.

The following example shows that Corollary 3.9 is sharp.

Example 3.10. *Consider the Lipschitz partition $\mathcal{P} = \{\mathbb{R}_+^2, \mathbb{R}_-^2\}$ of \mathbb{R}^2 in the upper and lower half plane with boundary $\Sigma = \mathbb{R}$. For constants $\alpha, \beta > 0$ the spectra of the operators $-\Delta_{\delta, \alpha}$ and $-\Delta_{\delta', \beta}$ can be computed via separation of variables; they are given by*

$$\sigma(-\Delta_{\delta, \alpha}) = \sigma_{\text{ess}}(-\Delta_{\delta, \alpha}) = [-\alpha^2/4, \infty)$$

and

$$\sigma(-\Delta_{\delta', \beta}) = \sigma_{\text{ess}}(-\Delta_{\delta', \beta}) = [-4/\beta^2, \infty),$$

respectively. Hence if $\beta > 4/\alpha$ then

$$\min \sigma_{\text{ess}}(-\Delta_{\delta', \beta}) = -\frac{4}{\beta^2} > -\frac{\alpha^2}{4} = \min \sigma_{\text{ess}}(-\Delta_{\delta, \alpha})$$

and it follows from Corollary 3.7 (ii) that there exists no unitary operator U in $L^2(\mathbb{R}^2)$ for which the operator inequality $U^{-1}(-\Delta_{\delta', \beta})U \leq -\Delta_{\delta, \alpha}$ holds.

Another situation which is worth to mention is the case of a Lipschitz partition of \mathbb{R}^2 which consists of a bounded domain and its complement, so that the chromatic number χ is again 2.

Example 3.11. *Consider the partition $\mathcal{P} = \{\Omega, \mathbb{R}^2 \setminus \bar{\Omega}\}$, where $\Omega \subset \mathbb{R}^2$ is a bounded domain with smooth boundary Σ , and let $\alpha, \beta > 0$ be constant. In this case*

$$\sigma_{\text{ess}}(-\Delta_{\delta, \alpha}) = \sigma_{\text{ess}}(-\Delta_{\delta', \beta}) = [0, \infty)$$

and $N(-\Delta_{\delta,\alpha}) \rightarrow +\infty$ as $\alpha \rightarrow +\infty$ according to [EY02, Theorem 1]. On the other hand we have $N(-\Delta_{\delta',\beta}) < \infty$ for any fixed $\beta > 0$ by [BLL13, Theorem 3.14 (ii)]. Hence it follows from Corollary 3.7 (iii) and Theorem 3.6 that for $\beta > 0$ there exists a sufficiently large $\alpha > 0$ such that the inequality $U^{-1}(-\Delta_{\delta',\beta})U \leq -\Delta_{\delta,\alpha}$ fails for any unitary operator of $L^2(\mathbb{R}^2)$.

In the next example, which forms a separate subsection, we discuss a particular situation with chromatic number $\chi = 3$.

3.4. An example: A symmetric star graph with three leads in \mathbb{R}^2 . We consider a symmetric star graph in \mathbb{R}^2 with three leads such that any two leads form an angle of degree $2\pi/3$, see Figure 3.1.




FIGURE 3.1. The star-graph $\Sigma = \Sigma_{12} \cup \Sigma_{23} \cup \Sigma_{13}$ separates the Euclidean space \mathbb{R}^2 into three congruent domains Ω_1 , Ω_2 and Ω_3 with bisector leads Σ_1 , Σ_2 and Σ_3 , respectively.

Let in the following $\alpha, \beta > 0$ be real constants, and let $-\Delta_{\delta,\alpha}$ and $-\Delta_{\delta',\beta}$ be the corresponding self-adjoint operators with δ and δ' -interactions, respectively, supported on the star graph. Then we have

$$(3.10) \quad \min \sigma(-\Delta_{\delta,\alpha}) = -\frac{\alpha^2}{3}$$

and

$$(3.11) \quad \min \sigma(-\Delta_{\delta',\beta}) \geq -\left(\frac{12\sqrt{3}-2}{9}\right)^2 \frac{1}{\beta^2}.$$

Whereas (3.10) is essentially a consequence of [LP08, Lemma 2.6] (and can be viewed as a strengthening of [BEW09, Theorem 3.2] in the present situation) the proof of (3.11) is of more computational nature. Both proofs are outsourced in the appendix.

Clearly the chromatic number of the partition of \mathbb{R}^2 corresponding to the star graph in Figure 3.1 is $\chi = 3$ and hence the operator inequality

$$U^{-1}(-\Delta_{\delta',\beta})U \leq -\Delta_{\delta,\alpha}$$

for the corresponding Laplacians in Theorem 3.6 is valid under the condition

$$(3.12) \quad 0 \leq \beta \leq \frac{4}{\alpha} \sin^2(\pi/3) = \frac{3}{\alpha}.$$

We point out that the assumption (3.12) can not be replaced by the weaker assumption

$$0 \leq \beta \leq \frac{4}{\alpha} \sin^2(\pi/2) = \frac{4}{\alpha},$$

which corresponds to the case $\chi = 2$ in Theorem 3.6. In fact, for $c^* := 4 - \frac{2\sqrt{3}}{9}$ we have $3 < c^* < 4$, and if we choose $\alpha, \beta > 0$ such that $\beta > c^* \frac{1}{\alpha}$ then we conclude $\min \sigma(-\Delta_{\delta, \alpha}) < \min \sigma(-\Delta_{\delta', \beta})$ from (3.10) and (3.11). This yields the following corollary.

Corollary 3.12. *Let $\alpha, \beta > 0$ and $\beta > c^* \frac{1}{\alpha}$, where $c^* = 4 - \frac{2\sqrt{3}}{9}$. Then there exists no unitary operator U in $L^2(\mathbb{R}^2)$ such that*

$$U^{-1}(-\Delta_{\delta', \beta})U \leq -\Delta_{\delta, \alpha}.$$

4. ESSENTIAL SPECTRA AND BOUND STATES OF SCHRÖDINGER OPERATORS WITH δ AND δ' -INTERACTIONS

In this section we discuss some spectral properties of the Schrödinger operators $-\Delta_{\delta, \alpha}$ and $-\Delta_{\delta', \beta}$, where the δ and δ' -interaction, respectively, is supported on certain Lipschitz partitions of \mathbb{R}^d . We are mainly interested in the following two situations: Lipschitz partitions with compact boundaries in Section 4.1 and Lipschitz partitions which are deformed on a compact subset of \mathbb{R}^d in Section 4.2. Special attention is paid to bound states in the cases $d = 2$ and $d = 3$ in Section 4.3.

4.1. Lipschitz partitions with compact boundary. Throughout this subsection we assume that the following hypothesis is satisfied.

Hypothesis 4.1. *Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a Lipschitz partition of \mathbb{R}^d , $d \geq 2$, such that the boundary $\Sigma = \cup_{k=1}^n \partial\Omega_k$ is compact.*

By Hypothesis 4.1 the partition $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ consists of $(n-1)$ bounded domains and one unbounded domain; cf. Figure 4.1. We shall call these type of partitions sometimes *compact Lipschitz partitions*.




FIGURE 4.1. Examples of compact Lipschitz partitions with chromatic numbers 3 and 4.

In the next theorem we show that under Hypothesis 4.1 the operators $-\Delta_{\delta,\alpha}$ and $-\Delta_{\delta',\beta}$ are compact perturbations of the free Laplacian $-\Delta_{\text{free}}$ defined on $H^2(\mathbb{R}^d)$. A variant of Theorem 4.2 (i) is also contained in [H89, Theorem 4] and in [BEKS94, Theorem 3.1]; cf. [BEL13] for a detailed proof in the present situation. We also mention that for a compact partition consisting of C^∞ -smooth domains it can be shown that the resolvent differences below belong to certain Schatten-von Neumann ideals depending on the space dimension d . We refer the reader to [BLL13] for more details.

Theorem 4.2. *Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a compact Lipschitz partition of \mathbb{R}^d with boundary Σ as in Hypothesis 4.1, let $\alpha, \beta : \Sigma \rightarrow \mathbb{R}$ be such that $\alpha, \beta^{-1} \in L^\infty(\Sigma)$, and let $-\Delta_{\delta,\alpha}$ and $-\Delta_{\delta',\beta}$ be the self-adjoint operators associated with \mathcal{P} . Then the following statements hold.*

- (i) *For all $\lambda \in \rho(-\Delta_{\text{free}}) \cap \rho(-\Delta_{\delta,\alpha})$ the resolvent difference*

$$(-\Delta_{\text{free}} - \lambda)^{-1} - (-\Delta_{\delta,\alpha} - \lambda)^{-1}$$

is a compact operator in $L^2(\mathbb{R}^d)$.

- (ii) *For all $\lambda \in \rho(-\Delta_{\text{free}}) \cap \rho(-\Delta_{\delta',\beta})$ the resolvent difference*

$$(-\Delta_{\text{free}} - \lambda)^{-1} - (-\Delta_{\delta',\beta} - \lambda)^{-1}$$

is a compact operator in $L^2(\mathbb{R}^d)$.

In particular, $\sigma_{\text{ess}}(-\Delta_{\delta,\alpha}) = \sigma_{\text{ess}}(-\Delta_{\delta',\beta}) = [0, \infty)$.

Proof. We shall only prove item (ii). The proof of item (i) is along the same lines and can also be found in the note [BEL13]. Let us fix $\lambda_0 < \min \sigma(-\Delta_{\delta',\beta})$ and set

$$W := (-\Delta_{\text{free}} - \lambda_0)^{-1} - (-\Delta_{\delta',\beta} - \lambda_0)^{-1}.$$

For $f, g \in L^2(\mathbb{R}^d)$ we define the functions

$$u := (-\Delta_{\text{free}} - \lambda_0)^{-1}f \quad \text{and} \quad v := (-\Delta_{\delta',\beta} - \lambda_0)^{-1}g.$$

Then we compute

$$\begin{aligned}
(Wf, g)_{L^2(\mathbb{R}^d)} &= ((-\Delta_{\text{free}} - \lambda_0)^{-1}f, g)_{L^2(\mathbb{R}^d)} - (f, (-\Delta_{\delta', \beta} - \lambda_0)^{-1}g)_{L^2(\mathbb{R}^d)} \\
(4.1) \qquad \qquad &= (u, (-\Delta_{\delta', \beta} - \lambda_0)v)_{L^2(\mathbb{R}^d)} - ((-\Delta_{\text{free}} - \lambda_0)u, v)_{L^2(\mathbb{R}^d)} \\
&= (u, -\Delta_{\delta', \beta}v)_{L^2(\mathbb{R}^d)} - (-\Delta_{\text{free}}u, v)_{L^2(\mathbb{R}^d)}.
\end{aligned}$$

Observe that $u \in H^2(\mathbb{R}^d) \subset \text{dom } \mathfrak{a}_{\delta', \beta}$ and that for any common boundary Σ_{kl} with $k, l = 1, 2, \dots, n$ and $k \neq l$ the condition $u_k|_{\Sigma_{kl}} = u_l|_{\Sigma_{kl}}$ holds. Hence we have

$$(4.2) \qquad (u, -\Delta_{\delta', \beta}v)_{L^2(\mathbb{R}^d)} = \sum_{k=1}^n (\nabla u_k, \nabla v_k)_{L^2(\Omega_k; \mathbb{C}^d)},$$

where we used the definition of $\mathfrak{a}_{\delta', \beta}$ from (3.3). Furthermore, we obtain with the help of Green's first identity (see e.g. [McL, Lemma 4.1])

$$(4.3) \qquad (-\Delta_{\text{free}}u, v)_{L^2(\mathbb{R}^d)} = \sum_{k=1}^n (\nabla u_k, \nabla v_k)_{L^2(\Omega_k; \mathbb{C}^d)} - \sum_{k=1}^n (\partial_{\nu_k} u_k|_{\partial\Omega_k}, v_k|_{\partial\Omega_k})_{L^2(\partial\Omega_k)};$$

here we also used that the restrictions u_k, v_k satisfy $u_k \in H^2(\Omega_k), v_k \in H^1(\Omega_k)$ and, hence, $\partial_{\nu_k} u_k|_{\partial\Omega_k}, v_k|_{\partial\Omega_k} \in H^{1/2}(\partial\Omega_k) \subset L^2(\partial\Omega_k)$. Combining (4.1) with (4.2) and (4.3) we obtain

$$(Wf, g)_{L^2(\mathbb{R}^d)} = \sum_{k=1}^n (\partial_{\nu_k} u_k|_{\partial\Omega_k}, v_k|_{\partial\Omega_k})_{L^2(\partial\Omega_k)}.$$

Let $\mathcal{G} := \bigoplus_{k=1}^n L^2(\partial\Omega_k)$ and $\mathcal{G}^{1/2} := \bigoplus_{k=1}^n H^{1/2}(\partial\Omega_k)$, and define the operators $T_1, T_2: L^2(\mathbb{R}^d) \rightarrow \mathcal{G}$ by

$$\begin{aligned}
T_1 f &:= \bigoplus_{k=1}^n \partial_{\nu_k} [(-\Delta_{\text{free}} - \lambda_0)^{-1}f]_k|_{\partial\Omega_k} = \bigoplus_{k=1}^n \partial_{\nu_k} u_k|_{\partial\Omega_k}, \\
T_2 g &:= \bigoplus_{k=1}^n [(-\Delta_{\delta', \beta} - \lambda_0)^{-1}g]_k|_{\partial\Omega_k} = \bigoplus_{k=1}^n v_k|_{\partial\Omega_k}.
\end{aligned}$$

As $(-\Delta_{\text{free}} - \lambda_0)^{-1}$ is continuous from $L^2(\mathbb{R}^d)$ into $H^2(\mathbb{R}^d)$ and $(-\Delta_{\delta', \beta} - \lambda_0)^{-1}$ is continuous from $L^2(\mathbb{R}^d)$ into $\text{dom } \mathfrak{a}_{\delta', \beta}$ it follows from the continuity of the trace maps that both operators T_1 and T_2 are continuous from $L^2(\mathbb{R}^d)$ into $\mathcal{G}^{1/2}$; cf. [McL, Theorem 3.37]. Since $\mathcal{G}^{1/2}$ is compactly embedded in \mathcal{G} both operators $T_1, T_2: L^2(\mathbb{R}^d) \rightarrow \mathcal{G}$ are compact. From $(Wf, g)_{L^2(\mathbb{R}^d)} = (T_1 f, T_2 g)_{\mathcal{G}}$ we conclude that

$$T_2 T_1^* = W = (-\Delta_{\text{free}} - \lambda_0)^{-1} - (-\Delta_{\delta', \beta} - \lambda_0)^{-1}$$

is a compact operator in $L^2(\mathbb{R}^d)$. Now a standard argument shows that the resolvent difference is compact for all $\lambda \in \rho(-\Delta_{\text{free}}) \cap \rho(-\Delta_{\delta', \beta})$, see, e.g., [BLL12a, Lemma 2.2].

Finally, note that $\sigma(-\Delta_{\text{free}}) = \sigma_{\text{ess}}(-\Delta_{\text{free}}) = [0, \infty)$ and hence the assertion on the essential spectra of $-\Delta_{\delta, \alpha}$ and $-\Delta_{\delta', \beta}$ follows from the compactness of the resolvent differences in (i) and (ii). \square

The next statement on the negative eigenvalues of $-\Delta_{\delta, \alpha}$ and $-\Delta_{\delta', \beta}$ is an immediate consequence of Corollary 3.7 and the fact that the essential spectra of $-\Delta_{\delta, \alpha}$ and $-\Delta_{\delta', \beta}$ coincide.

Corollary 4.3. *Let the assumptions be as in Theorem 4.2 and assume, in addition, that*

$$0 < \beta \leq \frac{4}{\alpha} \sin^2(\pi/\chi),$$

where χ is the chromatic number of the partition \mathcal{P} . Let $\{\lambda_k(-\Delta_{\delta,\alpha})\}_{k=1}^\infty$ and $\{\lambda_k(-\Delta_{\delta',\beta})\}_{k=1}^\infty$ be the negative eigenvalues of $-\Delta_{\delta,\alpha}$ and $-\Delta_{\delta',\beta}$, respectively, and let $N(-\Delta_{\delta,\alpha})$ and $N(-\Delta_{\delta',\beta})$ be their total multiplicities as in Definition 2.2. Then the following statements hold:

- (i) $\lambda_k(-\Delta_{\delta',\beta}) \leq \lambda_k(-\Delta_{\delta,\alpha})$ for all $k \in \mathbb{N}$;
- (ii) $N(-\Delta_{\delta,\alpha}) \leq N(-\Delta_{\delta',\beta})$.

Finally we show that the Schrödinger operator with a δ' -interaction of strength $\beta > 0$ has at least one negative eigenvalue.

Theorem 4.4. *Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a compact Lipschitz partition of \mathbb{R}^d with boundary Σ as in Hypothesis 4.1, let $\beta^{-1} \in L^\infty(\Sigma)$ be real, and let $-\Delta_{\delta',\beta}$ be the self-adjoint operator with δ' -interaction supported on Σ . If*

$$\int_{\partial\Omega_k} \beta^{-1}(x) d\sigma_k(x) > 0$$

holds for some bounded Ω_k , $k \in 1, \dots, n$, then $N(-\Delta_{\delta',\beta}) \geq 1$. In particular, if $\beta > 0$ is a real constant then $-\Delta_{\delta',\beta}$ has at least one negative eigenvalue.


Proof. Let $f = \chi_{\Omega_k}$ be the characteristic function of Ω_k . Then $f \in \text{dom } \mathbf{a}_{\delta',\beta}$, $\nabla f = 0$, and hence

$$\mathbf{a}_{\delta',\beta}[f] = - \int_{\partial\Omega_k} \beta^{-1}(x) d\sigma(x) < 0.$$

This implies $\inf \sigma(-\Delta_{\delta',\beta}) < 0$. □

Remark 4.5. There is no general analog of Theorem 4.4 for δ -interactions. In space dimensions $d \geq 3$ it follows implicitly from the Birman-Schwinger-type estimate in [BEKS94, Theorem 4.2 (iii)] that for $\|\alpha\|_\infty$ sufficiently small the operator $-\Delta_{\delta,\alpha}$ has no bound states. The existence of eigenvalues depends not only on α , but also on the geometry of the support of the interaction; an example in the case $d = 3$ is discussed in [EF09]. The picture is different in space dimension $d = 2$. In the simple case of a constant strength $\alpha > 0$ along the support of the interaction at least one bound state always exists, see [ET04].

4.2. Locally deformed partitions of \mathbb{R}^d . In this section we consider non-compact partitions consisting of finitely many Lipschitz domains.



$$\mathcal{P} = \{\Omega_k\}_{k=1}^7, \chi = 4$$

$$\mathcal{P}' = \{\Omega'_k\}_{k=1}^6, \chi = 3$$

FIGURE 4.2. A non-compact Lipschitz partition $\mathcal{P} = \{\Omega_k\}_{k=1}^7$ of \mathbb{R}^2 with chromatic number $\chi = 4$ and a local deformation $\mathcal{P}' = \{\Omega'_k\}_{k=1}^6$ with chromatic number $\chi = 3$.

Let in the following $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ and $\mathcal{P}' = \{\Omega'_k\}_{k=1}^{n'}$ be Lipschitz partitions of \mathbb{R}^d with boundaries Σ and Σ' , respectively. We say that \mathcal{P} and \mathcal{P}' are *local deformations of each other* if there exists a bounded domain \mathcal{B} such that

$$(4.4) \quad \Sigma \setminus \mathcal{B} = \Sigma' \setminus \mathcal{B},$$

see Figure 4.2. In addition it will be assumed that there exist $C^{1,1}$ components in the boundary Σ (and Σ') and that \mathcal{B} can be chosen in such a way that $\partial\mathcal{B} \cap \Sigma$ is contained in these components. The following hypothesis makes this more precise.

Hypothesis 4.6. *Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ and $\mathcal{P}' = \{\Omega'_k\}_{k=1}^{n'}$ be locally deformed Lipschitz partitions and let \mathcal{B} be a bounded domain with smooth boundary $\partial\mathcal{B}$ such that (4.4) holds. Let \mathcal{B}_0 and \mathcal{B}_1 be bounded domains such that $\overline{\mathcal{B}_0} \subset \mathcal{B}$, $\overline{\mathcal{B}} \subset \mathcal{B}_1$, and assume that*

$$\Gamma := (\mathcal{B}_1 \setminus \overline{\mathcal{B}_0}) \cap \Sigma = (\mathcal{B}_1 \setminus \overline{\mathcal{B}_0}) \cap \Sigma'$$

consists of $C^{1,1}$ components of a Lipschitz dissection of Σ , or equivalently, of Σ' .

In the next theorem we prove that the essential spectra of the Schrödinger operators $-\Delta_{\delta,\alpha}$ and $-\Delta_{\delta',\beta}$ do not change under local deformations of Lipschitz partitions. Our proof is partly inspired by [B62, Theorem 6.1 in English translation], where similar arguments were used for elliptic operators with Robin and mixed boundary conditions under local deformations of the boundary and local variations of the Robin coefficient.

Theorem 4.7. *Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ and $\mathcal{P}' = \{\Omega'_k\}_{k=1}^{n'}$ be Lipschitz partitions of \mathbb{R}^d which are local deformations of each other such that Hypothesis 4.6 holds. Let $\alpha, \beta^{-1} \in L^\infty(\Sigma)$ and $\alpha', \beta'^{-1} \in L^\infty(\Sigma')$ be real and assume that*

$$\alpha|_{\Sigma \setminus \mathcal{B}_0} = \alpha'|_{\Sigma' \setminus \mathcal{B}_0}, \quad \beta|_{\Sigma \setminus \mathcal{B}_0} = \beta'|_{\Sigma' \setminus \mathcal{B}_0} \quad \text{and} \quad \alpha|_\Gamma, \beta^{-1}|_\Gamma \in C^1(\Gamma).$$

Let $-\Delta_{\delta,\alpha}$, $-\Delta_{\delta',\beta}$, and $-\Delta'_{\delta,\alpha'}$, $-\Delta'_{\delta',\beta'}$ be the Schrödinger associated with the partitions \mathcal{P} and \mathcal{P}' , respectively. Then the following statements hold.

(i) For all $\lambda \in \rho(-\Delta_{\delta,\alpha}) \cap \rho(-\Delta'_{\delta,\alpha'})$ the resolvent difference

$$(-\Delta_{\delta,\alpha} - \lambda)^{-1} - (-\Delta'_{\delta,\alpha'} - \lambda)^{-1}$$

is a compact operator in $L^2(\mathbb{R}^d)$. In particular, $\sigma_{\text{ess}}(-\Delta_{\delta,\alpha}) = \sigma_{\text{ess}}(-\Delta'_{\delta,\alpha'})$.

(ii) For all $\lambda \in \rho(-\Delta_{\delta',\beta}) \cap \rho(-\Delta'_{\delta',\beta'})$ the resolvent difference

$$(-\Delta_{\delta',\beta} - \lambda)^{-1} - (-\Delta'_{\delta',\beta'} - \lambda)^{-1}$$

is a compact operator in $L^2(\mathbb{R}^d)$. In particular, $\sigma_{\text{ess}}(-\Delta_{\delta',\beta}) = \sigma_{\text{ess}}(-\Delta'_{\delta',\beta'})$.

Proof. The proof of Theorem 4.7 will be given only for the simple case that both Lipschitz partitions consist of two domains only, that is, $n = n' = 2$. The general case requires more notation but follows the same strategy. We verify (ii), the proof of (i) is similar. The fact that the essential spectra of $-\Delta_{\delta,\alpha}$ and $-\Delta'_{\delta,\alpha'}$, and $-\Delta_{\delta',\beta}$ and $-\Delta'_{\delta',\beta'}$ coincide is a direct consequence of the compactness of their resolvent differences in (i) and (ii).

Let us fix some notation; cf. Figure 4.3. Set

$$\Omega_{i1} := \Omega_i \cap \mathcal{B}, \quad \Omega_{i2} := \Omega_i \cap (\mathbb{R}^d \setminus \overline{\mathcal{B}}), \quad i = 1, 2,$$

denote the restrictions of functions f_k on Ω_k onto Ω_{kl} by f_{kl} , $k, l = 1, 2$, and let

$$\Sigma_1 := \Sigma \cap \mathcal{B}, \quad \Sigma_2 := \Sigma \cap (\mathbb{R}^d \setminus \overline{\mathcal{B}}).$$




FIGURE 4.3. The hypersurface $\partial\mathcal{B}$ splits the domain Ω_1 into the parts Ω_{11} and Ω_{12} , and the domain Ω_2 into the parts Ω_{21} and Ω_{22} . The hypersurface Σ splits into Σ_1 and Σ_2 .

We denote the restriction of β onto Σ_i by β_i , $i = 1, 2$. In the present situation the sesquilinear form $\mathfrak{a}_{\delta',\beta}$ in (3.3) is given by

$$\mathfrak{a}_{\delta',\beta}[f, g] = \sum_{k=1}^2 (\nabla f_k, \nabla g_k)_{L^2(\Omega_k; \mathbb{C}^d)} - (\beta^{-1}(f_1|_{\Sigma} - f_2|_{\Sigma}), g_1|_{\Sigma} - g_2|_{\Sigma})_{L^2(\Sigma)}$$

with $\text{dom } \mathbf{a}_{\delta',\beta} = H^1(\Omega_1) \oplus H^1(\Omega_2)$. Observe that the right hand side can also be written in the form

$$\sum_{k,l=1}^2 (\nabla f_{kl}, \nabla g_{kl})_{L^2(\Omega_{kl}; \mathbb{C}^d)} - \sum_{j=1}^2 (\beta_j^{-1}(f_{1j}|_{\Sigma_j} - f_{2j}|_{\Sigma_j}), g_{1j}|_{\Sigma_j} - g_{2j}|_{\Sigma_j})_{L^2(\Sigma_j)}.$$

Step I. We introduce an auxiliary sesquilinear form by

$$\begin{aligned} \mathbf{a}_{\delta',\beta,N}[f, g] &:= \sum_{k,l=1}^2 (\nabla f_{kl}, \nabla g_{kl})_{L^2(\Omega_{kl}; \mathbb{C}^d)} \\ &\quad - \sum_{j=1}^2 (\beta_j^{-1}(f_{1j}|_{\Sigma_j} - f_{2j}|_{\Sigma_j}), g_{1j}|_{\Sigma_j} - g_{2j}|_{\Sigma_j})_{L^2(\Sigma_j)}, \\ \text{dom } \mathbf{a}_{\delta',\beta,N} &= \bigoplus_{k,l=1}^2 H^1(\Omega_{kl}). \end{aligned}$$

As in the proof of Proposition 3.1 one verifies that $\mathbf{a}_{\delta',\beta,N}$ is a closed, densely defined form which is semibounded from below, and hence gives rise to a self-adjoint operator $-\Delta_{\delta',\beta,N}$ in $L^2(\mathbb{R}^d)$. Note that the functions in the domain of $-\Delta_{\delta',\beta,N}$ satisfy Neumann boundary conditions on $\partial\mathcal{B} \cap \Omega_i$, $i = 1, 2$, and the same δ' -type boundary conditions at Σ_i , $i = 1, 2$, as the functions in the domain of $-\Delta_{\delta',\beta}$. In this step we show that

$$(4.5) \quad (-\Delta_{\delta',\beta} - \lambda)^{-1} - (-\Delta_{\delta',\beta,N} - \lambda)^{-1}$$

is a compact operator in $L^2(\mathbb{R}^d)$ for all $\lambda \in \rho(-\Delta_{\delta',\beta}) \cap \rho(-\Delta_{\delta',\beta,N})$.

In fact, choose $\lambda_0 < \min\{\min \sigma(-\Delta_{\delta',\beta}), \min \sigma(-\Delta_{\delta',\beta,N})\}$ and let W be the resolvent difference in (4.5) with $\lambda = \lambda_0$. For $f, g \in L^2(\mathbb{R}^d)$ define

$$u := (-\Delta_{\delta',\beta} - \lambda_0)^{-1} f \quad \text{and} \quad v := (-\Delta_{\delta',\beta,N} - \lambda_0)^{-1} g.$$

A straightforward computation as in (4.1) yields

$$(4.6) \quad (Wf, g)_{L^2(\mathbb{R}^d)} = (u, -\Delta_{\delta',\beta,N} v)_{L^2(\mathbb{R}^d)} - (-\Delta_{\delta',\beta} u, v)_{L^2(\mathbb{R}^d)}.$$

As $u \in \text{dom}(-\Delta_{\delta',\beta}) \subset \text{dom } \mathbf{a}_{\delta',\beta} \subset \text{dom } \mathbf{a}_{\delta',\beta,N}$ we have for the first term on the right hand side

$$\begin{aligned} &(u, -\Delta_{\delta',\beta,N} v)_{L^2(\mathbb{R}^d)} \\ &= \sum_{k,l=1}^2 (\nabla u_{kl}, \nabla v_{kl})_{L^2(\Omega_{kl}; \mathbb{C}^d)} - \sum_{j=1}^2 (\beta_j^{-1}(u_{1j}|_{\Sigma_j} - u_{2j}|_{\Sigma_j}), v_{1j}|_{\Sigma_j} - v_{2j}|_{\Sigma_j})_{L^2(\Sigma_j)}. \end{aligned}$$

In order to rewrite the second term on the right hand side of (4.6) note first that for $u \in \text{dom}(-\Delta_{\delta',\beta})$ we have

$$\partial\nu_{j1}u_{j1}|_{\partial\mathcal{B} \cap \Omega_j} + \partial\nu_{j2}u_{j2}|_{\partial\mathcal{B} \cap \Omega_j} = 0, \quad j = 1, 2;$$

here the Neumann traces exist in $H^{1/2}(\partial\mathcal{B} \cap \Omega_j)$ due to the H^2 -regularity of the functions in $\text{dom}(-\Delta_{\delta',\beta})$ near $\partial\mathcal{B} \cap \Omega_j$ (which follows from $u_j \in H_{\text{loc}}^2(\Omega_j)$ and Lemma 3.5 (ii)). Moreover $u \in \text{dom}(-\Delta_{\delta',\beta})$ satisfies the boundary conditions

$$\partial\nu_{1j}u_{1j}|_{\Sigma_j} = \beta_j^{-1}(u_{1j}|_{\Sigma_j} - u_{2j}|_{\Sigma_j}) = -\partial\nu_{2j}u_{2j}|_{\Sigma_j}, \quad j = 1, 2,$$

by Theorem 3.3 (ii)-(c'). Hence we obtain for the second term on the right hand side of (4.6) when integrating by parts,

$$\begin{aligned} & (-\Delta_{\delta',\beta} u, v)_{L^2(\mathbb{R}^d)} \\ &= \sum_{k,l=1}^2 (\nabla u_{kl}, \nabla v_{kl})_{L^2(\Omega_{kl}; \mathbb{C}^d)} - \sum_{j=1}^2 (\beta_j^{-1} (u_{1j}|_{\Sigma_j} - u_{2j}|_{\Sigma_j}), v_{1j}|_{\Sigma_j} - v_{2j}|_{\Sigma_j})_{L^2(\Sigma_j)} \\ &\quad - \sum_{j=1}^2 (\partial_{\nu_j} u_{j1}|_{\partial\mathcal{B} \cap \Omega_j}, v_{j1}|_{\partial\mathcal{B} \cap \Omega_j} - v_{j2}|_{\partial\mathcal{B} \cap \Omega_j})_{L^2(\partial\mathcal{B} \cap \Omega_j)}. \end{aligned}$$

Thus (4.6) has the form

$$\begin{aligned} (4.7) \quad (Wf, g)_{L^2(\mathbb{R}^d)} &= \sum_{j=1}^2 (\partial_{\nu_j} u_{j1}|_{\partial\mathcal{B} \cap \Omega_j}, v_{j1}|_{\partial\mathcal{B} \cap \Omega_j} - v_{j2}|_{\partial\mathcal{B} \cap \Omega_j})_{L^2(\partial\mathcal{B} \cap \Omega_j)} \\ &= (T_1 f, T_2 g)_{L^2(\partial\mathcal{B})}, \end{aligned}$$

where the operators $T_1, T_2: L^2(\mathbb{R}^d) \rightarrow L^2(\partial\mathcal{B})$ are defined by

$$\begin{aligned} T_1 f &:= \bigoplus_{j=1}^2 \partial_{\nu_j} [(-\Delta_{\delta',\beta} - \lambda_0)^{-1} f]_{j1}|_{\partial\mathcal{B} \cap \Omega_j} = \bigoplus_{j=1}^2 \partial_{\nu_j} u_{j1}|_{\partial\mathcal{B} \cap \Omega_j} \\ T_2 g &:= \bigoplus_{j=1}^2 [(-\Delta_{\delta',\beta,N} - \lambda_0)^{-1} g]_{j1}|_{\partial\mathcal{B} \cap \Omega_j} - [(-\Delta_{\delta',\beta,N} - \lambda_0)^{-1} g]_{j2}|_{\partial\mathcal{B} \cap \Omega_j} \\ &= \bigoplus_{j=1}^2 [v_{j1}|_{\partial\mathcal{B} \cap \Omega_j} - v_{j2}|_{\partial\mathcal{B} \cap \Omega_j}]. \end{aligned}$$

Since $(-\Delta_{\delta',\beta,N} - \lambda_0)^{-1}$ is bounded from $L^2(\mathbb{R}^d)$ into $\text{dom } \mathfrak{a}_{\delta',\beta,N}$ it follows from [McL, Theorem 3.37] that the operator T_2 maps $L^2(\mathbb{R}^d)$ boundedly into

$$H^{1/2}(\partial\mathcal{B} \cap \Omega_1) \oplus H^{1/2}(\partial\mathcal{B} \cap \Omega_2),$$

which is compactly embedded in $L^2(\partial\mathcal{B} \cap \Omega_1) \oplus L^2(\partial\mathcal{B} \cap \Omega_2) = L^2(\partial\mathcal{B})$. Hence $T_2: L^2(\mathbb{R}^d) \rightarrow L^2(\partial\mathcal{B})$ is compact. We shall show below in Step II that the operator $T_1: L^2(\mathbb{R}^d) \rightarrow L^2(\partial\mathcal{B})$ is bounded, so that by (4.7)

$$T_2^* T_1 = W = (-\Delta_{\delta',\beta} - \lambda_0)^{-1} - (-\Delta_{\delta',\beta,N} - \lambda_0)^{-1}$$

is a compact operator in $L^2(\mathbb{R}^d)$. It then follows that the resolvent difference in (4.5) is compact for all $\lambda \in \rho(-\Delta_{\delta',\beta}) \cap \rho(-\Delta_{\delta',\beta,N})$, see, e.g. [BLL12a, Lemma 2.2].

Step II. We verify that $T_1: L^2(\mathbb{R}^d) \rightarrow L^2(\partial\mathcal{B})$ is bounded, which is essentially a consequence of [McL, Theorem 4.18 (ii)] and the H^2 -regularity of the functions in $\text{dom } (-\Delta_{\delta',\beta})$ near $\partial\mathcal{B} \cap \Omega_j$; cf. Lemma 3.5 (ii). More precisely, let $0 < s < t < 1$ and let \mathcal{B}_s and \mathcal{B}_t be bounded domains with smooth boundaries such that

$$\overline{\mathcal{B}}_0 \subset \mathcal{B}_s \subset \overline{\mathcal{B}}_s \subset \mathcal{B} \subset \overline{\mathcal{B}} \subset \mathcal{B}_t \subset \overline{\mathcal{B}}_t \subset \mathcal{B}_1.$$

Set $\mathcal{R}_j := (\mathcal{B}_t \setminus \overline{\mathcal{B}}_s) \cap \Omega_j$ and $\mathcal{S}_j := (\mathcal{B}_1 \setminus \overline{\mathcal{B}}_0) \cap \Omega_j$, $j = 1, 2$. Since $\Gamma = (\mathcal{B}_1 \setminus \overline{\mathcal{B}}_0) \cap \Sigma$ is $C^{1,1}$ we conclude for $u = (-\Delta_{\delta',\beta} - \lambda_0)^{-1} f$ from [McL, Theorem 4.18 (ii)] that

$$(4.8) \quad \|u_j|_{\mathcal{R}_j}\|_{H^2(\mathcal{R}_j)} \leq C_j (\|u_j|_{\mathcal{S}_j}\|_{H^1(\mathcal{S}_j)} + \|\partial_{\nu_j} u_j|_{\Gamma}\|_{H^{1/2}(\Gamma)} + \|f_j|_{\mathcal{S}_j}\|_{L^2(\mathcal{S}_j)})$$

holds for some constants C_j , $j = 1, 2$. The continuity of $(-\Delta_{\delta',\beta} - \lambda_0)^{-1}$ from $L^2(\mathbb{R}^d)$ into $\text{dom } \mathfrak{a}_{\delta',\beta}$ yields $\|u_j|_{\mathcal{S}_j}\|_{H^1(\mathcal{S}_j)} \leq C' \|f\|_{L^2(\mathbb{R}^d)}$ with some constant C' . Furthermore, the boundary conditions $\partial_{\nu_1} u_1|_{\Gamma} = \beta^{-1}(u_1|_{\Gamma} - u_2|_{\Gamma}) = -\partial_{\nu_2} u_2|_{\Gamma}$, the continuity of the trace and of $(-\Delta_{\delta',\beta} - \lambda_0)^{-1}$ from $L^2(\mathbb{R}^d)$ into $\text{dom } \mathfrak{a}_{\delta',\beta}$ yields

$$\begin{aligned} \|\partial_{\nu_j} u_j|_{\Gamma}\|_{H^{1/2}(\Gamma)} &\leq D(\|u_1|_{\Gamma}\|_{H^{1/2}(\Gamma)} + \|u_2|_{\Gamma}\|_{H^{1/2}(\Gamma)}) \\ &\leq D'(\|u_1|_{\mathcal{S}_1}\|_{H^1(\mathcal{S}_1)} + \|u_2|_{\mathcal{S}_2}\|_{H^1(\mathcal{S}_2)}) \leq D'' \|f\|_{L^2(\mathbb{R}^d)} \end{aligned}$$

with some constants D, D', D'' . If P_j denotes the orthogonal projection in $L^2(\mathbb{R}^d)$ onto $L^2(\mathcal{R}_j)$ then we conclude together with (4.8) that

$$\text{ran}(P_j(-\Delta_{\delta',\beta} - \lambda_0)^{-1}) \subset H^2(\mathcal{R}_j)$$

and that the operator $P_j(-\Delta_{\delta',\beta} - \lambda_0)^{-1}$ is bounded from $L^2(\mathbb{R}^d)$ into $H^2(\mathcal{R}_j)$ for $j = 1, 2$. Hence $f \mapsto \partial_{\nu_j} [(-\Delta_{\delta',\beta} - \lambda_0)^{-1} f]|_{\partial\mathcal{B} \cap \Omega_j}$ is bounded from $L^2(\mathbb{R}^d)$ into $H^{1/2}(\partial\mathcal{B} \cap \Omega_j)$, $j = 1, 2$, and, in particular, T_1 is bounded from $L^2(\mathbb{R}^d)$ into $L^2(\partial\mathcal{B})$.

Step III. As in Step I we introduce an auxiliary sesquilinear form by

$$\begin{aligned} \mathfrak{a}'_{\delta',\beta',N}[h, k] &:= \sum_{k,l=1}^2 (\nabla h_{kl}, \nabla k_{kl})_{L^2(\Omega'_{kl}; \mathbb{C}^d)} \\ &\quad - \sum_{j=1}^2 (\beta_j'^{-1}(h_{1j}|_{\Sigma'_j} - h_{2j}|_{\Sigma'_j}), k_{1j}|_{\Sigma'_j} - k_{2j}|_{\Sigma'_j})_{L^2(\Sigma'_j)}, \\ \text{dom } \mathfrak{a}'_{\delta',\beta',N} &= \bigoplus_{k,l=1}^2 H^1(\Omega'_{kl}), \end{aligned}$$

where $\Omega'_{i1} := \Omega'_i \cap \mathcal{B}$, $\Omega'_{i2} := \Omega'_i \cap (\mathbb{R}^d \setminus \overline{\mathcal{B}})$, $i = 1, 2$, g_{ij}, h_{ij} denote the corresponding restrictions of functions g, h , and $\Sigma'_1 := \Sigma' \cap \mathcal{B}$, $\Sigma'_2 := \Sigma' \cap (\mathbb{R}^d \setminus \overline{\mathcal{B}}) = \Sigma_2$. The form $\mathfrak{a}'_{\delta',\beta',N}$ is closed, densely defined and semibounded from below, and hence gives rise to a self-adjoint operator $-\Delta'_{\delta',\beta',N}$ in $L^2(\mathbb{R}^d)$. In the same way as in Step I and II one verifies that

$$(4.9) \quad (-\Delta'_{\delta',\beta'} - \lambda)^{-1} - (-\Delta'_{\delta',\beta',N} - \lambda)^{-1}$$

is compact for all $\lambda \in (-\Delta'_{\delta',\beta'}) \cap (-\Delta'_{\delta',\beta',N})$.

Step IV. Since the Lipschitz partitions \mathcal{P} and \mathcal{P}' are local deformations of each other and Hypothesis 4.6 holds the self-adjoint operators $-\Delta_{\delta',\beta,N}$ and $-\Delta'_{\delta',\beta',N}$ from Steps I-III admit the direct sum decompositions

$$-\Delta_{\delta',\beta,N} = H_1 \oplus H_2 \quad \text{and} \quad -\Delta'_{\delta',\beta',N} = H_1 \oplus H'_2$$

with respect to the decomposition $L^2(\mathbb{R}^d) = L^2(\mathbb{R}^d \setminus \overline{\mathcal{B}}) \oplus L^2(\mathcal{B})$. The operators H_2 and H'_2 acting in $L^2(\mathcal{B})$ have compact resolvents in view of the compact embeddings of the spaces $H^1(\mathcal{B} \cap \Omega_1) \oplus H^1(\mathcal{B} \cap \Omega_2)$ and $H^1(\mathcal{B} \cap \Omega'_1) \oplus H^1(\mathcal{B} \cap \Omega'_2)$ into $L^2(\mathcal{B})$. This implies the compactness of

$$(-\Delta_{\delta',\beta,N} - \lambda)^{-1} - (-\Delta'_{\delta',\beta',N} - \lambda)^{-1}$$

for all $\lambda \in \rho(-\Delta_{\delta',\beta,N}) \cap \rho(-\Delta'_{\delta',\beta',N})$ and hence assertion (ii) follows together with the compactness of the resolvent differences in (4.5) and (4.9). \square

The following corollary is an immediate consequence of Theorem 4.7 and the fact that for the Lipschitz partition $\mathcal{P}' = \{\mathbb{R}_+^d, \mathbb{R}_-^d\}$ and constants $\alpha, \beta > 0$ the essential spectra of $-\Delta'_{\delta, \alpha}$ and $-\Delta'_{\delta', \beta}$ can be computed by separation of variables:

$$\begin{aligned}\sigma(-\Delta'_{\delta, \alpha}) &= \sigma_{\text{ess}}(-\Delta'_{\delta, \alpha}) = [-\alpha^2/4, \infty), \\ \sigma(-\Delta'_{\delta', \beta}) &= \sigma_{\text{ess}}(-\Delta'_{\delta', \beta}) = [-4/\beta^2, \infty).\end{aligned}$$

Corollary 4.8. *Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a local deformation of the Lipschitz partition $\mathcal{P}' = \{\mathbb{R}_+^d, \mathbb{R}_-^d\}$ of \mathbb{R}^d and let $\alpha, \beta > 0$ be constant. Then the essential spectra of $-\Delta_{\delta, \alpha}$ and $-\Delta_{\delta', \beta}$ are given by*

$$\sigma_{\text{ess}}(-\Delta_{\delta, \alpha}) = [-\alpha^2/4, \infty) \quad \text{and} \quad \sigma_{\text{ess}}(-\Delta_{\delta', \beta}) = [-4/\beta^2, \infty).$$

The next corollary is a consequence of Theorem 2.3, Corollary 3.7 and Corollary 4.8.

Corollary 4.9. *Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a local deformation of the Lipschitz partition $\mathcal{P}' = \{\mathbb{R}_+^d, \mathbb{R}_-^d\}$, assume that the chromatic number of \mathcal{P} is $\chi = 2$ and that the constants $\alpha, \beta > 0$ satisfy*

$$\beta = \frac{4}{\alpha}, \quad \text{and hence} \quad \sigma_{\text{ess}}(-\Delta_{\delta, \alpha}) = \sigma_{\text{ess}}(-\Delta_{\delta', \beta}) = [-\alpha^2/4, \infty).$$

Let $\{\lambda_k(-\Delta_{\delta, \alpha})\}_{k=1}^\infty$ and $\{\lambda_k(-\Delta_{\delta', \beta})\}_{k=1}^\infty$ be the eigenvalues of $-\Delta_{\delta, \alpha}$ and $-\Delta_{\delta', \beta}$ below $-\alpha^2/4$, respectively, and let $N(-\Delta_{\delta, \alpha})$ and $N(-\Delta_{\delta', \beta})$ be their total multiplicities as in Definition 2.2. Then the following statements hold:

- (i) $\lambda_k(-\Delta_{\delta', \beta}) \leq \lambda_k(-\Delta_{\delta, \alpha})$ for all $k \in \mathbb{N}$;
- (ii) $N(-\Delta_{\delta, \alpha}) \leq N(-\Delta_{\delta', \beta})$.

4.3. Locally deformed partitions of \mathbb{R}^2 and \mathbb{R}^3 . In this subsection special attention is paid to bound states of $-\Delta_{\delta, \alpha}$ and $-\Delta_{\delta', \beta}$ induced by local deformations of certain Lipschitz partitions of \mathbb{R}^2 and \mathbb{R}^3 . We first characterize the essential spectra of $-\Delta_{\delta, \alpha}$ and $-\Delta_{\delta', \beta}$ associated with partitions, which are local deformations of a partition $\{\Omega, \mathbb{R}^2 \setminus \Omega\}$ with Ω being a wedge, see Figure 4.4.




FIGURE 4.4. A wedge $\Omega \subset \mathbb{R}^2$ with angle $\varphi \in (0, \pi]$ and boundary consisting of the two rays Σ_1 and Σ_2 ; the axis x_1 coincides with the ray Σ_1 .

Theorem 4.10. *Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a local deformation of the Lipschitz partition $\mathcal{P}' = \{\Omega, \mathbb{R}^2 \setminus \Omega\}$ of \mathbb{R}^2 , where Ω is a wedge in \mathbb{R}^2 and let $\alpha, \beta > 0$ be constant. Then the essential spectra of $-\Delta_{\delta, \alpha}$ and $-\Delta_{\delta', \beta}$ are given by*

$$\sigma_{\text{ess}}(-\Delta_{\delta, \alpha}) = [-\alpha^2/4, \infty) \quad \text{and} \quad \sigma_{\text{ess}}(-\Delta_{\delta', \beta}) = [-4/\beta^2, \infty).$$

Proof. According to Theorem 4.7 it suffices to show the statements for the operators $-\Delta'_{\delta,\alpha}$ and $-\Delta'_{\delta',\beta}$ associated with the Lipschitz partition $\mathcal{P}' = \{\Omega, \mathbb{R}^2 \setminus \overline{\Omega}\}$. In fact, the assertion for $-\Delta'_{\delta,\alpha}$ can be found in [EN03, Proposition 5.4], and hence we verify $\sigma_{\text{ess}}(-\Delta'_{\delta',\beta}) = [-4/\beta^2, \infty)$ only.

Step I. Decompose \mathbb{R}^2 into eight domains as in Figure 4.5, where $\Omega_1, \Omega'_1, \Omega_2, \Omega'_2$ coincide (up to rotations and translations) with $[0, l] \times \mathbb{R}_+$ for some $l > 0$; Ω_3, Ω'_3 are bounded Lipschitz domains, and Ω_4 and Ω_5 are wedges with angles φ and $2\pi - \varphi$, respectively. We choose this partition in such a way that Ω coincides with $\Omega_1 \cup \Omega_2 \cup \Omega_3 \cup \Omega_4$ up to a set of Lebesgue measure zero.




FIGURE 4.5. A partition of \mathbb{R}^2 into eight domains. The wedge Ω coincides with $\Omega_1 \cup \Omega_2 \cup \Omega_3 \cup \Omega_4$ up to a set of Lebesgue measure zero.

Let \mathcal{P} be the corresponding partition and set $\Sigma_k := \overline{\Omega_k} \cap \overline{\Omega'_k}$ for $k = 1, 2, 3$. Clearly, $\partial\Omega = \Sigma_1 \cup \Sigma_2 \cup \Sigma_3$. Observe that such a decomposition can be constructed for any $l > 0$. We use the notation $f_\Omega := f|_\Omega$. Consider the quadratic form

$$\mathfrak{a}'_{\delta',\beta,N}[f] := \sum_{\Omega \in \mathcal{P}} \|\nabla f_\Omega\|_{L^2(\Omega; \mathbb{C}^d)}^2 - \sum_{k=1}^3 \beta^{-1} \|f_{\Omega_k}|_{\Sigma_k} - f_{\Omega'_k}|_{\Sigma_k}\|_{L^2(\Sigma_k)}^2,$$

$$\text{dom } \mathfrak{a}'_{\delta',\beta,N} := \oplus_{\Omega \in \mathcal{P}} H^1(\Omega)$$

Similarly as in the proof of Proposition 3.1 one verifies that the form $\mathfrak{a}'_{\delta',\beta,N}$ is closed, densely defined, symmetric and semibounded from below. The corresponding self-adjoint operator $-\Delta'_{\delta',\beta,N}$ can be decomposed into the orthogonal sum of five self-adjoint operators

$$(4.10) \quad -\Delta'_{\delta',\beta,N} = \bigoplus_{k=1}^5 H_k,$$

where H_i acts in $L^2(\Omega_i) \oplus L^2(\Omega'_i)$, $i = 1, 2, 3$, and H_4 and H_5 are the self-adjoint Neumann Laplacians on the wedges Ω_4 and Ω_5 in $L^2(\Omega_4)$ and $L^2(\Omega_5)$, respectively. Hence we have

$$(4.11) \quad \sigma_{\text{ess}}(H_4) = \sigma_{\text{ess}}(H_5) = [0, +\infty).$$

The operator H_3 acts on a bounded domain and in view of the compact embedding of the space $H^1(\Omega_3) \oplus H^1(\Omega'_3)$ into $L^2(\Omega_3) \oplus L^2(\Omega'_3)$ we obtain

$$(4.12) \quad \sigma_{\text{ess}}(H_3) = \emptyset.$$

Separation of variables shows that the essential spectra of the operators H_1 and H_2 have the form

$$(4.13) \quad \sigma_{\text{ess}}(H_1) = \sigma_{\text{ess}}(H_2) = [\varepsilon(\beta, l), +\infty),$$

where $\varepsilon(\beta, l)$ is the principal eigenvalue of the self-adjoint one-dimensional Schrödinger operator on the interval $(-l, l)$ with Neumann boundary conditions at the endpoints $-l$ and l , and a δ' -interaction of strength β at the origin. According to [EJ13, Lemma 3.3]

$$(4.14) \quad \varepsilon(\beta, l) < -\frac{4}{\beta^2} \quad \text{and} \quad \lim_{l \rightarrow +\infty} \varepsilon(\beta, l) = -\frac{4}{\beta^2}.$$

From the decomposition (4.10) and the characterizations (4.11), (4.12), (4.13) we conclude

$$\sigma_{\text{ess}}(-\Delta'_{\delta', \beta, \mathbb{N}}) = [\varepsilon(\beta, l), +\infty).$$

Clearly, $\mathbf{a}'_{\delta', \beta, \mathbb{N}} \leq \mathbf{a}'_{\delta', \beta}$ holds in the sense of Definition 2.1 and hence

$$\min \sigma_{\text{ess}}(-\Delta'_{\delta', \beta}) \geq \varepsilon(\beta, l)$$

by Theorem 2.3 (ii). As we noted above, the construction in the proof can be realized for any $l > 0$. Thus by (4.14)

$$\min \sigma_{\text{ess}}(-\Delta'_{\delta', \beta}) \geq -\frac{4}{\beta^2}.$$

Step II. In view of Step I it suffices to show that for any $\lambda \in [-4/\beta^2, +\infty)$ there exists a singular sequence for the operator $-\Delta'_{\delta', \beta}$ corresponding to λ . Let us fix the axes (x_1, x_2) such that the axis x_1 coincides with the side Σ_1 of the wedge Ω , see Figure 4.4. Let us fix two functions $\varphi_1, \varphi_2 \in C_0^\infty([0, \infty))$ with $\text{supp } \varphi_1$ and $\text{supp } \varphi_2$ in $[0, 2)$ such that $\varphi_1(x) = \varphi_2(x) = 1$ in the vicinity of the point $x = 0$ and $0 \leq \varphi_2(x) \leq 1$. Consider the sequence of functions

$$\psi_{n,p}(x) := \frac{1}{\sqrt{n}} \varphi_1\left(\frac{1}{n}|x_1 - x_1^{(n)}|\right) \varphi_2\left(\frac{1}{n}|x_2|\right) \text{sign}(x_2) e^{-2|x_2|/\beta} e^{ipx_1}, \quad n \in \mathbb{N},$$

where $p \geq 0$ is arbitrary and the sequence $\{x_1^{(n)}\}$ tends to $+\infty$ sufficiently fast, so that the sequence of the supports $\text{supp } \psi_{n,p}$ does not intersect the ray Σ_2 of the wedge. We denote by $\psi_{n,p,\Omega}$ and $\psi_{n,p,\mathbb{R}^2 \setminus \overline{\Omega}}$ the restriction of $\psi_{n,p}$ onto Ω and $\mathbb{R}^2 \setminus \overline{\Omega}$, respectively. Computing the traces of $\psi_{n,p}$ from both sides of Σ_1 we find

$$\partial_\nu \psi_{n,p,\Omega}|_{\Sigma_1} = \frac{2}{\beta} \frac{1}{\sqrt{n}} \varphi_1\left(\frac{1}{n}|x_1 - x_1^{(n)}|\right) e^{ipx_1} = \frac{2}{\beta} \psi_{n,p,\Omega}|_{\Sigma_1} = -\frac{2}{\beta} \psi_{n,p,\mathbb{R}^2 \setminus \overline{\Omega}}|_{\Sigma_1}$$

with the normal ν pointing outwards of Ω . Thus we conclude from Theorem 3.3 (ii) that the functions $\psi_{n,p}$ are in $\text{dom}(-\Delta'_{\delta', \beta})$. Obviously, the sequence of the functions $\{\psi_{n,p}\}$ converges weakly to zero. Moreover, with the help of the dominated convergence theorem we get

$$\lim_{n \rightarrow \infty} \|\psi_{n,p}\|^2 = \|\varphi_1\|_{L^2(\mathbb{R})}^2 \int_{\mathbb{R}} e^{-4|x|/\beta} dx = \frac{\beta}{2} \|\varphi_1\|_{L^2(\mathbb{R})}^2 \neq 0.$$

One can check via direct computation that

$$-\Delta'_{\delta',\beta}\psi_{n,p} = \left(-\frac{4}{\beta^2} + p^2\right)\psi_{n,p} + O\left(\frac{1}{n}\right), \quad n \rightarrow \infty,$$

which yields

$$(4.15) \quad \|(-\Delta'_{\delta',\beta} + 4/\beta^2 - p^2)\psi_{n,p}\|_{L^2(\mathbb{R}^2)} \rightarrow 0, \quad n \rightarrow +\infty.$$

Therefore, the sequence

$$\tilde{\psi}_{n,p} := \frac{\psi_{n,p}}{\|\psi_{n,p}\|}, \quad n \in \mathbb{N},$$

is a singular sequence for the operator $-\Delta'_{\delta',\beta}$ corresponding to the point $-4/\beta^2 + p^2$. Since the choice of p is arbitrary, the claim is proven. \square

The next corollary is a consequence of Theorems 2.3 and 4.10.

Corollary 4.11. *Let $\mathcal{P} = \{\Omega_k\}_{k=1}^n$ be a local deformation of the Lipschitz partition $\mathcal{P}' = \{\Omega, \mathbb{R}^2 \setminus \overline{\Omega}\}$ with Ω being a wedge, assume that the chromatic number of \mathcal{P} is $\chi = 2$ and that the constants $\alpha, \beta > 0$ satisfy*

$$\beta = \frac{4}{\alpha}, \quad \text{and hence} \quad \sigma_{\text{ess}}(-\Delta_{\delta,\alpha}) = \sigma_{\text{ess}}(-\Delta_{\delta',\beta}) = [-\alpha^2/4, \infty).$$

Let $\{\lambda_k(-\Delta_{\delta,\alpha})\}_{k=1}^\infty$ and $\{\lambda_k(-\Delta_{\delta',\beta})\}_{k=1}^\infty$ be the eigenvalues of $-\Delta_{\delta,\alpha}$ and $-\Delta_{\delta',\beta}$ below $-\alpha^2/4$, respectively, and let $N(-\Delta_{\delta,\alpha})$ and $N(-\Delta_{\delta',\beta})$ be their total multiplicities as in Definition 2.2. Then the following statements hold:

- (i) $\lambda_k(-\Delta_{\delta',\beta}) \leq \lambda_k(-\Delta_{\delta,\alpha})$ for all $k \in \mathbb{N}$;
- (ii) $N(-\Delta_{\delta,\alpha}) \leq N(-\Delta_{\delta',\beta})$.

The following corollary shows the existence of negative bound states of $-\Delta_{\delta',\beta}$ for locally deformed broken lines in \mathbb{R}^2 . The assertion follows directly from [EI01, Theorem 5.2] and Corollary 4.11. We mention that in [EI01] more general weakly deformed curves were considered.

Corollary 4.12. *Let $\mathcal{P} = \{\Omega, \mathbb{R}^2 \setminus \overline{\Omega}\}$ be a local deformation of the Lipschitz partition $\mathcal{P}' = \{\Omega', \mathbb{R}^2 \setminus \overline{\Omega'}\}$, where Ω' is a wedge with angle $\varphi \in (0, \pi]$. In the case $\varphi = \pi$ let $\mathcal{P} \neq \mathcal{P}'$. Assume, in addition, that $\partial\Omega$ is piecewise C^1 -smooth. Then $N(-\Delta_{\delta',\beta}) \geq 1$ holds for any $\beta > 0$.*

In the next proposition we show the existence of bound states for δ and δ' -operators for special local deformations of the partition $\{\mathbb{R}_+^2, \mathbb{R}_-^2\}$.

Proposition 4.13. *Let $\Omega_1 \subset \mathbb{R}_+^2$ be a bounded Lipschitz domain and consider the Lipschitz partition $\mathcal{P} = \{\Omega_k\}_{k=1}^3$ of \mathbb{R}^2 , where $\Omega_2 = \mathbb{R}_+^2 \setminus \overline{\Omega_1}$ and $\Omega_3 = \mathbb{R}_-^2$ as in Figure 4.6. Let $\alpha, \beta > 0$ be constant and let the Schrödinger operators $-\Delta_{\delta,\alpha}$ and $-\Delta_{\delta',\beta}$ be associated with \mathcal{P} . Then the following statements hold.*

- (i) $\sigma_{\text{ess}}(-\Delta_{\delta,\alpha}) = [-\alpha^2/4, +\infty)$ and $N(-\Delta_{\delta,\alpha}) \geq 1$;
- (ii) $\sigma_{\text{ess}}(-\Delta_{\delta',\beta}) = [-4/\beta^2, +\infty)$ and $N(-\Delta_{\delta',\beta}) \geq 1$.




FIGURE 4.6. The partition of \mathbb{R}^2 via a straight line and a compact Lipschitz contour, which consists of a bounded domain Ω_1 and two unbounded domains Ω_2 and Ω_3 .

Proof. The characterization of the essential spectra in (i) and (ii) is a direct consequence of Corollary 4.8. We shall show the assertion $N(-\Delta_{\delta,\alpha}) \geq 1$ in (i) first. For this we can assume that Σ_{23} is the hyperplane defined by $x_2 = 0$, where $x = (x_1, x_2) \in \mathbb{R}^2$. Let $\varphi \in C_0^\infty(\mathbb{R})$ be equal to one in the neighbourhood of the origin and consider the sequence of functions

$$f_n(x) := \varphi\left(\frac{1}{n}x_1\right)e^{-(\alpha/2)|x_2|} \in H^1(\mathbb{R}^2), \quad n \in \mathbb{N},$$

and the sequence of real values

$$I_n := \mathbf{a}_{\delta,\alpha}[f_n] + \frac{\alpha^2}{4}\|f_n\|_{L^2(\mathbb{R}^2)}^2, \quad n \in \mathbb{N}.$$

From the definition of the form $\mathbf{a}_{\delta,\alpha}$ in (3.2),

$$\begin{aligned} \|f_n\|_{L^2(\mathbb{R}^2)}^2 &= \frac{2n}{\alpha}\|\varphi\|_{L^2(\mathbb{R})}^2, \\ \|\nabla f_n\|_{L^2(\mathbb{R}^2;\mathbb{C}^2)}^2 &= \frac{2}{n\alpha}\|\varphi'\|_{L^2(\mathbb{R})}^2 + \frac{\alpha n}{2}\|\varphi\|_{L^2(\mathbb{R})}^2, \end{aligned}$$

and $\|f_n|_{\Sigma_{23}}\|_{L^2(\Sigma_{23})}^2 = n\|\varphi\|_{L^2(\mathbb{R})}^2$ we obtain

$$\begin{aligned} I_n &= \|\nabla f_n\|_{L^2(\mathbb{R}^2;\mathbb{C}^2)}^2 - \alpha\|f_n|_{\Sigma_{12}}\|_{L^2(\Sigma_{12})}^2 - \alpha\|f_n|_{\Sigma_{23}}\|_{L^2(\Sigma_{23})}^2 + \frac{\alpha^2}{4}\|f_n\|_{L^2(\mathbb{R}^2)}^2 \\ &= \frac{2}{\alpha n}\|\varphi'\|_{L^2(\mathbb{R})}^2 - \alpha\|f_n|_{\Sigma_{12}}\|_{L^2(\Sigma_{12})}^2. \end{aligned}$$

Let $D > 0$ be the distance from Σ_{23} to the farthest point of Σ_{12} . For large $n \in \mathbb{N}$

$$I_n \leq \frac{2}{\alpha n}\|\varphi'\|_{L^2(\mathbb{R})}^2 - \alpha e^{-\alpha D}|\Sigma_{12}|,$$

and hence for sufficiently large $n \in \mathbb{N}$ we obtain that $I_n < 0$, which proves the existence of at least one bound state for the operator $-\Delta_{\delta,\alpha}$.

In order to show that $-\Delta_{\delta',\beta}$ has at least one bound state we note that $-\Delta_{\delta,4\beta-1}$ has at least one bound state by the considerations above. Hence Corollary 4.11 implies $N(-\Delta_{\delta',\beta}) \geq 1$. \square

The next result shows the existence of negative bound states of $-\Delta_{\delta',\beta}$ for certain hypersurfaces in \mathbb{R}^3 . The assertion follows directly from and [EK03, Theorem 4.3]

and Corollary 4.9. We mention that in [EK03] more general weakly deformed planes were considered.

Corollary 4.14. *Let $\mathcal{P} = \{\Omega, \mathbb{R}^3 \setminus \bar{\Omega}\}$ be a local deformation of the Lipschitz partition $\mathcal{P}' = \{\mathbb{R}_+^3, \mathbb{R}_-^3\}$ such that $\mathcal{P} \neq \mathcal{P}'$. If, in addition, $\partial\Omega$ is C^4 -smooth and admits a global natural parametrization in the sense of [dC, §2-3, Definition 2] then $N(-\Delta_{\delta',\beta}) \geq 1$ for all sufficiently small $\beta > 0$.*

5. APPENDIX: SOBOLEV SPACES ON WEDGES AND A SYMMETRIC STAR GRAPH WITH THREE LEADS IN \mathbb{R}^2

In this appendix we verify the statements

$$\min(-\Delta_{\delta,\alpha}) = -\frac{\alpha^2}{3} \quad \text{and} \quad \min(-\Delta_{\delta',\beta}) \geq -\left(\frac{12\sqrt{3}-2}{9}\right)^2 \frac{1}{\beta^2}$$

from (3.10) and (3.11), where $\alpha, \beta > 0$ are real constants and the δ and δ' -interaction is supported on the symmetric star graph with three leads in Figure 3.1. We first provide some useful estimates and decompositions for H^1 -functions on wedges.

5.1. Sobolev spaces on wedges in \mathbb{R}^2 . Let Ω be a wedge with angle $\varphi \in (0, \pi]$ as in the figure below. The estimates for functions $f \in H^1(\Omega)$ in Lemma 5.1 and Lemma 5.2 below will be used in the proofs of (3.10) and (3.11).




FIGURE 5.1. A wedge $\Omega \subset \mathbb{R}^2$ with angle $\varphi \in (0, \pi]$ and boundary consisting of the two rays Σ_1 and Σ_2 ; the ray Σ separates Ω into two wedges Ω_1 and Ω_2 .

The first lemma is a reformulation of [LP08, Lemma 2.6].

Lemma 5.1. *Let $\Omega \subset \mathbb{R}^2$ be a wedge with angle $\varphi \in (0, \pi]$ and boundary $\partial\Omega$. Then for every $f \in H^1(\Omega)$ and all $\gamma > 0$ the estimate*

$$\|\nabla f\|_{L^2(\Omega; \mathbb{C}^2)}^2 - \gamma \|f|_{\partial\Omega}\|_{L^2(\partial\Omega)}^2 \geq -\frac{\gamma^2}{\sin^2(\varphi/2)} \|f\|_{L^2(\Omega)}^2$$

holds. For $\varphi \in (0, \pi)$ the estimate is sharp.

We provide a variant of Lemma 5.1 which will be useful in the proof of (3.11). We note for completeness that the estimate below is not sharp for $\varphi \in (0, \pi)$.

Lemma 5.2. *Let $\Omega \subset \mathbb{R}^2$ be a wedge with angle $\varphi \in (0, \pi]$ and boundary $\partial\Omega$. Let Σ be a ray separating Ω into two wedges as in Figure 5.1. Then for every $f \in H^1(\Omega)$ with $f|_{\Sigma} = 0$ and all $\gamma > 0$ the estimate*

$$\|\nabla f\|_{L^2(\Omega; \mathbb{C}^2)}^2 - \gamma \|f|_{\partial\Omega}\|_{L^2(\partial\Omega)}^2 \geq -\gamma^2 \|f\|_{L^2(\Omega)}^2$$

holds.

Proof. Let $f \in H^1(\Omega)$ with $f|_{\Sigma} = 0$ and denote the restrictions of f to the wedges Ω_1 and Ω_2 by f_1 and f_2 , respectively. Extend the wedge Ω_1 with degree $\varphi_1 < \varphi$ to the half-plane \mathbb{R}_+^2 by gluing the wedge Ω'_1 with the angle $\pi - \varphi_1$ as in Figure 5.2.




FIGURE 5.2. Extension of the wedge Ω_1 to the half-plane \mathbb{R}_+^2 via the wedge Ω'_1 .

As $f_1 \in H^1(\Omega_1)$ and $f_1|_{\Sigma} = 0$ we can extend f_1 by zero to $\tilde{f} \in H^1(\mathbb{R}_+^2)$. Then for $\gamma > 0$

$$\begin{aligned} \|\nabla f_1\|_{L^2(\Omega_1; \mathbb{C}^2)}^2 - \gamma \|f_1|_{\Sigma_1}\|_{L^2(\Sigma_1)}^2 &= \|\nabla \tilde{f}\|_{L^2(\mathbb{R}_+^2; \mathbb{C}^2)}^2 - \gamma \|\tilde{f}\|_{\partial \mathbb{R}_+^2}^2 \\ &\geq -\gamma^2 \|\tilde{f}\|_{L^2(\mathbb{R}_+^2)}^2 = -\gamma^2 \|f_1\|_{L^2(\Omega_1)}^2 \end{aligned}$$

holds by Lemma 5.1. The same argument shows that for $\gamma > 0$ the function $f_2 \in H^1(\Omega_2)$ satisfies

$$\|\nabla f_2\|_{L^2(\Omega_2; \mathbb{C}^2)}^2 - \gamma \|f_2|_{\Sigma_2}\|_{L^2(\Sigma_2)}^2 \geq -\gamma^2 \|f_2\|_{L^2(\Omega_2)}^2.$$


Summing up the above estimates we obtain the estimate in the lemma. \square

It turns out to be useful in the proof of (3.11) to decompose functions in $H^1(\Omega)$ as sums of even and odd functions with respect to the angle bisector of the wedge Ω .

Lemma 5.3. *Let $\Omega \subset \mathbb{R}^2$ be a wedge with angle $\varphi \in (0, \pi]$ and let Σ be the angle bisector which separates Ω into two wedges with angles $\varphi/2 \in (0, \pi/2]$. Then every $f \in H^1(\Omega)$ can be decomposed into the sum $f_o + f_e$ such that the following conditions (a)-(e) hold:*

- (a) $f_o, f_e \in H^1(\Omega)$;
- (b) $(f_o, f_e)_{L^2(\Omega)} = 0$;
- (c) $(\nabla f_o, \nabla f_e)_{L^2(\Omega; \mathbb{C}^2)} = 0$;
- (d) $f_e|_{\Sigma_1} = f_e|_{\Sigma_2}$ and $f_o|_{\Sigma_1} = -f_o|_{\Sigma_2}$;
- (e) $f_o|_{\Sigma} = 0$.

Proof. Fix the Cartesian coordinate system such that the vertex of the wedge Ω is the origin and the angle bisector is the ordinate axis as in Figure 5.3

FIGURE 5.3. The wedge Ω in the Cartesian coordinate system.

Let $C_0^\infty(\bar{\Omega}) := \{f|_\Omega : f \in C_0^\infty(\mathbb{R}^2)\}$ and note that for $h_1, h_2 \in C_0^\infty(\bar{\Omega})$ with

$$h_1(x, y) = h_1(-x, y) \quad \text{and} \quad h_2(x, y) = -h_2(-x, y)$$

the equality

$$(5.1) \quad (h_1, h_2)_{L^2(\Omega)} = 0$$

holds. Let us introduce the mappings

$$P_{\text{even},0}: C_0^\infty(\bar{\Omega}) \rightarrow C_0^\infty(\bar{\Omega}), \quad (P_{\text{even},0}f)(x, y) := \frac{f(x, y) + f(-x, y)}{2},$$

$$P_{\text{odd},0}: C_0^\infty(\bar{\Omega}) \rightarrow C_0^\infty(\bar{\Omega}), \quad (P_{\text{odd},0}f)(x, y) := \frac{f(x, y) - f(-x, y)}{2},$$

and define $f_e := P_{\text{even},0}f$ and $f_o := P_{\text{odd},0}f$. Then obviously, $f = f_e + f_o$, and f_e and f_o satisfy (d) and (e) by their definition. Condition (b) holds according to (5.1). Computing partial derivatives we get

$$\partial_1 f_e = \frac{(\partial_1 f)(x, y) - (\partial_1 f)(-x, y)}{2}, \quad \partial_1 f_o = \frac{(\partial_1 f)(x, y) + (\partial_1 f)(-x, y)}{2},$$

$$\partial_2 f_e = \frac{(\partial_2 f)(x, y) + (\partial_2 f)(-x, y)}{2}, \quad \partial_2 f_o = \frac{(\partial_2 f)(x, y) - (\partial_2 f)(-x, y)}{2}.$$

Hence, according to (5.1) we conclude

$$(\partial_1 f_e, \partial_1 f_o)_{L^2(\Omega)} = 0 \quad \text{and} \quad (\partial_2 f_e, \partial_2 f_o)_{L^2(\Omega)} = 0.$$

Therefore, condition (c) holds as well. Note that

$$\|P_{\text{even},0}f\|_{H^1(\Omega)} \leq \|f\|_{H^1(\Omega)} \quad \text{and} \quad \|P_{\text{odd},0}f\|_{H^1(\Omega)} \leq \|f\|_{H^1(\Omega)},$$

where $\|f\|_{H^1(\Omega)}^2 = \|f\|_{L^2(\Omega)}^2 + \|\nabla f\|_{L^2(\Omega; \mathbb{C}^2)}^2$. Hence, the operators $P_{\text{even},0}$ and $P_{\text{odd},0}$ can be extended by continuity to operators $P_{\text{even}}: H^1(\Omega) \rightarrow H^1(\Omega)$ and $P_{\text{odd}}: H^1(\Omega) \rightarrow H^1(\Omega)$ with $\text{dom } P_{\text{even}} = \text{dom } P_{\text{odd}} = H^1(\Omega)$. For $f \in H^1(\Omega)$ define $f_e := P_{\text{even}}f$ and $f_o := P_{\text{odd}}f$. Clearly, $f = f_e + f_o$ holds, and the conditions (a), (b), (c), (d) and (e) are satisfied. This completes the proof of the lemma. \square

5.2. Proof of (3.10). The assertion $\min \sigma(-\Delta_{\delta, \alpha}) = -\frac{\alpha^2}{3}$ is essentially a consequence of Lemma 5.1. In fact, recall first that the operator $-\Delta_{\delta, \alpha}$ corresponds to the quadratic form

$$\mathfrak{a}_{\delta, \alpha}[f] = \|\nabla f\|_{L^2(\mathbb{R}^2; \mathbb{C}^2)}^2 - \alpha \|f|_\Sigma\|_{L^2(\Sigma)}^2, \quad \text{dom } \mathfrak{a}_{\delta, \alpha} = H^1(\mathbb{R}^2).$$

Let $f \in \text{dom } \mathfrak{a}_{\delta, \alpha}$ such that $\|f\|_{L^2} = 1$, and denote the restrictions by $f_k := f|_{\Omega_k}$, $k = 1, 2, 3$. From Lemma 5.1 with $\gamma = \alpha/2$ and $\varphi = 2\pi/3$ we obtain the estimates

$$(5.2) \quad \|\nabla f_k\|_{L^2(\Omega_k; \mathbb{C}^2)}^2 - \frac{\alpha}{2} \|f_k|_{\partial\Omega_k}\|_{L^2(\partial\Omega_k)}^2 \geq -\frac{\alpha^2}{3} \|f_k\|_{L^2(\Omega_k)}^2$$

for $k = 1, 2, 3$. Since

$$\mathfrak{a}_{\delta, \alpha}[f] = \sum_{k=1}^3 \|\nabla f_k\|_{L^2(\Omega_k; \mathbb{C}^2)}^2 - \frac{\alpha}{2} \sum_{k=1}^3 \|f_k|_{\partial\Omega_k}\|_{L^2(\partial\Omega_k)}^2$$

and $\sum_{k=1}^3 \|f_k\|_{L^2(\Omega_k)}^2 = 1$ we conclude from (5.2) that

$$\mathfrak{a}_{\delta, \alpha}[f] \geq -\frac{\alpha^2}{3},$$

and hence $\min \sigma(-\Delta_{\delta, \alpha}) \geq -\frac{\alpha^2}{3}$. Furthermore, according to Lemma 5.1 there exist $f_k \in H^1(\Omega_k)$ such that equality holds in (5.2). This yields

$$\min \sigma(-\Delta_{\delta, \alpha}) = -\frac{\alpha^2}{3}$$

and completes the proof of (3.10). We remark that the estimate $\min \sigma(-\Delta_{\delta, \alpha}) \leq -\frac{\alpha^2}{3}$ follows also from [BEW09, Theorem 3.2].

5.3. Proof of (3.11). The proof of the estimate

$$\min(-\Delta_{\delta', \beta}) \geq -\left(\frac{12\sqrt{3}-2}{9}\right)^2 \frac{1}{\beta^2}$$

is carried out in three steps, followed by a separate proof of the inequality (5.3) below.

Step I. Recall first that $-\Delta_{\delta', \beta}$ corresponds to the quadratic form

$$\begin{aligned} \mathfrak{a}_{\delta', \beta}[f] &= \sum_{k=1}^3 \|\nabla f_k\|_{L^2(\Omega_k; \mathbb{C}^2)}^2 - \beta^{-1} \|f_1|_{\Sigma_{12}} - f_2|_{\Sigma_{12}}\|_{L^2(\Sigma_{12})}^2 \\ &\quad - \beta^{-1} \|f_2|_{\Sigma_{23}} - f_3|_{\Sigma_{23}}\|_{L^2(\Sigma_{23})}^2 - \beta^{-1} \|f_3|_{\Sigma_{13}} - f_1|_{\Sigma_{13}}\|_{L^2(\Sigma_{13})}^2, \end{aligned}$$

$$\text{dom } \mathfrak{a}_{\delta', \beta} = \bigoplus_{k=1}^3 H^1(\Omega_k),$$

where $f_k = f|_{\Omega_k}$, $k = 1, 2, 3$. We split the problem into two separate problems for odd and even components. For $f \in \text{dom } \mathfrak{a}_{\delta', \beta}$ with $\|f\|_{L^2(\mathbb{R}^2)} = 1$ we decompose the restrictions $f_k = f|_{\Omega_k}$ as in Lemma 5.3. Let $\{f_{k,e}\}_{k=1}^3$ and $\{f_{k,o}\}_{k=1}^3$ be the corresponding even and odd components, and let

$$\begin{aligned} \theta_1 &:= f_{1,e}|_{\Sigma_{12}} = f_{1,e}|_{\Sigma_{13}}, & \eta_1 &:= f_{1,o}|_{\Sigma_{12}} = f_{1,o}|_{\Sigma_{13}}, \\ \theta_2 &:= f_{2,e}|_{\Sigma_{23}} = f_{1,e}|_{\Sigma_{12}}, & \eta_2 &:= f_{1,o}|_{\Sigma_{23}} = f_{1,o}|_{\Sigma_{12}}, \\ \theta_3 &:= f_{3,e}|_{\Sigma_{13}} = f_{1,e}|_{\Sigma_{23}}, & \eta_3 &:= f_{1,o}|_{\Sigma_{13}} = f_{1,o}|_{\Sigma_{23}}. \end{aligned}$$

Using Lemma 5.3 we obtain

$$\mathfrak{a}_{\delta', \beta}[f] = \sum_{k=1}^3 \|\nabla f_{k,e}\|_{L^2(\Omega_k; \mathbb{C}^2)}^2 + \sum_{k=1}^3 \|\nabla f_{k,o}\|_{L^2(\Omega_k; \mathbb{C}^2)}^2 - \frac{1}{\beta} S$$

where the value S is given by

$$\|\theta_1 - \theta_2 + \eta_1 + \eta_2\|_{L^2(\mathbb{R}_+)}^2 + \|\theta_2 - \theta_3 + \eta_2 + \eta_3\|_{L^2(\mathbb{R}_+)}^2 + \|\theta_3 - \theta_1 + \eta_3 + \eta_1\|_{L^2(\mathbb{R}_+)}^2;$$

here we have identified $L^2(\Sigma_{ij}) = L^2(\mathbb{R}_+)$, $i, j = 1, 2, 3$. We shall show later that the above term can be estimated by

$$(5.3) \quad S \leq (4 - \omega(1 - t)) \sum_{k=1}^3 \|\theta_k\|_{L^2(\mathbb{R}_+)}^2 + (4 + 3\omega t^{-1}) \sum_{k=1}^3 \|\eta_k\|_{L^2(\mathbb{R}_+)}^2$$

for all $t > 0$ and all $\omega \in [0, 1]$. With the help of this inequality we find

$$(5.4) \quad \mathfrak{a}_{\delta', \beta}[f] \geq C_e + C_o,$$

where C_e and C_o are given by

$$C_e := \sum_{k=1}^3 \|\nabla f_{k,e}\|_{L^2(\Omega_k; \mathbb{C}^2)}^2 - \frac{1}{\beta} (4 - \omega(1 - t)) \sum_{k=1}^3 \|\theta_k\|_{L^2(\mathbb{R}_+)}^2$$

and

$$C_o := \sum_{k=1}^3 \|\nabla f_{k,o}\|_{L^2(\Omega_k; \mathbb{C}^2)}^2 - \frac{1}{\beta} (4 + 3\omega t^{-1}) \sum_{k=1}^3 \|\eta_k\|_{L^2(\mathbb{R}_+)}^2.$$

Step II. In this step we estimate C_e and C_o . We start with C_e . Applying Lemma 5.1 with $\gamma = \frac{1}{2\beta}(4 - \omega(1 - t))$ and $\varphi = 2\pi/3$ to the functions $\{f_{k,e}\}_{k=1}^3$ we get

$$\|\nabla f_{k,e}\|_{L^2(\Omega_k; \mathbb{C}^2)}^2 - \frac{1}{\beta} (4 - \omega(1 - t)) \|\theta_k\|_{L^2(\mathbb{R}_+)}^2 \geq -\frac{1}{3\beta^2} (4 - \omega(1 - t))^2 \|f_{k,e}\|_{L^2(\Omega_k)}^2$$

for $k = 1, 2, 3$. Summing up these three inequalities we find

$$(5.5) \quad C_e \geq -\frac{1}{3\beta^2} (4 - \omega(1 - t))^2 \sum_{k=1}^3 \|f_{k,e}\|_{L^2(\Omega_k)}^2$$

for all $t > 0$ and all $\omega \in [0, 1]$. Next we estimate C_o . Note that $f_{k,o}|_{\Sigma_k} = 0$ for $k = 1, 2, 3$, and hence we can apply Lemma 5.2 with $\gamma = \frac{1}{2\beta}(4 + 3\omega t^{-1})$ to the functions $\{f_{k,o}\}_{k=1}^3$. This yields

$$\|\nabla f_{k,o}\|_{L^2(\Omega_k; \mathbb{C}^2)}^2 - \frac{1}{\beta} (4 + 3\omega t^{-1}) \|\eta_k\|_{L^2(\mathbb{R}_+)}^2 \geq -\frac{1}{4\beta^2} (4 + 3\omega t^{-1})^2 \|f_{k,o}\|_{L^2(\Omega_k)}^2$$

for $k = 1, 2, 3$. Summing up these three inequalities gives

$$(5.6) \quad C_o \geq -\frac{1}{4\beta^2} (4 + 3\omega t^{-1})^2 \sum_{k=1}^3 \|f_{k,o}\|_{L^2(\Omega_k)}^2$$

for all $t > 0$ and all $\omega \in [0, 1]$.

Step III. Note that

$$\sum_{k=1}^3 (\|f_{k,e}\|_{L^2(\Omega_k)}^2 + \|f_{k,o}\|_{L^2(\Omega_k)}^2) = \|f\|_{L^2(\mathbb{R}^2)}^2 = 1.$$

Thus, (5.4), (5.5) and (5.6) imply

$$(5.7) \quad \mathfrak{a}_{\delta', \beta}[f] \geq -\inf_{t>0} \min_{\omega \in [0, 1]} \max \left\{ \frac{1}{3} (4 - \omega(1 - t))^2, \frac{1}{4} \left(4 + \frac{3\omega}{t}\right)^2 \right\} \frac{1}{\beta^2}$$

and for (3.11) to hold it remains to show that the value on the right hand side is equal to $-\left(\frac{12\sqrt{3}-2}{9}\right)^2 \frac{1}{\beta^2}$. For this consider the functions

$$M_1(\omega, t) := \frac{1}{3}(4 - \omega(1-t))^2 \quad \text{and} \quad M_2(\omega, t) := \frac{1}{4}\left(4 + \frac{3\omega}{t}\right)^2.$$

We have

$$M_1(0, 1) = \frac{16}{3}, \quad M_2(0, 1) = 4, \quad \text{and} \quad M_1(\omega, t) \geq \frac{16}{3}, \quad t \geq 1, \omega \in [0, 1],$$

and therefore

$$(5.8) \quad \min_{t \in [1, +\infty)} \min_{\omega \in [0, 1]} \max\{M_1(\omega, t), M_2(\omega, t)\} = \frac{16}{3}.$$

Suppose now that $t \in (0, 1)$ is fixed. Then $M_1(\cdot, t)$ is continuous and decreasing, whereas $M_2(\cdot, t)$ is continuous and increasing, and a straightforward computation shows that for

$$\omega_* := \frac{(8 - 4\sqrt{3})t}{3\sqrt{3} + 2(1-t)t} \in [0, 1]$$

we have $M_1(\omega_*, t) = M_2(\omega_*, t)$. Hence for $t \in (0, 1)$ fixed we find

$$\begin{aligned} \min_{\omega \in [0, 1]} \max\{M_1(\omega, t), M_2(\omega, t)\} &= M_1(\omega_*, t) = M_2(\omega_*, t) \\ &= \frac{1}{4}\left(4 + \frac{3(8 - 4\sqrt{3})}{3\sqrt{3} + 2(1-t)t}\right)^2. \end{aligned}$$

Next, we minimize with respect to $t \in (0, 1)$. Clearly, the above value is minimal in $t \in (0, 1)$ if $t = 1/2$. Therefore we obtain

$$(5.9) \quad \min_{t \in (0, 1)} \min_{\omega \in [0, 1]} \max\{M_1(\omega, t), M_2(\omega, t)\} = M_2(\omega_*, 1/2) = \left(\frac{12\sqrt{3}-2}{9}\right)^2.$$

Now (5.8) and (5.9) together imply

$$\min_{t > 0} \min_{\omega \in [0, 1]} \max\{M_1(\omega, t), M_2(\omega, t)\} = \min\left\{\frac{16}{3}, \left(\frac{12\sqrt{3}-2}{9}\right)^2\right\} = \left(\frac{12\sqrt{3}-2}{9}\right)^2,$$

and hence the assertion (3.11) on the minimum of the spectrum of $-\Delta_{\delta', \beta}$ follows from (5.7).

Proof of the estimate (5.3). Let $\theta_1, \theta_2, \theta_3 \in L^2(\mathbb{R}_+)$ and $\eta_1, \eta_2, \eta_3 \in L^2(\mathbb{R}_+)$. We shall not use an index for the norm in $L^2(\mathbb{R}_+)$ in this proof. From

$$\begin{aligned} \|\theta_1 - \theta_2 + \eta_1 + \eta_2\|^2 &\leq 2\|\theta_1 + \eta_1\|^2 + 2\|\theta_2 - \eta_2\|^2, \\ \|\theta_2 - \theta_3 + \eta_2 + \eta_3\|^2 &\leq 2\|\theta_2 + \eta_2\|^2 + 2\|\theta_3 - \eta_3\|^2, \\ \|\theta_3 - \theta_1 + \eta_3 + \eta_1\|^2 &\leq 2\|\theta_3 + \eta_3\|^2 + 2\|\theta_1 - \eta_1\|^2, \end{aligned}$$

and the parallelogram identity we conclude

$$(5.10) \quad S \leq 2 \sum_{k=1}^3 (\|\theta_k - \eta_k\|^2 + \|\theta_k + \eta_k\|^2) \leq 4 \sum_{k=1}^3 (\|\theta_k\|^2 + \|\eta_k\|^2).$$

On the other hand we have

$$\begin{aligned}
(5.11) \quad S &= \|\theta_1 - \theta_2\|^2 + \|\theta_2 - \theta_3\|^2 + \|\theta_3 - \theta_1\|^2 + \|\eta_1 + \eta_2\|^2 + \|\eta_2 + \eta_3\|^2 + \|\eta_3 + \eta_1\|^2 \\
&\quad + 2\operatorname{Re}[(\theta_1 - \theta_2, \eta_1 + \eta_2)] + 2\operatorname{Re}[(\theta_2 - \theta_3, \eta_2 + \eta_3)] + 2\operatorname{Re}[(\theta_3 - \theta_1, \eta_3 + \eta_1)] \\
&= \|\theta_1 - \theta_2\|^2 + \|\theta_2 - \theta_3\|^2 + \|\theta_3 - \theta_1\|^2 + \|\eta_1 + \eta_2\|^2 + \|\eta_2 + \eta_3\|^2 + \|\eta_3 + \eta_1\|^2 \\
&\quad + 2\operatorname{Re}[(\theta_1, \eta_2 - \eta_3)] + 2\operatorname{Re}[(\theta_2, \eta_3 - \eta_1)] + 2\operatorname{Re}[(\theta_3, \eta_1 - \eta_2)]
\end{aligned}$$

and the Cauchy-Schwarz inequality together with the inequality $2ab \leq ta^2 + \frac{1}{t}b^2$, $a, b > 0$, $t > 0$, yields

$$\begin{aligned}
\left| 2\operatorname{Re}[(\theta_1, \eta_2 - \eta_3)] \right| &\leq 2\|\theta_1\| \cdot \|\eta_2 - \eta_3\| \leq t\|\theta_1\|^2 + \frac{1}{t}\|\eta_2 - \eta_3\|^2, \\
\left| 2\operatorname{Re}[(\theta_2, \eta_3 - \eta_1)] \right| &\leq 2\|\theta_2\| \cdot \|\eta_3 - \eta_1\| \leq t\|\theta_2\|^2 + \frac{1}{t}\|\eta_3 - \eta_1\|^2, \\
\left| 2\operatorname{Re}[(\theta_3, \eta_1 - \eta_2)] \right| &\leq 2\|\theta_3\| \cdot \|\eta_1 - \eta_2\| \leq t\|\theta_3\|^2 + \frac{1}{t}\|\eta_1 - \eta_2\|^2.
\end{aligned}$$

Combining the latter with (5.11) and making use of $\|\eta_i + \eta_j\|^2 \leq 2\|\eta_i\|^2 + 2\|\eta_j\|^2$, $i, j = 1, 2, 3$, we arrive at

$$\begin{aligned}
(5.12) \quad S &\leq \|\theta_1 - \theta_2\|^2 + \|\theta_2 - \theta_3\|^2 + \|\theta_3 - \theta_1\|^2 + 4\|\eta_1\|^2 + 4\|\eta_2\|^2 + 4\|\eta_3\|^2 \\
&\quad + t(\|\theta_1\|^2 + \|\theta_2\|^2 + \|\theta_3\|^2) + \frac{1}{t}(\|\eta_1 - \eta_2\|^2 + \|\eta_2 - \eta_3\|^2 + \|\eta_3 - \eta_1\|^2)
\end{aligned}$$

for all $t > 0$. Moreover, as

$$\begin{aligned}
\|\theta_1 - \theta_2\|^2 + \|\theta_2 - \theta_3\|^2 + \|\theta_3 - \theta_1\|^2 &= 3\|\theta_1\|^2 + 3\|\theta_2\|^2 + 3\|\theta_3\|^2 - \|\theta_1 + \theta_2 + \theta_3\|^2 \\
&\leq 3\|\theta_1\|^2 + 3\|\theta_2\|^2 + 3\|\theta_3\|^2,
\end{aligned}$$

and analogously

$$\|\eta_1 - \eta_2\|^2 + \|\eta_2 - \eta_3\|^2 + \|\eta_3 - \eta_1\|^2 \leq 3\|\eta_1\|^2 + 3\|\eta_2\|^2 + 3\|\eta_3\|^2,$$

we can further estimate (5.12) by

$$(5.13) \quad S \leq (3+t) \sum_{k=1}^3 \|\theta_k\|^2 + \sum_{k=1}^3 \left(4 + \frac{3}{t}\right) \|\eta_k\|^2.$$

In order to obtain the estimate (5.3) let $\omega \in [0, 1]$, consider $S = \omega S + (1-\omega)S$ and estimate ωS as in (5.13) and $(1-\omega)S$ as in (5.10). \square

REFERENCES

- [AGHH] S. Albeverio, F. Gesztesy, R. Hoegh-Krohn, and H. Holden, *Solvable models in quantum mechanics*, 2nd edition, AMS Chelsea Publishing, Providence, 2005.
- [AKMN13] S. Albeverio, A. Kostenko, M. M. Malamud, and H. Neidhardt, Spherical Schrödinger operators with δ -type interactions, *J. Math. Phys.* **54** (2013), 052103.
- [AGS87] J.-P. Antoine, F. Gesztesy and J. Shabani, Exactly solvable models of sphere interactions in quantum mechanics, *J. Phys. A* **20** (1987), 3687–3712.
- [AH77] K. Appel and W. Haken, W. Every planar map is four colorable. I. Discharging, *Illinois J. Math.* **21** (1977), 429–490.
- [AHK77] K. Appel, K. W. Haken, and J. Koch, Every planar map is four colorable. II. Reducibility. *Illinois J. Math.* **21** (1977), 491–567.
- [BEL13] J. Behrndt, P. Exner, and V. Lotoreichik, Essential spectrum of Schrödinger operators with δ -interactions on the union of compact Lipschitz hypersurfaces, *to appear in Proc. Appl. Math. Mech.* **13** (2013)
- [BL07] J. Behrndt and M. Langer, Boundary value problems for elliptic partial differential operators on bounded domains, *J. Funct. Anal.* **243** (2007), 536–565.

- [BLL12a] J. Behrndt, M. Langer and V. Lotoreichik, Spectral estimates for resolvent differences of self-adjoint elliptic operators, *to appear in Integral Equations Operator Theory*.
- [BLL13] J. Behrndt, M. Langer, and V. Lotoreichik, Schrödinger operators with δ and δ' -potentials supported on hypersurfaces, *Ann. Henri Poincaré*, **14** (2013), 385–423.
- [B62] M. Sh. Birman, Perturbations of the continuous spectrum of a singular elliptic operator by varying the boundary and the boundary conditions, *Vestnik Leningrad. Univ.* **17** (1962), 22–55 (in Russian); translated in: *Amer. Math. Soc. Transl.* **225** (2008), 19–53.
- [BS87] M. Sh. Birman and M. Z. Solomjak, *Spectral theory of selfadjoint operators in Hilbert spaces*, Dordrecht, Holland, 1987.
- [BEH08] J. Blank, P. Exner, and M. Havlíček, *Hilbert space operators in quantum physics*, 2nd edition, Springer, New York, 2008.
- [BEKS94] J. F. Brasche, P. Exner, Yu. A. Kuperin and P. Šeba, Schrödinger operators with singular interactions, *J. Math. Anal. Appl.* **184** (1994), 112–139.
- [BN11] J. F. Brasche and L. Nizhnik, One-dimensional Schrödinger operators with δ' -interactions on a set of Lebesgue measure zero, arXiv:1112.2545.
- [BEW09] B. M. Brown, M. S. P. Eastham and I. G. Wood, Estimates for the lowest eigenvalue of a star graph, *J. Math. Anal. Appl.* **354** (2009), 24–30.
- [dC] M. P. do Carmo, *Differential geometry of curves and surfaces*, Englewood Cliffs, N.J., 1976.
- [CK11] J. Čislo and S. Kondej, Upper bound for the number of bound states induced by the curvature of singular potential, *Rep. Math. Phys.* **68** (2011), 225–240.
- [EE] D. Edmunds and W. Evans, *Spectral theory and differential operators*, Clarendon Press, New York, 1987.
- [E08] P. Exner, Leaky quantum graphs: a review, in: *Analysis on graphs and its applications*. Selected papers based on the Isaac Newton Institute for Mathematical Sciences programme, Cambridge, UK, 2007. *Proc. Symp. Pure Math.* **77** (2008), 523–564.
- [EF09] P. Exner and M. Fraas, On geometric perturbations of critical Schrödinger operators with a surface interaction, *J. Math. Phys.* **50** (2009), 112101.
- [EJ13] P. Exner and M. Jex, Spectral asymptotics of a strong δ' interaction on a planar loop, arXiv: 1304.7696.
- [EI01] P. Exner and I. Ichinose, Geometrically induced spectrum in curved leaky wires, *J. Phys. A* **34** (2001), 1439–1450.
- [EK03] P. Exner and S. Kondej, Bound states due to a strong δ interaction supported by a curved surface, *J. Phys. A* **36** (2003), 443–457.
- [EK05] P. Exner and S. Kondej, Scattering by local deformations of a straight leaky wire, *J. Phys. A* **38** (2005), 4865–4874.
- [EK08] P. Exner and S. Kondej, Hiatus perturbation for a singular Schrödinger operator with an interaction supported by a curve in \mathbb{R}^3 , *J. Math. Phys.* **49** (2008), 032111.
- [EN03] P. Exner and K. Němcová, Leaky quantum graphs: approximations by point-interaction Hamiltonians, *J. Phys. A* **36** (2003), 10173–10193.
- [EP12] P. Exner and K. Pankrashkin, Strong coupling asymptotics for a singular Schroedinger operator with an interaction supported by an open arc, arXiv:1207.2271
- [ET04] P. Exner and M. Tater, Spectra of soft ring graphs. Special section on quantum graphs. *Waves Random Media* **14** (2004), 47–60.
- [EY02] P. Exner and K. Yoshitomi, Asymptotics of eigenvalues of the Schrödinger operator with a strong δ -interaction on a loop, *J. Geom. Phys.* **41** (2002), 344–358.
- [HT] D. Haroske and H. Triebel, *Distributions, Sobolev spaces, elliptic equations*, European Mathematical Society (EMS), Zürich, 2008.
- [H89] J. Herczyński, On Schrödinger operators with distributional potentials, *J. Operator Theory* **21** (1989), 273–295.
- [K] T. Kato, *Perturbation theory for linear operators*, Springer-Verlag, Berlin, 1995.
- [K12] S. Kondej, Resonances induced by broken symmetry in a system with a singular potential, *Ann. Henri Poincaré* **13** (2012), 1451–1467.
- [KV07] S. Kondej and I. Veselić, Lower bounds on the lowest spectral gap of singular potential Hamiltonians, *Ann. Henri Poincaré* **8** (2007), 109–134.
- [KM10] A. Kostenko and M. M. Malamud, 1-D Schrödinger operators with local point interactions on a discrete set, *J. Differential Equations* **249** (2010), 253–304.
- [LP08] M. Levitin and L. Parnovski, On the principal eigenvalue of a Robin problem with a large parameter, *Math. Nachr.* **281** (2008), 272–281.

- [LLP10] I. Lobanov, V. Lotoreichik, and I. Yu. Popov, Lower bound on the spectrum of the Schrödinger operator in the plane with delta-potential supported by a curve, *Theor. Math. Phys.* **162** (2010), 332–340.
- [MS12] M. M. Malamud and K. Schmüdgen, Spectral theory of Schrödinger operators with infinitely many point interactions and radial positive definite functions, *J. Funct. Anal.* **263** (2012), 3144–3194
- [M87] J. Marschall, The trace of Sobolev-Slobodeckij spaces on Lipschitz domains, *Manuscripta Math.* **58** (1987), 47–65.
- [McL] W. McLean, *Strongly elliptic systems and boundary integral equations*, Cambridge University Press, Cambridge, 2000.
- [MT01] B. Mohar and C. Thomassen, *Graphs on surfaces*, Johns Hopkins University Press, Baltimore, 2001.
- [Ne] J. Nečas, *Les méthodes directes en théorie des équations elliptiques*, Masson et Cie, Éditeurs, Paris, 1967.
- [O10] O. Ogurisu, On the number of negative eigenvalues of a multi-dimensional Schrödinger operator with point interactions. *Methods Funct. Anal. Topology* **16** (2010), 383–392.
- [RS-IV] M. Reed and B. Simon, *Methods of modern mathematical physics IV. Analysis of operators*, Academic Press, 1978.
- [S88] J. Shabani, Some properties of the Hamiltonian describing a finite number of δ' -interactions with support on concentric spheres, *Nuovo Cimento B (11)* **101** (1988), 429–437.
- [St] E. Stein, *Singular integrals and differentiability properties of functions*, Princeton University Press, Princeton, 1970.
- [W00] J. Weidmann, *Lineare Operatoren in Hilberträumen. Teil I: Grundlagen*, Teubner, Stuttgart, 2000 (German).